

Package Insert

ACCORD BORTEZOMIB 3.5 (Bortezomib Powder for Solution for Injection 3.5 mg/Vial)

- **Name and strength of active ingredient**

Bortezomib 3.5 mg/Vial

- **Product Description**

ACCORD BORTEZOMIB 3.5: A white to off-white cake or powder in a clear glass vial. The reconstituted solution is clear and colourless.

- **Pharmacodynamics & Pharmacokinetics**

Pharmacodynamic properties

Pharmacotherapeutic group: Antineoplastic agents, other antineoplastic agents, ATC code: L01XX32.

Mechanism of action

Bortezomib is a proteasome inhibitor. It is specifically designed to inhibit the chymotrypsin-like activity of the 26S proteasome in mammalian cells. The 26S proteasome is a large protein complex that degrades ubiquitinated proteins. The ubiquitin-proteasome pathway plays an essential role in regulating the turnover of specific proteins, thereby maintaining homeostasis within cells. Inhibition of the 26S proteasome prevents this targeted proteolysis and affects multiple signalling cascades within the cell, ultimately resulting in cancer cell death.

Bortezomib is highly selective for the proteasome. At 10 µM concentrations, bortezomib does not inhibit any of a wide variety of receptors and proteases screened and is more than 1,500-fold more selective for the proteasome than for its next preferable enzyme. The kinetics of proteasome inhibition were evaluated *in vitro*, and bortezomib was shown to dissociate from the proteasome with a $t_{1/2}$ of 20 minutes, thus demonstrating that proteasome inhibition by bortezomib is reversible.

Bortezomib mediated proteasome inhibition affects cancer cells in a number of ways, including, but not limited to, altering regulatory proteins, which control cell cycle progression and nuclear factor kappa B (NF-κB) activation. Inhibition of the proteasome results in cell cycle arrest and apoptosis. NF-κB is a transcription factor whose activation is required for many aspects of tumourigenesis, including cell growth and survival, angiogenesis, cell-cell interactions, and metastasis. In myeloma, bortezomib affects the ability of myeloma cells to interact with the bone marrow microenvironment.

Experiments have demonstrated that bortezomib is cytotoxic to a variety of cancer cell types and that cancer cells are more sensitive to the pro-apoptotic effects of proteasome inhibition than normal cells. Bortezomib causes reduction of tumour growth *in vivo* in many preclinical tumour models, including multiple myeloma.

Data from *in vitro*, *ex-vivo*, and animal models with bortezomib suggest that it increases osteoblast differentiation and activity and inhibits osteoclast function. These effects have been observed in patients with multiple myeloma affected by an advanced osteolytic disease and treated with bortezomib.

Pharmacokinetic properties

Absorption

Following intravenous bolus administration of a 1.0 mg/m² and 1.3 mg/m² dose to 11 patients with multiple myeloma and creatinine clearance values greater than 50 ml/min, the mean first-dose maximum plasma concentrations of bortezomib were 57 and 112 ng/ml, respectively. In subsequent doses, mean maximum observed plasma concentrations ranged from 67 to 106 ng/ml for the 1.0 mg/m² dose and 89 to 120 ng/ml for the 1.3 mg/m² dose.

Following an intravenous bolus or subcutaneous injection of a 1.3 mg/m² dose to patients with multiple myeloma (n=14 in the intravenous group, n=17 in the subcutaneous group), the total systemic exposure after repeat dose administration (AUC_{last}) was equivalent for subcutaneous and intravenous administrations. The C_{max} after subcutaneous administration (20.4 ng/ml) was lower than intravenous (223 ng/ml). The AUC_{last} geometric mean ratio was 0.99 and 90% confidence intervals were 80.18%-122.80%.

Distribution

The mean distribution volume (V_d) of bortezomib ranged from 1,659 l to 3,294 l following single- or repeated-dose intravenous administration of 1.0 mg/m² or 1.3 mg/m² to patients with multiple myeloma. This suggests that bortezomib distributes widely to peripheral tissues. Over a bortezomib concentration range of 0.01 to 1.0 µg/ml, the *in vitro* protein binding averaged 82.9% in human plasma. The fraction of bortezomib bound to plasma proteins was not concentration-dependent.

Biotransformation

In vitro studies with human liver microsomes and human cDNA-expressed cytochrome P450 isozymes indicate that bortezomib is primarily oxidatively metabolised via cytochrome P450 enzymes, 3A4, 2C19, and 1A2. The major metabolic pathway is deboronation to form two deboronated metabolites that subsequently undergo hydroxylation to several metabolites. Deboronated-bortezomib metabolites are inactive as 26S proteasome inhibitors.

Elimination

The mean elimination half-life (t_{1/2}) of bortezomib upon multiple dosing ranged from 40-193 hours. Bortezomib is eliminated more rapidly following the first dose compared to subsequent doses. Mean total body clearances were 102 and 112 l/h following the first dose for doses of 1.0 mg/m² and 1.3 mg/m², respectively, and ranged from 15 to 32 l/h and 18 to 32 l/h following subsequent doses for doses of 1.0 mg/m² and 1.3 mg/m², respectively.

Special populations

Hepatic impairment

The effect of hepatic impairment on the pharmacokinetics of bortezomib was assessed in a Phase I study during the first treatment cycle, including 61 patients primarily with solid tumors and varying degrees of hepatic impairment at bortezomib doses ranging from 0.5 to 1.3 mg/m².

When compared to patients with normal hepatic function, mild hepatic impairment did not alter dose-normalised bortezomib AUC. However, the dose-normalised mean AUC values were increased by approximately 60% in patients with moderate or severe hepatic impairment. A lower starting dose is recommended in patients with moderate or severe hepatic impairment, and those patients should be closely monitored.

Renal impairment

A pharmacokinetic study was conducted in patients with various degrees of renal impairment who were classified according to their creatinine clearance values (CrCL) into the following groups: Normal (CrCL ≥ 60 ml/min/1.73 m², n=12), Mild (CrCL=40-59 ml/min/1.73 m², n=10), Moderate (CrCL=20-39 ml/min/1.73 m², n=9), and Severe (CrCL < 20 ml/min/1.73 m², n=3). A group of dialysis patients who were dosed after dialysis was also included in the study (n=8). Patients were administered intravenous doses of 0.7 to 1.3 mg/m² of bortezomib twice weekly. Exposure of bortezomib (dose-normalised AUC and C_{max}) was comparable among all the groups.

• Indication

Bortezomib Accord is indicated:

- for the treatment of patients with multiple myeloma.
- for the treatment of patients with mantle cell lymphoma who have received at least one prior therapy.

• Recommended Dosage

Treatment must be initiated and administered under the supervision of a physician qualified and experienced in the use of chemotherapeutic agents. Bortezomib Accord must be reconstituted by a healthcare professional.

Posology for treatment of progressive multiple myeloma (patients who have received at least one prior therapy)

Monotherapy

Bortezomib Accord is administered via intravenous or subcutaneous injection at the recommended dose of 1.3 mg/m² body surface area twice weekly for two weeks on days 1, 4, 8, and 11 in a 21-day treatment cycle. This 3-week period is considered a treatment cycle. It is recommended that patients receive 2 cycles of bortezomib following a confirmation of a complete response. It is also recommended that responding patients who do not achieve a complete remission receive a total of 8 cycles of bortezomib therapy. At least 72 hours should elapse between consecutive doses of bortezomib.

Dose adjustments during treatment and re-initiation of treatment for monotherapy

Bortezomib treatment must be withheld at the onset of any Grade 3 non-haematological or any Grade 4 haematological toxicities, excluding neuropathy as discussed below.

Once the symptoms of the toxicity have resolved, bortezomib treatment may be re-initiated at a 25% reduced dose (1.3 mg/m² reduced to 1.0 mg/m²; 1.0 mg/m² reduced to 0.7 mg/m²). If the toxicity is not resolved or if it recurs at the lowest dose, discontinuation of bortezomib must be considered unless the benefit of treatment clearly outweighs the risk.

Neuropathic pain and/or peripheral neuropathy

Patients who experience bortezomib-related neuropathic pain and/or peripheral neuropathy are to be managed as presented in Table 1. Patients with pre-existing severe neuropathy may be treated with bortezomib only after careful risk/benefit assessment.

Table 1: Recommended posology modifications for Bortezomib Accord-related neuropathy*

Severity of neuropathy	Posology modification
Grade 1 (asymptomatic; loss of deep tendon reflexes or paresthesia) with no pain or loss of function	None
Grade 1 with pain or Grade 2 (moderate symptoms; limiting instrumental Activities of Daily Living	Reduce Bortezomib Accord to 1.0 mg/m ² or

(ADL)**)	Change Bortezomib Accord treatment schedule to 1.3 mg/m ² once per week
Grade 2 with pain or Grade 3 (severe symptoms; limiting self care ADL***)	Withhold Bortezomib Accord treatment until symptoms of toxicity have resolved. When toxicity resolves re-initiate Bortezomib Accord treatment and reduce dose to 0.7 mg/m ² once per week.
Grade 4 (life-threatening consequences; urgent intervention indicated) and/or severe autonomic neuropathy	Discontinue Bortezomib Accord

* Based on posology modifications in Phase II and III multiple myeloma studies and post-marketing experience. Grading based on NCI Common Toxicity Criteria CTCAE v 4.0.

** *Instrumental ADL*: refers to preparing meals, shopping for groceries or clothes, using telephone, managing money, etc;

*** *Self care ADL*: refers to bathing, dressing and undressing, feeding self, using the toilet, taking medicinal products, and not bedridden.

Combination therapy with pegylated liposomal doxorubicin

Bortezomib Accord is administered via intravenous or subcutaneous injection at the recommended dose of 1.3 mg/m² body surface area twice weekly for two weeks on days 1, 4, 8, and 11 in a 21-day treatment cycle. This 3-week period is considered a treatment cycle. At least 72 hours should elapse between consecutive doses of Bortezomib Accord.

Pegylated liposomal doxorubicin is administered at 30 mg/m² on day 4 of the Bortezomib Accord treatment cycle as a 1 hour intravenous infusion administered after the Bortezomib Accord injection.

Up to 8 cycles of this combination therapy can be administered as long as patients have not progressed and tolerate treatment. Patients achieving a complete response can continue treatment for at least 2 cycles after the first evidence of complete response, even if this requires treatment for more than 8 cycles. Patients whose levels of paraprotein continue to decrease after 8 cycles can also continue for as long as treatment is tolerated and they continue to respond.

For additional information concerning pegylated liposomal doxorubicin, see the corresponding Summary of Product Characteristics.

Combination with dexamethasone

Bortezomib Accord is administered via intravenous or subcutaneous injection at the recommended dose of 1.3 mg/m² body surface area twice weekly for two weeks on days 1, 4, 8, and 11 in a 21 day treatment cycle. This 3-week period is considered a treatment cycle. At least 72 hours should elapse between consecutive doses of Bortezomib Accord.

Dexamethasone is administered orally at 20 mg on days 1, 2, 4, 5, 8, 9, 11, and 12 of the Bortezomib Accord treatment cycle.

Patients achieving a response or a stable disease after 4 cycles of this combination therapy can continue to receive the same combination for a maximum of 4 additional cycles.

For additional information concerning dexamethasone, see the corresponding Summary of Product Characteristics.

Dose adjustments for combination therapy for patients with progressive multiple myeloma

For Bortezomib Accord dosage adjustments for combination therapy follow dose modification guidelines described under monotherapy above.

Posology for previously untreated multiple myeloma patients not eligible for haematopoietic stem cell transplantation

Combination therapy with melphalan and prednisone

Bortezomib Accord is administered via intravenous or subcutaneous injection in combination with oral melphalan and oral prednisone as shown in Table 2. A 6-week period is considered a treatment cycle. In Cycles 1-4, Bortezomib Accord is administered twice weekly on days 1, 4, 8, 11, 22, 25, 29 and 32. In Cycles 5-9, Bortezomib Accord is administered once weekly on days 1, 8, 22 and 29. At least 72 hours should elapse between consecutive doses of Bortezomib Accord.

Melphalan and prednisone should both be given orally on days 1, 2, 3 and 4 of the first week of each Bortezomib Accord treatment cycle.

Nine treatment cycles of this combination therapy are administered.

Table 2: Recommended posology for Bortezomib Accord in combination with melphalan and prednisone

Twice weekly Bortezomib Accord (cycles 1-4)												
Week	1				2		3	4		5		6
Bz (1.3 mg/m ²)	Day 1	--	--	Day 4	Day 8	Day 11	rest period	Day 22	Day 25	Day 29	Day 32	rest period
M (9 mg/m ²)	Day 1	Day 2	Day 3	Day 4	--	--	rest period	--	--	--	--	rest period
P (60 mg/m ²)	1	2	3	4			period					period

Once weekly Bortezomib Accord (cycles 5-9)												
Week	1				2		3	4		5		6
Bz (1.3 mg/m ²)	Day 1	--	--	--	Day 8		rest period	Day 22		Day 29		rest period
M (9 mg/m ²)	Day 1	Day 2	Day 3	Day 4	--		rest period	--				rest period
P (60 mg/m ²)	1	2	3	4			period					period

Bz= Bortezomib Accord; M=melphalan, P=prednisone

Dose adjustments during treatment and re-initiation of treatment for combination therapy with melphalan and prednisone

Prior to initiating a new cycle of therapy:

- Platelet counts should be $\geq 70 \times 10^9/l$ and the absolute neutrophils count should be $\geq 1.0 \times 10^9/l$
- Non-haematological toxicities should have resolved to Grade 1 or baseline

Table 3: Posology modifications during subsequent cycles of Bortezomib Accord therapy in combination with melphalan and prednisone

Toxicity	Posology modification or delay
<i>Haematological toxicity during a cycle</i>	
• If prolonged Grade 4 neutropenia or thrombocytopenia, or thrombocytopenia with bleeding is observed in the previous cycle	Consider reduction of the melphalan dose by 25% in the next cycle.
• If platelet counts $\leq 30 \times 10^9/l$ or ANC $\leq 0.75 \times 10^9/l$ on a Bortezomib Accord dosing day (other than day 1)	Bortezomib Accord therapy should be withheld
• If several Bortezomib Accord doses in a cycle are withheld (≥ 3 doses during twice weekly administration or ≥ 2 doses during weekly administration)	Bortezomib Accord dose should be reduced by 1 dose level (from 1.3 mg/m ² to 1 mg/m ² , or from 1 mg/m ² to 0.7 mg/m ²)
<i>Grade ≥ 3 non-haematological toxicities</i>	Bortezomib Accord therapy should be withheld until symptoms of the toxicity have resolved to Grade 1 or baseline. Then, Bortezomib Accord may be reinitiated with one dose level reduction (from

Toxicity	Posology modification or delay
	1.3 mg/m ² to 1 mg/m ² , or from 1 mg/m ² to 0.7 mg/m ²). For bortezomib-related neuropathic pain and/or peripheral neuropathy, hold and/or modify Bortezomib Accord as outlined in Table 1.

For additional information concerning melphalan and prednisone, see the corresponding Summary of Product Characteristics.

Posology for previously untreated multiple myeloma patients eligible for haematopoietic stem cell transplantation (induction therapy)

Combination therapy with dexamethasone

Bortezomib Accord is administered via intravenous or subcutaneous injection at the recommended dose of 1.3 mg/m² body surface area twice weekly for two weeks on days 1, 4, 8, and 11 in a 21-day treatment cycle. This 3-week period is considered a treatment cycle.

At least 72 hours should elapse between consecutive doses of Bortezomib Accord.

Dexamethasone is administered orally at 40 mg on days 1, 2, 3, 4, 8, 9, 10 and 11 of the Bortezomib Accord treatment cycle.

Four treatment cycles of this combination therapy are administered.

Combination therapy with dexamethasone and thalidomide

Bortezomib Accord is administered via intravenous or subcutaneous injection at the recommended dose of 1.3 mg/m² body surface area twice weekly for two weeks on days 1, 4, 8, and 11 in a 28-day treatment cycle. This 4-week period is considered a treatment cycle.

At least 72 hours should elapse between consecutive doses of Bortezomib Accord.

Dexamethasone is administered orally at 40 mg on days 1, 2, 3, 4, 8, 9, 10 and 11 of the Bortezomib Accord treatment cycle.

Thalidomide is administered orally at 50 mg daily on days 1-14 and if tolerated the dose is increased to 100 mg on days 15-28, and thereafter may be further increased to 200 mg daily from cycle 2 (see Table 4).

Four treatment cycles of this combination are administered. It is recommended that patients with at least partial response receive 2 additional cycles.

Table 4: Posology for Bortezomib Accord combination therapy for patients with previously untreated multiple myeloma eligible for haematopoietic stem cell transplantation

Bz+ Dx	Cycles 1 to 4				
	Week	1	2	3	
	Bz (1.3 mg/m ²)	Day 1, 4	Day 8, 11	Rest Period	
	Dx 40 mg	Day 1, 2, 3, 4	Day 8, 9, 10, 11	-	
Bz+ Dx+T	Cycles 1				
	Week	1	2	3	4
	Bz (1.3 mg/m ²)	Day 1, 4	Day 8, 11	Rest Period	Rest Period
	T 50 mg	Daily	Daily	-	-
	T 100 mg ^a	-	-	Daily	Daily
	Dx 40 mg	Day 1, 2, 3, 4	Day 8, 9, 10, 11	-	-
	Cycles 2 to 4 ^b				
	Bz (1.3 mg/m ²)	Day 1, 4	Day 8, 11	Rest Period	Rest Period
	T 200 mg ^a	Daily	Daily	Daily	Daily
	Dx 40 mg	Day 1, 2, 3, 4	Day 8, 9, 10, 11	-	-

Bz= Bortezomib Accord; Dx=dexamethasone; T=thalidomide

^a Thalidomide dose is increased to 100 mg from week 3 of Cycle 1 only if 50 mg is tolerated and to 200 mg from cycle 2 onwards if 100 mg is tolerated.

^b Up to 6 cycles may be given to patients who achieve at least a partial response after 4 cycles

Dosage adjustments for transplant eligible patients

For Bortezomib Accord dosage adjustments for neuropathy refer to Table 1.

In addition, when Bortezomib Accord is given in combination with other chemotherapeutic medicinal products, appropriate dose reductions for these products should be considered in the event of toxicities according to the recommendations in the Summary of Product Characteristics.

Posology for patients with previously untreated mantle cell lymphoma (MCL)

Combination therapy with rituximab, cyclophosphamide, doxorubicin and prednisone (BzR-CAP)

Bortezomib Accord is administered via intravenous or subcutaneous injection at the recommended dose of 1.3 mg/m² body surface area twice weekly for two weeks on days 1, 4, 8, and 11, followed by a 10-day rest period on days 12-21. This 3-week period is considered a treatment cycle. Six bortezomib cycles are recommended, although for patients with a response first documented at cycle 6, two additional bortezomib cycles may be given. At least 72 hours should elapse between consecutive doses of Bortezomib Accord.

The following medicinal products are administered on day 1 of each bortezomib 3 week treatment cycle as intravenous infusions: rituximab at 375 mg/m², cyclophosphamide at 750 mg/m² and doxorubicin at 50 mg/m².

Prednisone is administered orally at 100 mg/m² on days 1, 2, 3, 4 and 5 of each bortezomib treatment cycle.

Dose adjustments during treatment for patients with previously untreated mantle cell lymphoma

Prior to initiating a new cycle of therapy:

- Platelet counts should be $\geq 100,000$ cells/ μ L and the absolute neutrophils count (ANC) should be $\geq 1,500$ cells/ μ L
- Platelet counts should be $\geq 75,000$ cells/ μ L in patients with bone marrow infiltration or splenic sequestration
- Haemoglobin ≥ 8 g/dL
- Non-haematological toxicities should have resolved to Grade 1 or baseline.

Bortezomib treatment must be withheld at the onset of any $\geq$ Grade 3 bortezomib-related non-haematological toxicities (excluding neuropathy) or $\geq$ Grade 3 haematological toxicities. For dose adjustments, see Table 5 below.

Granulocyte colony stimulating factors may be administered for haematologic toxicity according to local standard practice. Prophylactic use of granulocyte colony stimulating factors should be considered in case of repeated delays in cycle administration. Platelet transfusion for the treatment of thrombocytopenia should be considered when clinically appropriate.

Table 5: Dose adjustments during treatment for patients with previously untreated mantle cell lymphoma

Toxicity	Posology modification or delay
<i>Haematological toxicity</i>	
• $\geq$ Grade 3 neutropenia with fever, Grade 4 neutropenia lasting more than 7 days, a platelet count $< 10,000$ cells/μL	Bortezomib Accord therapy should be withheld for up to 2 weeks until the patient has an ANC ≥ 750 cells/μL and a platelet count $\geq 25,000$ cells/μL. • If, after Bortezomib Accord has been held, the

Toxicity	Posology modification or delay
<i>Haematological toxicity</i>	
	toxicity does not resolve, as defined above, then Bortezomib Accord must be discontinued. If toxicity resolves i.e. patient has an ANC ≥ 750 cells/μL and a platelet count $\geq 25,000$ cells/μL, Bortezomib Accord may be reinitiated at a dose reduced by one dose level (from 1.3 mg/m² to 1 mg/m², or from 1 mg/m² to 0.7 mg/m²).
If platelet counts $< 25,000$ cells/μL. or ANC < 750 cells/μL on a Bortezomib Accord dosing day (other than Day 1 of each cycle)	Bortezomib Accord therapy should be withheld
<i>Grade ≥ 3 non-haematological toxicities considered to be related to Bortezomib Accord</i>	Bortezomib Accord therapy should be withheld until symptoms of the toxicity have resolved to Grade 2 or better. Then, Bortezomib Accord may be reinitiated at a dose reduced by one dose level (from 1.3 mg/m ² to 1 mg/m ² , or from 1 mg/m ² to 0.7 mg/m ²). For bortezomib-related neuropathic pain and/or peripheral neuropathy, hold and/or modify Bortezomib Accord as outlined in Table 1.

In addition, when bortezomib is given in combination with other chemotherapeutic medicinal products, appropriate dose reductions for these medicinal products should be considered in the event of toxicities, according to the recommendations in the respective Summary of Product Characteristics.

Special populations

Elderly

There is no evidence to suggest that dose adjustments are necessary in patients over 65 years of age with multiple myeloma or with mantle cell lymphoma.

There are no studies on the use of bortezomib in elderly patients with previously untreated multiple myeloma who are eligible for high-dose chemotherapy with haematopoietic stem cell transplantation. Therefore no dose recommendations can be made in this population.

In a study in previously untreated mantle cell lymphoma patients, 42.9% and 10.4% of patients exposed to bortezomib were in the range 65-74 years and ≥ 75 years of age, respectively. In patients aged ≥ 75 years, both regimens, BzR-CAP as well as R-CHOP, were less tolerated.

Hepatic impairment

Patients with mild hepatic impairment do not require a dose adjustment and should be treated per the recommended dose. Patients with moderate or severe hepatic impairment should be started on Bortezomib Accord at a reduced dose of 0.7 mg/m² per injection during the first treatment cycle, and a subsequent dose escalation to 1.0 mg/m² or further dose reduction to 0.5 mg/m² may be considered based on patient tolerability.

Table 6: Recommended starting dose modification for Bortezomib Accord in patients with hepatic impairment

Grade of hepatic impairment*	Bilirubin level	SGOT (AST) levels	Modification of starting dose
Mild	$\leq 1.0 \times \text{ULN}$	$> \text{ULN}$	None
	$> 1.0 \times -1.5 \times \text{ULN}$	Any	None
Moderate	$> 1.5 \times -3 \times \text{ULN}$	Any	Reduce Bortezomib Accord to 0.7 mg/m^2 in the first treatment cycle. Consider dose escalation to 1.0 mg/m^2 or further dose reduction to 0.5 mg/m^2 in subsequent cycles based on patient tolerability.
Severe	$> 3 \times \text{ULN}$	Any	

Abbreviations: SGOT=serum glutamic oxaloacetic transaminase;

AST=aspartate aminotransferase; ULN=upper limit of the normal range.

* Based on NCI Organ Dysfunction Working Group classification for categorising hepatic impairment (mild, moderate, severe).

Renal impairment

The pharmacokinetics of bortezomib are not influenced in patients with mild to moderate renal impairment (Creatinine Clearance [CrCL] $> 20 \text{ ml/min/1.73 m}^2$); therefore, dose adjustments are not necessary for these patients. It is unknown if the pharmacokinetics of bortezomib are influenced in patients with severe renal impairment not undergoing dialysis (CrCL $< 20 \text{ ml/min/1.73 m}^2$). Since dialysis may reduce bortezomib concentrations, Bortezomib Accord should be administered after the dialysis procedure.

Paediatric population

The safety and efficacy of bortezomib in children below 18 years of age have not been established. No data are available.

Method of administration

Bortezomib Accord is available for intravenous or subcutaneous administration.

Bortezomib Accord should not be given by other routes. Intrathecal administration has resulted in death.

Intravenous injection

Bortezomib Accord is administered as a 3-5 second bolus intravenous injection through a peripheral or central intravenous catheter followed by a flush with sodium chloride 9 mg/ml (0.9%) solution for injection. At least 72 hours should elapse between consecutive doses of Bortezomib Accord.

Subcutaneous injection

Bortezomib Accord is administered subcutaneously through the thighs (right or left) or abdomen (right or left). The solution should be injected subcutaneously, at a $45-90^\circ$ angle.

Injection sites should be rotated for successive injections.

If local injection site reactions occur following Bortezomib Accord subcutaneous injection, either a less concentrated Bortezomib Accord solution (Bortezomib Accord 3.5 mg to be reconstituted to 1 mg/ml instead of 2.5 mg/ml) may be administered subcutaneously or a switch to intravenous injection is recommended.

When Bortezomib Accord is given in combination with other medicinal products, refer to the Summary of Product Characteristics of these products for instructions for administration.

- **Route of Administration**

Parenteral (SC and IV)

- **Contraindications**

Hypersensitivity to the active substance, to boron or to any of the excipients listed in section A2.
Acute diffuse infiltrative pulmonary and pericardial disease.

When Bortezomib Accord is given in combination with other medicinal products, refer to their Summaries of Product Characteristics for additional contraindications.

- **Warnings & Precautions**

When Bortezomib Accord is given in combination with other medicinal products, the Summary of Product Characteristics of these other medicinal products must be consulted prior to initiation of treatment with Bortezomib Accord. When thalidomide is used, particular attention to pregnancy testing and prevention requirements is needed.

Intrathecal administration

Bortezomib Accord is for intravenous or subcutaneous use. Bortezomib Accord should not be administered intrathecally.

Gastrointestinal toxicity

Gastrointestinal toxicity, including nausea, diarrhoea, vomiting and constipation are very common with bortezomib treatment. Therefore, patients who experience constipation should be closely monitored.

Haematological toxicity

Bortezomib treatment is very commonly associated with haematological toxicities (thrombocytopenia, neutropenia and anaemia).

Gastrointestinal and intracerebral haemorrhage, have been reported in association with bortezomib treatment. Therefore, platelet counts should be monitored prior to each dose of bortezomib. Bortezomib therapy should be withheld when the platelet count is $< 25,000/\mu\text{l}$ or, in the case of combination with melphalan and prednisone, when the platelet count is $\leq 30,000/\mu\text{l}$. Potential benefit of the treatment should be carefully weighed against the risks, particularly in case of moderate to severe thrombocytopenia and risk factors for bleeding.

Complete blood counts (CBC) with differential and including platelet counts should be frequently monitored throughout treatment with bortezomib. Platelet transfusion should be considered when clinically appropriate.

Since patients with neutropenia are at increased risk of infections, they should be monitored for signs and symptoms of infection and treated promptly. Granulocyte colony stimulating factors may be administered for haematologic toxicity according to local standard practice. Prophylactic use of granulocyte colony stimulating factors should be considered in case of repeated delays in cycle administration.

Herpes zoster virus reactivation

Antiviral prophylaxis is recommended in patients being treated with bortezomib.

Hepatitis B Virus (HBV) reactivation and infection

When rituximab is used in combination with bortezomib, HBV screening must always be performed in patients at risk of infection with HBV before initiation of treatment. Carriers of hepatitis B and patients with a history of hepatitis B must be closely monitored for clinical and laboratory signs of active HBV infection during and following rituximab combination treatment with bortezomib. Antiviral prophylaxis should be considered. Refer to the Summary of Product Characteristics of rituximab for more information.

Progressive multifocal leukoencephalopathy (PML)

Very rare cases with unknown causality of John Cunningham (JC) virus infection, resulting in PML and death, have been reported in patients treated with bortezomib. Patients diagnosed with PML had prior or concurrent immunosuppressive therapy. Most cases of PML were diagnosed within 12 months of their first dose of bortezomib. Patients should be monitored at regular intervals for any new or worsening neurological symptoms or signs that may be suggestive of PML as part of the differential diagnosis of CNS problems. If a diagnosis of PML is suspected, patients should be referred to a specialist in PML and appropriate diagnostic measures for PML should be initiated. Discontinue bortezomib if PML is diagnosed.

Peripheral neuropathy

Treatment with bortezomib is very commonly associated with peripheral neuropathy, which is predominantly sensory. However, cases of severe motor neuropathy with or without sensory peripheral neuropathy have been reported. The incidence of peripheral neuropathy increases early in the treatment and has been observed to peak during cycle 5.

It is recommended that patients be carefully monitored for symptoms of neuropathy such as a burning sensation, hyperesthesia, hypoesthesia, paraesthesia, discomfort, neuropathic pain or weakness.

Patients experiencing new or worsening peripheral neuropathy should undergo neurological evaluation and may require a change in the dose, schedule or route of administration to subcutaneous. Neuropathy has been managed with supportive care and other therapies.

Early and regular monitoring for symptoms of treatment-emergent neuropathy with neurological evaluation should be considered in patients receiving bortezomib in combination with medicinal products known to be associated with neuropathy (e.g. thalidomide) and appropriate dose reduction or treatment discontinuation should be considered.

In addition to peripheral neuropathy, there may be a contribution of autonomic neuropathy to some adverse reactions such as postural hypotension and severe constipation with ileus. Information on autonomic neuropathy and its contribution to these undesirable effects is limited.

Seizures

Seizures have been uncommonly reported in patients without previous history of seizures or epilepsy. Special care is required when treating patients with any risk factors for seizures.

Hypotension

Bortezomib treatment is commonly associated with orthostatic/postural hypotension. Most adverse reactions are mild to moderate in nature and are observed throughout treatment. Patients who developed orthostatic hypotension on bortezomib (injected intravenously) did not have evidence of orthostatic

hypotension prior to treatment with bortezomib. Most patients required treatment for their orthostatic hypotension. A minority of patients with orthostatic hypotension experienced syncopal events. Orthostatic/postural hypotension was not acutely related to bolus infusion of bortezomib. The mechanism of this event is unknown although a component may be due to autonomic neuropathy. Autonomic neuropathy may be related to bortezomib or bortezomib may aggravate an underlying condition such as diabetic or amyloidotic neuropathy. Caution is advised when treating patients with a history of syncope receiving medicinal products known to be associated with hypotension; or who are dehydrated due to recurrent diarrhoea or vomiting. Management of orthostatic/postural hypotension may include adjustment of antihypertensive medicinal products, rehydration or administration of mineralocorticosteroids and/or sympathomimetics. Patients should be instructed to seek medical advice if they experience symptoms of dizziness, light-headedness or fainting spells.

Posterior Reversible Encephalopathy Syndrome (PRES)

There have been reports of *PRES* in patients receiving bortezomib. *PRES* is a rare, often reversible, rapidly evolving neurological condition, which can present with seizure, hypertension, headache, lethargy, confusion, blindness, and other visual and neurological disturbances. Brain imaging, preferably Magnetic Resonance Imaging (MRI), is used to confirm the diagnosis. In patients developing *PRES*, bortezomib should be discontinued.

Heart failure

Acute development or exacerbation of congestive heart failure, and/or new onset of decreased left ventricular ejection fraction has been reported during bortezomib treatment. Fluid retention may be a predisposing factor for signs and symptoms of heart failure. Patients with risk factors for or existing heart disease should be closely monitored.

Electrocardiogram investigations

There have been isolated cases of QT-interval prolongation in clinical studies, causality has not been established.

Pulmonary disorders

There have been rare reports of acute diffuse infiltrative pulmonary disease of unknown aetiology such as pneumonitis, interstitial pneumonia, lung infiltration, and acute respiratory distress syndrome (ARDS) in patients receiving bortezomib. Some of these events have been fatal. A pre-treatment chest radiograph is recommended to serve as a baseline for potential post-treatment pulmonary changes.

In the event of new or worsening pulmonary symptoms (e.g., cough, dyspnoea), a prompt diagnostic evaluation should be performed and patients treated appropriately. The benefit/risk ratio should be considered prior to continuing bortezomib therapy.

Renal impairment

Renal complications are frequent in patients with multiple myeloma. Patients with renal impairment should be monitored closely.

Hepatic impairment

Bortezomib is metabolised by liver enzymes. Bortezomib exposure is increased in patients with moderate or severe hepatic impairment; these patients should be treated with bortezomib at reduced doses and closely monitored for toxicities.

Hepatic reactions

Rare cases of hepatic failure have been reported in patients receiving bortezomib and concomitant medicinal products and with serious underlying medical conditions. Other reported hepatic reactions

include increases in liver enzymes, hyperbilirubinaemia, and hepatitis. Such changes may be reversible upon discontinuation of bortezomib.

Tumour lysis syndrome

Because bortezomib is a cytotoxic agent and can rapidly kill malignant plasma cells and MCL cells, the complications of tumour lysis syndrome may occur. The patients at risk of tumour lysis syndrome are those with high tumour burden prior to treatment. These patients should be monitored closely and appropriate precautions taken.

Concomitant medicinal products

Patients should be closely monitored when given bortezomib in combination with potent CYP3A4-inhibitors. Caution should be exercised when bortezomib is combined with CYP3A4- or CYP2C19 substrates.

Normal liver function should be confirmed and caution should be exercised in patients receiving oral hypoglycemics.

Potentially immunocomplex-mediated reactions

Potentially immunocomplex-mediated reactions, such as serum-sickness-type reaction, polyarthritis with rash and proliferative glomerulonephritis have been reported uncommonly. Bortezomib should be discontinued if serious reactions occur.

Interaction with other medicaments

In vitro studies indicate that bortezomib is a weak inhibitor of the cytochrome P450 (CYP) isozymes 1A2, 2C9, 2C19, 2D6 and 3A4. Based on the limited contribution (7%) of CYP2D6 to the metabolism of bortezomib, the CYP2D6 poor metaboliser phenotype is not expected to affect the overall disposition of bortezomib.

A drug-drug interaction study assessing the effect of ketoconazole, a potent CYP3A4 inhibitor, on the pharmacokinetics of bortezomib (injected intravenously), showed a mean bortezomib AUC increase of 35% (CI₉₀% [1.032 to 1.772]) based on data from 12 patients. Therefore, patients should be closely monitored when given bortezomib in combination with potent CYP3A4 inhibitors (e.g. ketoconazole, ritonavir).

In a drug-drug interaction study assessing the effect of omeprazole, a potent CYP2C19 inhibitor, on the pharmacokinetics of bortezomib (injected intravenously), there was no significant effect on the pharmacokinetics of bortezomib based on data from 17 patients.

A drug-drug interaction study assessing the effect of rifampicin, a potent CYP3A4 inducer, on the pharmacokinetics of bortezomib (injected intravenously), showed a mean bortezomib AUC reduction of 45% based on data from 6 patients. Therefore, the concomitant use of bortezomib with strong CYP3A4 inducers (e.g., rifampicin, carbamazepine, phenytoin, phenobarbital and St. John's Wort) is not recommended, as efficacy may be reduced.

In the same drug-drug interaction study assessing the effect of dexamethasone, a weaker CYP3A4 inducer, on the pharmacokinetics of bortezomib (injected intravenously), there was no significant effect on the pharmacokinetics of bortezomib based on data from 7 patients.

A drug-drug interaction study assessing the effect of melphalan-prednisone on the pharmacokinetics of bortezomib (injected intravenously), showed a mean bortezomib AUC increase of 17% based on data from 21 patients. This is not considered clinically relevant.

During clinical trials, hypoglycemia and hyperglycemia were uncommonly and commonly reported in diabetic patients receiving oral hypoglycemics. Patients on oral antidiabetic agents receiving bortezomib treatment may require close monitoring of their blood glucose levels and adjustment of the dose of their antidiabetics.

- **Statement on usage during pregnancy and lactation**

Contraception in males and females

Male and female patients of childbearing potential must use effective contraceptive measures during and for 3 months following treatment.

Pregnancy

No clinical data are available for bortezomib with regard to exposure during pregnancy. The teratogenic potential of bortezomib has not been fully investigated.

In non-clinical studies, bortezomib had no effects on embryonal/foetal development in rats and rabbits at the highest maternally tolerated doses. Animal studies to determine the effects of bortezomib on parturition and post-natal development were not conducted. Bortezomib should not be used during pregnancy unless the clinical condition of the woman requires treatment with bortezomib. If bortezomib is used during pregnancy, or if the patient becomes pregnant while receiving this medicinal product, the patient should be informed of potential for hazard to the foetus.

Thalidomide is a known human teratogenic active substance that causes severe life-threatening birth defects. Thalidomide is contraindicated during pregnancy and in women of childbearing potential unless all the conditions of the thalidomide pregnancy prevention programme are met. Patients receiving bortezomib in combination with thalidomide should adhere to the pregnancy prevention programme of thalidomide. Refer to the Summary of Product Characteristics of thalidomide for additional information.

Breast-feeding

It is not known whether bortezomib is excreted in human milk. Because of the potential for serious adverse reactions in breast-fed infants, breast-feeding should be discontinued during treatment with bortezomib.

Fertility

Fertility studies were not conducted with bortezomib.

- **Adverse Effects/ Undesirable Effects**

Summary of the safety profile

Serious adverse reactions uncommonly reported during treatment with bortezomib include cardiac failure, tumour lysis syndrome, pulmonary hypertension, posterior reversible encephalopathy syndrome, acute diffuse infiltrative pulmonary disorders and rarely autonomic neuropathy. The most commonly reported adverse reactions during treatment with bortezomib are nausea, diarrhoea, constipation, vomiting, fatigue, pyrexia, thrombocytopenia, anaemia, neutropenia, peripheral neuropathy (including sensory), headache, paraesthesia, decreased appetite, dyspnoea, rash, herpes zoster and myalgia.

Tabulated summary of adverse reactions

Multiple Myeloma

Undesirable effects in Table 7 were considered by the investigators to have at least a possible or probable causal relationship to bortezomib. These adverse reactions are based on an integrated data set of 5,476 patients of whom 3,996 were treated with bortezomib at 1.3 mg/m² and included in Table 7.

Overall, bortezomib was administered for the treatment of multiple myeloma in 3,974 patients.

Adverse reactions are listed below by system organ class and frequency grouping. Frequencies are defined as: Very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$), not known (cannot be estimated from the available data). Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness. Table 7 has been generated using Version 14.1 of the MedDRA.

Post-marketing adverse reactions not seen in clinical trials are also included.

Table 7: Adverse reactions in patients treated with Multiple Myeloma treated with bortezomib as single agent or in combination

System Organ Class	Incidence	Adverse reaction
Infections and infestations	Common	Herpes zoster (inc disseminated & ophthalmic), Pneumonia*, Herpes simplex*, Fungal infection*
	Uncommon	Infection*, Bacterial infections*, Viral infections*, Sepsis (inc septic shock)*, Bronchopneumonia, Herpes virus infection*, Meningoencephalitis herpetic [#] , Bacteraemia (inc staphylococcal), Hordeolum, Influenza, Cellulitis, Device related infection, Skin infection*, Ear infection*, Staphylococcal infection, Tooth infection*
	Rare	Meningitis (inc bacterial), Epstein-Barr virus infection, Genital herpes, Tonsillitis, Mastoiditis, Post viral fatigue syndrome
Neoplasms benign, malignant and unspecified (incl cysts and polyps)	Rare	Neoplasm malignant, Leukaemia plasmacytic, Renal cell carcinoma, Mass, Mycosis fungoides, Neoplasm benign*
Blood and lymphatic system disorders	Very Common	Thrombocytopenia*, Neutropenia*, Anaemia*
	Common	Leukopenia*, Lymphopenia*
	Uncommon	Pancytopenia*, Febrile neutropenia, Coagulopathy*, Leukocytosis*, Lymphadenopathy, Haemolytic anaemia [#]
	Rare	Disseminated intravascular coagulation, Thrombocytosis*, Hyperviscosity syndrome, Platelet disorder NOS, Thrombocytopenic purpura, Blood disorder NOS, Haemorrhagic diathesis, Lymphocytic infiltration
Immune system Disorders	Uncommon	Angioedema [#] , Hypersensitivity*
	Rare	Anaphylactic shock, Amyloidosis, Type III immune complex mediated reaction
Endocrine disorders	Uncommon	Cushing's syndrome*, Hyperthyroidism*, Inappropriate antidiuretic hormone secretion
	Rare	Hypothyroidism
Metabolism and nutrition disorders	Very Common	Decreased appetite
	Common	Dehydration, Hypokalaemia*, Hyponatraemia*, Blood glucose abnormal*, Hypocalcaemia*, Enzyme abnormality*

System Organ Class	Incidence	Adverse reaction
	Uncommon	Tumour lysis syndrome, Failure to thrive*, Hypomagnesaemia*, Hypophosphataemia*, Hyperkalaemia*, Hypercalcaemia*, Hyponatraemia*, Uric acid abnormal*, Diabetes mellitus*, Fluid retention
	Rare	Hypermagnesaemia*, Acidosis, Electrolyte imbalance*, Fluid overload, Hypochloraemia*, Hypovolaemia, Hyperchloraemia*, Hyperphosphataemia*, Metabolic disorder, Vitamin B complex deficiency, Vitamin B12 deficiency, Gout, Increased appetite, Alcohol intolerance
Psychiatric disorders	Common	Mood disorders and disturbances*, Anxiety disorder*, Sleep disorders and disturbances*
	Uncommon	Mental disorder*, Hallucination*, Psychotic disorder*, Confusion*, Restlessness
	Rare	Suicidal ideation*, Adjustment disorder, Delirium, Libido decreased
Nervous system disorders	Very Common	Neuropathies*, Peripheral sensory neuropathy, Dysaesthesia*, Neuralgia*
	Common	Motor neuropathy*, Loss of consciousness (inc syncope), Dizziness*, Dysgeusia*, Lethargy, Headache*
	Uncommon	Tremor, Peripheral sensorimotor neuropathy, Dyskinesia*, Cerebellar coordination and balance disturbances*, Memory loss (exc dementia)*, Encephalopathy*, Posterior Reversible Encephalopathy Syndrome#, Neurotoxicity, Seizure disorders*, Post herpetic neuralgia, Speech disorder*, Restless legs syndrome, Migraine, Sciatica, Disturbance in attention, Reflexes abnormal*, Parosmia
	Rare	Cerebral haemorrhage*, Haemorrhage intracranial (inc subarachnoid)*, Brain oedema, Transient ischaemic attack, Coma, Autonomic nervous system imbalance, Autonomic neuropathy, Cranial palsy*, Paralysis*, Paresis*, Presyncope, Brain stem syndrome, Cerebrovascular disorder, Nerve root lesion, Psychomotor hyperactivity, Spinal cord compression, Cognitive disorder NOS, Motor dysfunction, Nervous system disorder NOS, Radiculitis, Drooling, Hypotonia, Guillain-Barré syndrome, Demyelinating polyneuropathy
Eye disorders	Common	Eye swelling*, Vision abnormal*, Conjunctivitis*
	Uncommon	Eye haemorrhage*, Eyelid infection*, Eye inflammation*, Diplopia, Dry eye*, Eye irritation*, Eye pain, Lacrimation increased, Eye discharge
	Rare	Corneal lesion*, Exophthalmos, Retinitis, Scotoma, Eye disorder (inc. eyelid) NOS, Dacryoadenitis acquired, Photophobia, Photopsia, Optic neuropathy#, Different degrees of visual impairment (up to blindness)*
Ear and labyrinth disorders	Common	Vertigo*
	Uncommon	Dysacusis (inc tinnitus)*, Hearing impaired (up to and inc deafness), Ear discomfort*
	Rare	Ear haemorrhage, Vestibular neuronitis, Ear disorder NOS
Cardiac disorders	Uncommon	Cardiac tamponade#, Cardio-pulmonary arrest*, Cardiac fibrillation (inc atrial), Cardiac failure (inc left and right

System Organ Class	Incidence	Adverse reaction
		ventricular)*, Arrhythmia*, Tachycardia*, Palpitations, Angina pectoris, Pericarditis (inc pericardial effusion)*, Cardiomyopathy*, Ventricular dysfunction*, Bradycardia
	Rare	Atrial flutter, Myocardial infarction*, Atrioventricular block*, Cardiovascular disorder (inc cardiogenic shock), Torsade de pointes, Angina unstable, Cardiac valve disorders*, Coronary artery insufficiency, Sinus arrest
Vascular disorders	Common	Hypotension*, Orthostatic hypotension, Hypertension*
	Uncommon	Cerebrovascular accident [#] , Deep vein thrombosis*, Haemorrhage*, Thrombophlebitis (inc superficial), Circulatory collapse (inc hypovolaemic shock), Phlebitis, Flushing*, Haematoma (inc perirenal)*, Poor peripheral circulation*, Vasculitis, Hyperaemia (inc ocular)*
	Rare	Peripheral embolism, Lymphoedema, Pallor, Erythromelalgia, Vasodilatation, Vein discolouration, Venous insufficiency
Respiratory, thoracic and mediastinal disorders	Common	Dyspnoea*, Epistaxis, Upper/lower respiratory tract infection*, Cough*
	Uncommon	Pulmonary embolism, Pleural effusion, Pulmonary oedema (inc acute), Pulmonary alveolar haemorrhage [#] , Bronchospasm, Chronic obstructive pulmonary disease*, Hypoxaemia*, Respiratory tract congestion*, Hypoxia, Pleurisy*, Hiccups, Rhinorrhoea, Dysphonia, Wheezing
	Rare	Respiratory failure, Acute respiratory distress syndrome, Apnoea, Pneumothorax, Atelectasis, Pulmonary hypertension, Haemoptysis, Hyperventilation, Orthopnoea, Pneumonitis, Respiratory alkalosis, Tachypnoea, Pulmonary fibrosis, Bronchial disorder*, Hypocapnia*, Interstitial lung disease, Lung infiltration, Throat tightness, Dry throat, Increased upper airway secretion, Throat irritation, Upper-airway cough syndrome
Gastrointestinal disorders	Very Common	Nausea and vomiting symptoms*, Diarrhoea*, Constipation
	Common	Gastrointestinal haemorrhage (inc mucosal)*, Dyspepsia, Stomatitis*, Abdominal distension, Oropharyngeal pain*, Abdominal pain (inc gastrointestinal and splenic pain)*, Oral disorder*, Flatulence
	Uncommon	Pancreatitis (inc chronic)*, Haematemesis, Lip swelling*, Gastrointestinal obstruction (inc ileus)*, Abdominal discomfort, Oral ulceration*, Enteritis*, Gastritis*, Gingival bleeding, Gastrooesophageal reflux disease*, Colitis (inc clostridium difficile)*, Colitis ischaemic [#] , Gastrointestinal inflammation*, Dysphagia, Irritable bowel syndrome, Gastrointestinal disorder NOS, Tongue coated, Gastrointestinal motility disorder*, Salivary gland disorder*
	Rare	Pancreatitis acute, Peritonitis*, Tongue oedema*, Ascites, Oesophagitis, Cheilitis, Faecal incontinence, Anal sphincter atony, Faecaloma*, Gastrointestinal ulceration and perforation*, Gingival hypertrophy, Megacolon, Rectal discharge, Oropharyngeal blistering*, Lip pain, Periodontitis,

System Organ Class	Incidence	Adverse reaction
		Anal fissure, Change of bowel habit, Proctalgia, Abnormal Faeces
Hepatobiliary disorders	Common	Hepatic enzyme abnormality*
	Uncommon	Hepatotoxicity (inc liver disorder), Hepatitis*, Cholestasis
	Rare	Hepatic failure, Hepatomegaly, Budd-Chiari syndrome, Cytomegalovirus hepatitis, Hepatic haemorrhage, Cholelithiasis
Skin and subcutaneous tissue disorders	Common	Rash*, Pruritus*, Erythema, Dry skin
	Uncommon	Erythema multiforme, Urticaria, Acute febrile neutrophilic dermatosis, Toxic skin eruption, Toxic epidermal necrolysis [#] , Stevens-Johnson syndrome [#] , Dermatitis*, Hair disorder*, Petechiae, Ecchymosis, Skin lesion, Purpura, Skin mass*, Psoriasis, Hyperhidrosis, Night sweats, Decubitus ulcer [#] , Acne*, Blister*, Pigmentation disorder*
	Rare	Skin reaction, Jessner's lymphocytic infiltration, Palmar-plantar erythrodysesthesia syndrome, Haemorrhage subcutaneous, Livedo reticularis, Skin induration, Papule, Photosensitivity reaction, Seborrhoea, Cold sweat, Skin disorder NOS, Erythrosis, Skin ulcer, Nail disorder
Musculoskeletal and connective tissue disorders	Very Common	Musculoskeletal pain*
	Common	Muscle spasms*, Pain in extremity, Muscular weakness
	Uncommon	Muscle twitching, Joint swelling, Arthritis*, Joint stiffness, Myopathies*, Sensation of heaviness
	Rare	Rhabdomyolysis, Temporomandibular joint syndrome, Fistula, Joint effusion, Pain in jaw, Bone disorder, Musculoskeletal and connective tissue infections and inflammations*, Synovial cyst
Renal and urinary disorders	Common	Renal impairment*
	Uncommon	Renal failure acute, Renal failure chronic*, Urinary tract infection*, Urinary tract signs and symptoms*, Haematuria*, Urinary retention, Micturition disorder*, Proteinuria, Azotaemia, Oliguria*, Pollakiuria
	Rare	Bladder irritation
Reproductive system and breast disorders	Uncommon	Vaginal haemorrhage, Genital pain*, Erectile dysfunction,
	Rare	Testicular disorder*, Prostatitis, Breast disorder female, Epididymal tenderness, Epididymitis, Pelvic pain, Vulval ulceration
Congenital, familial and genetic disorders	Rare	Aplasia, Gastrointestinal malformation, Ichthyosis
General disorders and administration site conditions	Very Common	Pyrexia*, Fatigue, Asthenia
	Common	Oedema (inc peripheral), Chills, Pain*, Malaise*
	Uncommon	General physical health deterioration*, Face oedema*, Injection site reaction*, Mucosal disorder*, Chest pain, Gait disturbance, Feeling cold, Extravasation*, Catheter related complication*, Change in thirst*, Chest discomfort, Feeling of body temperature change*, Injection site pain*
	Rare	Death (inc sudden), Multi-organ failure, Injection site haemorrhage*, Hernia(inc hiatus)*, Impaired healing*, Inflammation, Injection site phlebitis*, Tenderness, Ulcer,

System Organ Class	Incidence	Adverse reaction
		Irritability, Non-cardiac chest pain, Catheter site pain, Sensation of foreign body
Investigations	Common	Weight decreased
	Uncommon	Hyperbilirubinaemia*, Protein analyses abnormal*, Weight increased, Blood test abnormal*, C-reactive protein increased
	Rare	Blood gases abnormal*, Electrocardiogram abnormalities (inc QT prolongation)*, International normalised ratio abnormal*, Gastric pH decreased, Platelet aggregation increased, Troponin I increased, Virus identification and serology*, Urine analysis abnormal*
Injury, poisoning and procedural complications	Uncommon	Fall, Contusion
	Rare	Transfusion reaction, Fractures*, Rigors*, Face injury, Joint injury*, Burns, Laceration, Procedural pain, Radiation injuries*
Surgical and medical procedures	Rare	Macrophage activation

NOS=not otherwise specified

* Grouping of more than one MedDRA preferred term.

Postmarketing adverse reaction

Mantle Cell Lymphoma (MCL)

The safety profile of bortezomib in 240 MCL patients treated with bortezomib at 1.3 mg/m² in combination with rituximab, cyclophosphamide, doxorubicin, and prednisone (BzR-CAP) versus 242 patients treated with rituximab, cyclophosphamide, doxorubicin, vincristine, and prednisone [R-CHOP] was relatively consistent to that observed in patients with multiple myeloma with main differences described below. Additional adverse drug reactions identified associated with the use of the combination therapy (BzR-CAP) were hepatitis B infection (< 1%) and myocardial ischaemia (1.3%). The similar incidences of these events in both treatment arms, indicated that these adverse drug reactions are not attributable to bortezomib alone. Notable differences in the MCL patient population as compared to patients in the multiple myeloma studies were a $\geq 5\%$ higher incidence of the haematological adverse reactions (neutropenia, thrombocytopenia, leukopenia, anemia, lymphopenia), peripheral sensory neuropathy, hypertension, pyrexia, pneumonia, stomatitis, and hair disorders.

Adverse drug reactions identified as those with a $\geq 1\%$ incidence, similar or higher incidence in the BzR-CAP arm and with at least a possible or probable causal relationship to the components of the BzR-CAP arm, are listed in Table 8 below. Also included are adverse drug reactions identified in the BzR-CAP arm that were considered by investigators to have at least a possible or probable causal relationship to bortezomib based on historical data in the multiple myeloma studies.

Adverse reactions are listed below by system organ class and frequency grouping. Frequencies are defined as: Very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$), not known (cannot be estimated from the available data). Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness. Table 8 has been generated using Version 16 of the MedDRA.

Table 8 Adverse reactions in patients with Mantle Cell Lymphoma treated with BzR-CAP

System Organ Class	Incidence	Adverse reaction
Infections and	Very Common	Pneumonia*

System Organ Class	Incidence	Adverse reaction
infestations	Common	Sepsis (inc septic shock)*, Herpes zoster (inc disseminated & ophthalmic), Herpes virus infection*, Bacterial infections*, Upper/lower respiratory tract infection*, Fungal infection*, Herpes simplex*
	Uncommon	Hepatitis B, Infection*, Bronchopneumonia
Blood and lymphatic system disorders	Very Common	Thrombocytopenia*, Febrile neutropenia, Neutropenia*, Leukopenia*, Anaemia*, Lymphopenia*
	Uncommon	Pancytopenia*
Immune system Disorders	Common	Hypersensitivity*
	Uncommon	Anaphylactic reaction
Metabolism and nutrition disorders	Very Common	Decreased appetite
	Common	Hypokalaemia*, Blood glucose abnormal*, Hyponatraemia*, Diabetes mellitus*, Fluid retention
	Uncommon	Tumour lysis syndrome
Psychiatric disorders	Common	Sleep disorders and disturbances*
Nervous system disorders	Very Common	Peripheral sensory neuropathy, Dysaesthesia*, Neuralgia*
	Common	Neuropathies*, Motor neuropathy*, Loss of consciousness (inc syncope), Encephalopathy*, Peripheral sensorimotor neuropathy, Dizziness*, Dysgeusia*, Autonomic neuropathy
	Uncommon	Autonomic nervous system imbalance
Eye disorders	Common	Vision abnormal*
Ear and labyrinth disorders	Common	Dysacusis (inc tinnitus)*
	Uncommon	Vertigo*, Hearing impaired (up to and inc deafness)
Cardiac disorders	Common	Cardiac fibrillation (inc atrial), Arrhythmia*, Cardiac failure (inc left and right ventricular)*, Myocardial ischaemia, Ventricular dysfunction*
	Uncommon	Cardiovascular disorder (inc cardiogenic shock)
Vascular disorders	Common	Hypertension*, Hypotension*, Orthostatic hypotension
Respiratory, thoracic and mediastinal disorders	Common	Dyspnoea*, Cough*, Hiccups
	Uncommon	Acute respiratory distress syndrome, Pulmonary embolism, Pneumonitis, Pulmonary hypertension, Pulmonary oedema (inc acute)
Gastrointestinal disorders	Very Common	Nausea and vomiting symptoms*, Diarrhoea*, Stomatitis*, Constipation
	Common	Gastrointestinal haemorrhage (inc mucosal)*, Abdominal distension, Dyspepsia, Oropharyngeal pain*, Gastritis*, Oral ulceration*, Abdominal discomfort, Dysphagia, Gastrointestinal inflammation*, Abdominal pain (inc gastrointestinal and splenic pain)*, Oral disorder*
	Uncommon	Colitis (inc clostridium difficile)*
Hepatobiliary disorders	Common	Hepatotoxicity (inc liver disorder)
	Uncommon	Hepatic failure
Skin and subcutaneous tissue disorders	Very Common	Hair disorder*
	Common	Pruritus*, Dermatitis*, Rash*
Musculoskeletal and connective tissue disorders	Common	Muscle spasms*, Musculoskeletal pain*, Pain in extremity

System Organ Class	Incidence	Adverse reaction
Renal and urinary disorders	Common	Urinary tract infection*
General disorders and administration site conditions	Very Common	Pyrexia*, Fatigue, Asthenia
	Common	Oedema (inc peripheral), Chills, Injection site reaction*, Malaise*
Investigations	Common	Hyperbilirubinaemia*, Protein analyses abnormal*, Weight decreased, Weight increased

* Grouping of more than one MedDRA preferred term.

Description of selected adverse reactions

Herpes zoster virus reactivation

Multiple Myeloma

Antiviral prophylaxis was administered to 26% of the patients in the Bz+M+P arm. The incidence of herpes zoster among patients in the Bz+M+P treatment group was 17% for patients not administered antiviral prophylaxis compared to 3% for patients administered antiviral prophylaxis.

Mantle cell lymphoma

Antiviral prophylaxis was administered to 137 of 240 patients (57%) in the BzR-CAP arm. The incidence of herpes zoster among patients in the BzR-CAP arm was 10.7% for patients not administered antiviral prophylaxis compared to 3.6% for patients administered antiviral prophylaxis.

Hepatitis B Virus (HBV) reactivation and infection

Mantle cell lymphoma

HBV infection with fatal outcomes occurred in 0.8% (n=2) of patients in the non-bortezomib treatment group (rituximab, cyclophosphamide, doxorubicin, vincristine, and prednisone; R-CHOP) and 0.4% (n=1) of patients receiving bortezomib in combination with rituximab, cyclophosphamide, doxorubicin, and prednisone (BzR-CAP). The overall incidence of hepatitis B infections was similar in patients treated with BzR-CAP or with R-CHOP (0.8% vs 1.2% respectively).

Peripheral neuropathy in combination regimens

Multiple Myeloma

In trials in which bortezomib was administered as induction treatment in combination with dexamethasone (study IFM-2005-01), and dexamethasone-thalidomide (study MMY-3010), the incidence of peripheral neuropathy in the combination regimens is presented in the table below:

Table 9: Incidence of peripheral neuropathy during induction treatment by toxicity and treatment discontinuation due to peripheral neuropathy

	<u>IFM-2005-01</u>		<u>MMY-3010</u>	
	VDDx (N=239)	BzDx (N=239)	TDx (N=126)	BzTDx (N=130)
Incidence of PN (%)				
All GradePN	3	15	12	45
≥ Grade 2 PN	1	10	2	31
≥ Grade 3 PN	< 1	5	0	5
Discontinuation due to PN (%)	< 1	2	1	5

VDDx=vincristine, doxorubicin, dexamethasone; BzDx= bortezomib, dexamethasone; TDx=thalidomide, dexamethasone; BzTDx= bortezomib, thalidomide, dexamethasone; PN=peripheral neuropathy

Note: Peripheral neuropathy included the preferred terms: neuropathy peripheral, peripheral motor neuropathy, peripheral sensory neuropathy, and polyneuropathy.

Mantle cell lymphoma

In study LYM-3002 in which bortezomib was administered with cyclophosphamide, doxorubicin, and prednisone (R-CAP), the incidence of peripheral neuropathy in the combination regimens is presented in the table below:

Table 10: Incidence of peripheral neuropathy in study LYM-3002 by toxicity and treatment discontinuation due to peripheral neuropathy

	BzR-CAP (N=240)	R-CHOP (N=242)
Incidence of PN (%)		
All GradePN	30	29
≥ Grade 2 PN	18	9
≥ Grade 3 PN	8	4
Discontinuation due to PN (%)	2	< 1

BzR-CAP=Bortezomib, rituximab, cyclophosphamide, doxorubicin, and prednisone; R-CHOP= rituximab, cyclophosphamide, doxorubicin, vincristine, and prednisone; PN=peripheral neuropathy

Peripheral neuropathy included the preferred terms: peripheral sensory neuropathy, neuropathy peripheral, peripheral motor neuropathy, and peripheral sensorimotor neuropathy

Elderly MCL patients

42.9% and 10.4% of patients in the BzR-CAP arm were in the range 65-74 years and ≥ 75 years of age, respectively. Although in patients aged ≥ 75 years, both BzR-CAP and R-CHOP were less tolerated, the serious adverse event rate in the BzR-CAP groups was 68%, compared to 42% in the RCHOP group.

Notable differences in the safety profile of bortezomib administered subcutaneously versus intravenously as single agent

In the Phase III study patients who received bortezomib subcutaneously compared to intravenous administration had 13% lower overall incidence of treatment emergent adverse reactions that were Grade 3 or higher in toxicity, and a 5% lower incidence of discontinuation of bortezomib. The overall incidence of diarrhoea, gastrointestinal and abdominal pain, asthenic conditions, upper respiratory tract infections and peripheral neuropathies were 12%-15% lower in the subcutaneous group than in the intravenous group. In addition, the incidence of Grade 3 or higher peripheral neuropathies was 10% lower, and the discontinuation rate due to peripheral neuropathies 8% lower for the subcutaneous group as compared to the intravenous group.

Six percent of patients had an adverse local reaction to subcutaneous administration, mostly redness. Cases resolved in a median of 6 days, dose modification was required in two patients. Two (1%) of the patients had severe reactions; 1 case of pruritus and 1 case of redness.

The incidence of death on treatment was 5% in the subcutaneous treatment group and 7% in the intravenous treatment group. Incidence of death from “Progressive disease” was 18% in the subcutaneous group and 9% in the intravenous group.

Retreatment of patients with relapsed multiple myeloma

In a study in which bortezomib retreatment was administered in 130 patients with relapsed multiple myeloma, who previously had at least partial response on a bortezomib-containing regimen, the most common all-grade adverse events occurring in at least 25% of patients were thrombocytopenia (55%), neuropathy (40%), anaemia (37%), diarrhoea (35%), and constipation (28%). All grade peripheral neuropathy and grade ≥ 3 peripheral neuropathy were observed in 40% and 8.5% of patients, respectively.

- **Overdose and Treatment**

In patients, overdose more than twice the recommended dose has been associated with the acute onset of symptomatic hypotension and thrombocytopenia with fatal outcomes.

There is no known specific antidote for bortezomib overdose. In the event of an overdose, the patient's vital signs should be monitored and appropriate supportive care given to maintain blood pressure (such as fluids, pressors, and/or inotropic agents) and body temperature.

- **Incompatibilities**

This medicinal product must not be mixed with other medicinal products except those mentioned in section **Special precautions for disposal and other handling**.

- **Special precautions for disposal and other handling**

General precautions

Bortezomib is a cytotoxic agent. Therefore, caution should be used during handling and preparation of Bortezomib Accord. Use of gloves and other protective clothing to prevent skin contact is recommended.

Aseptic technique must be strictly observed throughout the handling of Bortezomib Accord, since it contains no preservative.

There have been fatal cases of inadvertent intrathecal administration of bortezomib. Bortezomib Accord is for intravenous or subcutaneous use. Bortezomib Accord should not be administered intrathecally.

Instructions for reconstitution

Bortezomib Accord must be reconstituted by a healthcare professional.

Intravenous injection

Each 10 ml vial of Bortezomib Accord must be reconstituted with 3.5 ml of sodium chloride 9 mg/ml (0.9%) solution for injection. Dissolution of the lyophilised powder is completed in less than 2 minutes.

After reconstitution, each ml solution contains 1 mg bortezomib. The reconstituted solution is clear and colourless, with a final pH of 4 to 7.

The reconstituted solution must be inspected visually for particulate matter and discolouration prior to administration. If any discolouration or particulate matter is observed, the reconstituted solution must be discarded.

Subcutaneous injection

Each 10 ml vial of Bortezomib Accord should be reconstituted with 1.4 ml of sodium chloride 9 mg/ml (0.9%) solution for injection. Dissolution of the lyophilised powder is completed in less than 2 minutes.

After reconstitution, each ml solution contains 2.5 mg bortezomib. The reconstituted solution is clear and colourless, with a final pH of 4 to 7. The reconstituted solution must be inspected visually for particulate matter and discolouration prior to administration. If any discolouration or particulate matter is observed, the reconstituted solution must be discarded.

Disposal

Bortezomib Accord is for single use only. Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

- **Storage Conditions**

Store below 30°C. Keep the vial in the outer carton in order to protect from light.

For storage conditions after reconstitution of the medicinal product, see section **Shelf life**.

- **Shelf life**

Unopened vial

3 years

Intravenous administration

The chemical and physical in-use stability of the reconstituted solution at a concentration of 1 mg/ml has been demonstrated for 3 days at 20°C-25°C stored in the original vial and/or a syringe. From a microbiological point of view, unless the method of opening/reconstitution/dilution precludes the risk of microbial contamination, the reconstituted solution should be used immediately after preparation. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user.

Subcutaneous administration

The chemical and physical in-use stability of the reconstituted solution of 2.5 mg/ml has been demonstrated for 8 hours at 20°C-25°C stored in the original vial and/or a syringe. From a microbiological point of view, unless the method of opening/reconstitution/dilution precludes the risk of microbial contamination, the reconstituted solution should be used immediately after preparation. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user.

- **Dosage forms and packaging available**

Accord Bortezomib 3.5 is available in 1 vial pack.

- **Name and address of manufacturer**

INTAS PHARMACEUTICALS LIMITED

Plot 5, 6, 7, Pharmez,
Sarkhej-Bavla Highway,
Matoda, Sanand Taluka,
Ahmedabad, Gujarat, India.

- **Date of revision of PI**

April-2022