

VISCLEAR EYE DROP 0.5MG/ML

Ingredient(s):

Each ml contains:

Tetrahydrozoline Hydrochloride	0.5mg
Benzalkonium Chloride (as preservative)	0.1mg

Pharmacology (Summary of Pharmacodynamic and Pharmacokinetic):**Pharmacodynamic:**

The drug directly stimulates α -adrenergic receptors of the sympathetic nervous system and exerts little or no effect on β -adrenergic receptors. Following topical application of tetrahydrozoline to the conjunctiva, small arterioles are constricted and conjunctival congestion is temporarily relieved, but reactive hyperaemia may occur. The drug also may produce mydriasis when applied to the conjunctiva, but this effect usually minimal with the concentrations used as ocular decongestants.

Pharmacokinetic:

Following topical application of Tetrahydrozoline hydrochloride solutions to the conjunctiva local vasoconstriction usually occurs within a few minutes and may persist for 4-8 hours. Occasionally, enough tetrahydrozoline may be absorbed to produce systemic effects. Information on the distribution and elimination of the drug in human is not available.

Indication(s):

Provides temporary relief from soreness, burning, itching, irritation and congestion of the eyes in a variety of allergy and eye irritation conditions caused by allergies, smog, smoke, dust, wind and other eye irritants.

Dosage and Administration(s):

Place one or two drops in the affected eye(s) up to four times daily or as directed by a physician. Not recommended for use in children under 6 years of age except on medical advice. To avoid contamination of this product, do not touch dropper tip to other surfaces and always replace cap after using. Patients should be cautioned to remove contact lenses before using this product, and that contact lenses should not be reinserted until 15 minutes have elapsed after the instillation of the drops.

Mode of Administration(s):

Ophthalmic administration.

Contraindication(s):

Contraindicated in patients with a known hypersensitivity to any component of the product. This product should not be used in the presence of narrow-angle glaucoma and serious eye infections. If the solution changes colour or become cloudy, it should not be used.

Warning and Precautions(s):

Patients should be warned that if they experience eye pain, changes in vision, continued redness or irritation of the eye, or if the condition worsens or persists for more than 72 hours, that they should discontinue use and consult a doctor, as this may be sign of a more serious condition. Overuse of this product may produce increased redness of the eye.

This product should be used with caution in patients with glaucoma, other serious eye diseases, hyperthyroidism, heart disease, hypertension, or diabetes mellitus, or in those receiving monoamine oxidase inhibitors.

This product should not be substituted for contact lens fluid.

Interaction with Other Medicaments(s):

None known.

Pregnancy and Lactation(s):

Use in pregnancy: Safe use of this product during pregnancy has not been established

Use in lactation: None available.

Side Effects(s):

The incidence of serious adverse effects is low in patients receiving therapeutic dosages of topical tetrahydrozoline hydrochloride. Excessive dosage and/or prolonged or too frequent use may irritate conjunctiva, especially in children, and cause adverse systemic effects.

Use of tetrahydrozoline in the eye may cause blurred vision, irritation, and mydriasis. Conjunctival application of tetrahydrozoline, especially when high concentrations are used in geriatric patients, may liberate pigment granules, presumably from the iris. Prolonged use may worsen symptoms and should be avoided. In children, excessive dosage of the drug may cause severe drowsiness accompanied by profuse sweating.

Ophthalmic use of tetrahydrozoline may occasionally cause systemic sympathomimetic effects such as headache, hypertension, weakness, sweating, palpitation, and tremors. Overdosage of the drug may produce CNS depression with drowsiness, decreased body temperature, bradycardia, shock like hypotension, apnea, and coma.

Symptoms and Treatment of Overdose(s):

In case of accidental ingestion, seek professional advice. Symptoms of overdosage may include CNS depression with drowsiness, decreased body temperature, bradycardia, shock-like hypotension, apnea and coma.

Storage Condition(s):

Store at temperature below 30°C. Protect from light and moisture.

Shelf Life(s):

2 years from the date of manufacture.

Discard 1 month after open.

Product description & Packaging available:

A clear and colorless solution.

Plastic bottle of 5ml x 1, 5ml x 10, 5ml x 12, 5ml x 20, 5ml x 24.



Manufacturer and Product Registration Holder :

Y.S.P. INDUSTRIES (M) SDN. BHD. (192593 U)

Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan, Malaysia.

Ordering Line: 1 800 88 3027

Product Info: 1 800 88 3679

Date of Revision: 04 May 16