



Pegalate
(Macrogol Plus Electrolytes 13.72g)

PRODUCT DESCRIPTION

Before dissolve: A white crystalline powder in single-dose sachets with lemon odour.
After dissolve: Clear colourless solution

DOSAGE FORM

POWDER in form of POWDER TO BE DISSOLVED IN WATER

PHARMACODYNAMIC

Macrogol 3350 acts by virtue of its osmotic action in the gut, which induces a laxative effect. Macrogol 3350 increases the stool volume, which triggers colon motility via neuromuscular pathways. The physiological consequence is an improved propulsive colonic transportation of the softened stools and a facilitation of the defecation. Electrolytes combined with macrogol 3350 are exchanged across the intestinal barrier (mucosa) with serum electrolytes and excreted in faecal water without net gain or loss of sodium, potassium and water.

PHARMACOKINETIC

Macrogol 3350 is unchanged along the gut. It is virtually unabsorbed from the gastro-intestinal tract. Any macrogol 3350 that is absorbed is excreted via the urine.

INDICATION

It is a laxative for the treatment of chronic constipation and faecal impaction in adults, adolescents and elderly. It is not recommended for children below 12 years of age.

RECOMMENDED DOSE

Constipation:

take 1-3 sachets daily in divided dose according to the severity of your constipation. Each sachet should be dissolved in 125 ml water.

Faecal impaction:

Before you take **PEGALATE ORAL SACHET** for faecal impaction, it should be confirmed that you have this condition.

A dose of 8 sachets a day of **PEGALATE ORAL SACHET** is needed for the treatment of faecal impaction. The 8 sachets should be taken within 6 hours for up to 3 days if required. If you have a heart condition do not take more than 2 sachets in any one hour. Dissolve 8 sachets in 1 liter of water.

ROUTE OF ADMINISTRATION

Oral

CONTRAINDICATION

Intestinal perforation or obstruction due to structural or functional disorder of the gut wall, ileus, severe inflammatory conditions of the intestinal tract, such as Crohn's disease and ulcerative colitis and toxic megacolon.

Hypersensitivity to the active ingredients or to any of the excipients.

WARNING AND PRECAUTIONS

Diagnosis of impaction/faecal loading of the rectum should be confirmed by physical or radiological examination of the abdomen and rectum.

Mild adverse drug reactions are possible. If patients develop any symptoms indicating shifts of fluids/electrolytes (e.g. oedema, shortness of breath, increasing fatigue, dehydration, cardiac failure) Macrogol should be stopped immediately and electrolytes measured and any abnormality should be treated appropriately.

The absorption of other medicinal products could transiently be reduced to an increase in gastrointestinal transit rate induced by Macrogol e.g. anti-epileptics.

INTERACTIONS WITH OTHER MEDICAMENTS

Macrogol raises the solubility of medicinal products that are soluble in alcohol and relatively insoluble in water.

There is a possibility that the absorption of other medicinal products could be transiently reduced during use with Macrogol. There have been isolated reports of decreased efficacy with some concomitantly administered medicinal products, e.g. anti-epileptics.

PREGNANCY AND LACTATION

There is no experience of the use of Macrogol during pregnancy and lactation and it should only be used if considered essential by the physician.

160 x 200 mm

Front Page

ARTWORK DETAIL LABEL

Product	Pegalate - (Macrogol Plus Electrolytes 13.72g)				
Buyer/Country	UNIMED, MALAYSIA	Component	Pack Insert		
Dimension	160 x 200 mm			Pack	--
New Item Code	1035714	Old Item Code	NA		
Colour Shades	Black			No. of Colours	1
Change Control No.	NA			Artwork Version	1.1
Design/Style	Front & Back Printing. To be supplied in folded size: 80 x 50mm				
Substrate	60 GSM Paper.				
Special Instructions	Printing clarity to be clear & sharp.				
Autocartonator Requirements	NA				
Caution to the printer: Before processing, please ensure that the ARTWORK received for printing is exactly in line with APPROVED ARTWORK provided to you. In case of any FONTS/DESIGN are Mis-matching with the APPROVED ARTWORK, please inform PDC for further action. DO NOT MAKE ANY CHANGE TO THE ARTWORK WITHOUT WRITTEN INSTRUCTIONS FROM PDC.					

RD/PD/003/F-08/R0

SIDE EFFECTS

Reactions related to the gastrointestinal tract occur most commonly.

These reactions may occur as a consequence of expansion of the contents of the gastrointestinal tract, and an increase in motility due to the pharmacologic effects of Compound Macrogol. Mild diarrhoea usually responds to dose reduction.

System Organ Class	Adverse Event
Immune system disorders	Allergic reactions, including anaphylaxis, angioedema, dyspnoea, rash, urticaria, and pruritus.
Skin and subcutaneous tissue disorders	Erythema
Metabolism and nutrition disorders	Electrolyte disturbances, particularly hyperkalaemia and hypokalaemia.
Nervous system disorders	Headache
Gastrointestinal disorders	Abdominal pain,diarrhoea, vomiting, nausea, dyspepsia, abdominal distension, borborygmi, flatulence, anorectal discomfort.
General disorders and administration site conditions	Peripheral oedema

SIGN AND SYMPTOM OF OVERDOSE AND TREATMENT

Severe pain or distension can be treated by nasogastric aspiration. Extensive fluid loss by diarrhea or vomiting may require correction of electrolyte disturbances.

STORAGE CONDITION

Sachet: Do not store above 30°C.
Reconstituted solution: Store at 2 - 8°C (in a refrigerator and covered) and use within six hours.

PRESENTATION

13.72g of white crystalline powder (PEGALATE) is filled in sachet. 20 Sachets are packed in a carton with a package insert.

CONTENTS OF THE PACK

Each sachet contains:	
Polyethylene glycol	13125.00mg
Sodium Chloride	350.70 mg
Sodium Hydrogen Carbonate	178.50mg
Potassium Chloride	46.60mg

Following reconstitution in 125 ml of water, Each sachet provides:

sodium	65mmol/l(1.49mg/ml)
chloride	53mmol/l(1.88mg/ml)
hydrogen carbonate (bicarbonate)	17mmol/l (1.04mg/ml)
potassium	5.4mmol/l (0.21mg/ml)

Manufactured by:

Strides Pharma Science Limited
S-No. 36/7, Suragajakkanahalli,
Indlavadi Cross, Anekal Taluk,
Bangalore - 562 106, INDIA

Product Registration Holder:

UNIMED SDN.BHD
No. 53, Jalan Tembaga SD5/2B,
Bandar Sri Damansara,
52200 Kuala Lumpur.

Revised Date: 05/2021

160 x 200 mm



Back Page

ARTWORK DETAIL LABEL

Product	Pegalate - (Macrogol Plus Electrolytes 13.72g)				
Buyer/Country	UNIMED, MALAYSIA	Component	Pack Insert		
Dimension	160 x 200 mm			Pack	--
New Item Code	1035714	Old Item Code	NA		
Colour Shades	Black			No. of Colours	1
Change Control No.	NA			Artwork Version	1.1
Design/Style	Front & Back Printing. To be supplied in folded size: 80 x 50mm				
Substrate	60 GSM Paper.				
Special Instructions	Printing clarity to be clear & sharp.				
Autocartonator Requirements	NA				
Caution to the printer: Before processing, please ensure that the ARTWORK received for printing is exactly in line with APPROVED ARTWORK provided to you. In case of any FONTS/DESIGN are Mis-matching with the APPROVED ARTWORK, please inform PDC for further action. DO NOT MAKE ANY CHANGE TO THE ARTWORK WITHOUT WRITTEN INSTRUCTIONS FROM PDC.					