

T-Z 5% w/v Oral Suspension

T-Z 5% w/v Oral Suspension appears to be an opaque, white colour liquid.

Composition:

Each ml contains

Toltrazuril	50 mg
Sodium Benzoate (as Preservative)	2.1mg

Pharmacodynamics:

Toltrazuril is a triazinon derivative. It acts against coccidia of the genus *Isospora* and *Eimeria*. It is acting against all intracellular development stages of coccidia of the merogony (asexual multiplication) and gamogony (sexual phase). All stages are destroyed, thus the mode of action is coccidiocidal.

Pharmacokinetics:

Piglets

After oral administration toltrazuril is slowly absorbed with a bioavailability of $\geq 70\%$. The main metabolite is characterised as toltrazuril sulfone. The elimination of toltrazuril is slow with a half-life elimination time around 3 days. The major route of excretion is via the faeces.

Calves and Lambs

After oral administration in cattle toltrazuril is slowly absorbed. The main metabolite is characterised as toltrazuril sulfone. The major route of excretion is via the faeces.

Indications:

Piglets: For the prevention of clinical signs of coccidiosis in neonatal piglets on farms with a confirmed history of coccidiosis caused by *Isospora suis*.

Calves: For the prevention of clinical signs of coccidiosis and reduction of coccidia shedding in housed calves replacing cows producing milk for human consumption (dairy cows) on farms with a confirmed history of coccidiosis caused by *Eimeria bovis* or *Eimeria zuernii*.

Lambs: For the prevention of clinical signs of coccidiosis and reduction of coccidia shedding in lambs on farms with a confirmed history of coccidiosis caused by *Eimeria crandallis* and *Eimeria ovinoidalis*.

Recommended dosage:

Piglets

Individual animal treatment. Each pig to be treated on day 3-5 of life with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight.

Calves

Each animal should be treated with a single oral dose of 15 mg toltrazuril/kg body weight corresponding to 3.0 ml oral suspension per 10 kg body weight. For the treatment of a group of animals of the same breed and same or similar age, the dosing should be done according to the heaviest animal of this group.

Lambs

Each animal should be treated with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight.

Mode of Administration:

Oral administration.

Contraindication:

For environmental reasons: Do not use in calves weighing more than 80 kg bodyweight. Do not use in fattening units such as veal or beef calves

Warning and Precautions:

Frequent and repeated use of antiprotozoals from the same class may lead to the development of resistance. It is recommended to treat all animals in a pen. Hygienic measures may reduce the risk of coccidiosis. To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period. To alter the course of an established clinical coccidial infection, in individual animals already showing signs of diarrhoea, additional supportive therapy may be required.

Special precautions for use

i. Special precautions for use in animals

Not applicable.

ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wash any splashes from skin or eyes immediately with water.

iii. Other Precautions

The major metabolite of toltrazuril, toltrazuril sulfone (ponazuril), has been shown to be both persistent (half-life >1 year) and mobile in soil and to be toxic to plants.

For environmental reasons:

In order to prevent any adverse effects on plants and possible contamination of groundwater manure from treated calves must not be spread onto land without dilution with manure from untreated cows. Manure from treated calves must be diluted with at least 3 times the weight of manure from mature cows before it can be spread onto land. Lambs kept throughout the whole life span indoors under an intensive rearing system must not be treated beyond the age of 6 weeks or body weight of more than 20 kg at treatment. Manure from these animals should only be applied to the same piece of land every third year.

TO BE PRESCRIBED AND TREATED WITH BY REGISTERED VETERINARY SURGEONS ONLY.

Interactions with Other Medicaments:

None known.

Statement on Usage During Pregnancy and Lactation:

Laboratory studies in rat and rabbit have not produced any evidence of a teratogenic, foetotoxic or maternotoxic effect.

Side effects:

None known.

Overdose and Treatment:

A threefold overdose is well tolerated by healthy piglets and calves without signs of intolerance. No signs of overdose have been observed in lamb safety studies with threefold overdose at a single treatment and twofold overdose treatment on 2 consecutive days.

Withdrawal Period (s):

Piglets: Meat and offal: 77 days

Calves: Meat and offal: 63 days

Lambs: Meat and offal: 42 days

Not permitted for use in lactating animals producing milk for human consumption.

Storage Condition:

Store below 30°C. Protect from heat and sunlight exposure.

Keep out of reach of children/Jauhkan dari kanak-kanak.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Shelf life after first opening: 28 days.

Shelf life after reconstitution: 24 hours.

Packaging Available:

100ml

Date of revision:

12-06-2017

Product Holder/ Manufacturer:

NAM PHARMA SDN. BHD. (768370-H)

5, Lebuhr Perusahaan Klebang 11,
Taman Perindustrian Antarabangsa IGB,
31200 Chemor, Perak, Malaysia.

Tel: +605-2931838, 1839 Fax: +605-2931833