

PACKAGE INSERT

Product name

NW Povidone Iodine 10%w/v Solution

Name and Strength of Active Ingredient (s)

Povidone iodine.....10% w/v

Product Description

A deep brown liquid.

Pharmacodynamics

Povidone iodine is an antiseptic and disinfectants which liberates iodine in contact with the skin.

Pharmacokinetics

Povidone iodine is an iodophor with a powerful broad-spectrum germicidal activity against a wide range of bacteria, viruses, fungi, protozoa and spores.

Indication

Use to prevent and treat infections that may occur in minor scrapes and cuts.

Recommended Dosage

Adults and infants 1 month and over: No special dosage recommendation. Use medication for full time of treatment, do not use for more than ten days. Clean and dry the affected area carefully and then apply the solution 2 to 3 times a day.

Infants under 1 month: Use is not recommended.

Route of Administration

Topical

Warning and Precautions

- Avoid application of povidone iodine into the eyes and nose.
- NW Povidone Iodine 10%w/v Solution can be absorbed systemically during the topical treatment of burns and large areas of broken skin. The degree of absorption is proportional to the depth and extent of the burn or broken skin.
- Prolonged treatment with NW Povidone Iodine 10%w/v Solution of patients with severe and extensive burns may cause metabolic acidosis, hypernatraemia and renal impairment.

- Special caution is needed in patients with pre-existing renal insufficiency and in neonates and infants up to 6 months of age, pregnant and breast-feeding women. In such cases benefit/risk assessment should be performed and monitoring of thyroid function should be considered.

Interaction with Other Medicaments

Absorption of iodine from povidone iodine may interfere with thyroid function tests by causing an increase in protein bound iodine levels. Contamination with povidone iodine of several types of tests for the detection of occult blood in urine may produce false positive results.

Do not mix or co-administer with disinfectants and/or antiseptics or other agents used for the treatment of wounds. There is a risk of (transient) hypothyroidism after regular or prolonged use in patients on concomitant lithium therapy.

Pregnancy and Lactation

Pregnancy - Not recommended for use in pregnant women.

Breast-feeding – Not recommended for use in nursing mothers.

Side Effects

Povidone iodine may produce local skin reactions although it is considered to be less irritant than iodine.

The application of povidone iodine to large wounds or severe burns may produce systemic adverse effects such as metabolic acidosis, hypernatraemia and impairment of renal function.

Contraindication

Contraindicated in patients with allergic to the ingredients and thyroid disorder.

Overdose and treatment

Excess iodine can produce goitre and hypothyroidism or hyperthyroidism. Systemic absorption of iodine after repeated application of povidone iodine to large areas of wounds or burns may lead to a number of adverse effects: Metallic taste in mouth, increased salivation, burning or pain in the throat or mouth, irritation and swelling of the eyes, pulmonary oedema, skin reactions, gastrointestinal upset and diarrhoea, metabolic acidosis, hypernatraemia and renal impairment.

Deliberate or accidental ingestion of large quantities of povidone iodine will result in high blood levels of iodine and gastrointestinal toxicity including vomiting and diarrhoea. Symptomatic and supportive treatment should be started with special attention to monitoring electrolyte balance, renal function, thyroid function and liver function. Iodine can be removed by dialysis.

Effects on Ability to Drive and Use Machine

Not applicable

Instructions for use and handling, and disposal

Store on the original packaging

Any unused product or waste material should be discarded in accordance with the local requirements. No additional environment precautions for disposal are necessary.

Use this medicine only as directed

Medicines should be kept out of the reach of children.

Storage Condition

Store below 30°C in closed container.

Keep away from heat, moisture, and direct light.

Do not keep expired product.

Shelf Life

3 Years

Dosage forms and packaging available:

30ml, 50ml, 100ml, 500ml and 1000ml in Amber PET plastic bottle with white PP cap.

Name and Address of Manufacturer/Marketing Authorization Holder

Natural Wellness Industries Sdn. Bhd.

No. 78, Jalan Kilang Midah,

Taman Midah,

56000 Kuala Lumpur,

Wilayah Persekutuan Kuala Lumpur,

Malaysia

Date of Revision of Package Insert

17/07/2023