

I-Dectin 0.4% w/v Oral Solution

I-Dectin 0.4% w/v Oral Solution appears to be a yellow color, clear solution with good water miscibility.

Composition:

Each ml contains

Ivermectin	4 mg
Preservatives	
Methyl paraben	1 mg
Propyl paraben	0.1 mg

Pharmacodynamics:

Ivermectin belongs to the avermectin family, which are a macrocyclic lactone group of endectocides. Compounds of the class bind selectively and with high affinity to glutamate-gated chloride ion channels which occur in invertebrate nerve and muscle cells. This leads to an increase in the permeability of the cell membrane to chloride ions with hyperpolarization of the nerve or muscle cell, resulting in paralysis and death of the parasite. Compounds of this class may also interact with other ligand-gated chloride channels, such as those gated by the neurotransmitter gamma-aminobutyric acid (GABA).

The margin of safety for compounds of this class is attributable to the fact that mammals do not have glutamate-gated chloride channels, the macrocyclic lactones have a low affinity for other mammalian ligand-gated chloride channels and they do not readily cross the blood-brain barrier.

Pharmacokinetics:

Ivermectin binds extensively to plasma proteins. Due to its high lipophilic nature, Ivermectin is extensively distributed. It tends to accumulate in fat tissue, which acts as a drug reservoir and the highest levels of Ivermectin are found in liver and fat. Ivermectin is only partially metabolized. Ivermectin is mainly eliminated in the faeces as unaltered drug and faecal excretion accounts for 90% of the dose administered with <2% of the dose excreted in urine. Ivermectin is also excreted by the mammary gland.

Indications:

I-Dectin 0.4% w/v Oral Solution is indicated for the treatment and control of gastrointestinal roundworms, kidney worms, and lungworms in pigs. Ivermectin in water at the recommended dose level, it provides effective control of the following parasites of pigs:

Gastrointestinal Roundworms: Adults and fourth stage larvae of large roundworm (*Ascaris suum*), small brown stomach worm (*Hyostrogylus rubidus*), nodule worm (*Oesophagostomum* spp.) and adults of stomach worm (*Ascarops strongylina*) and intestinal threadworm (*Strongyloides ransomi*).

Kidney Worms: Adults and fourth stage larvae of *Stephanurus dentatus*.

Lungworms: Adult *Metastrongylus* spp.

Mode of Administration:

Oral Administration in drinking water.

Recommended dosage:

Growing pigs and Adult pigs including Sows and Boars:

This product is suitable for mixing with water.

The recommended dose level is 2.5mL I-DECTIN 0.4% W/V ORAL SOLUTION per 100kg liveweight (corresponding to 100µg/kg liveweight) daily for 7 days.

Instruction for use:

1. Generally, treated pigs should not be moved to clean quarters or exposed to infected pigs for at least one week after completion of treatment.
2. Pigs exposed to contaminated premises, soil or pasture may need re-treatment if reinfection occurs.

RECOMMENDED TREATMENT PROGRAMS

Breeding Animals

At the time of initiating any parasite control program, it is important to treat all animals in the herd. After the initial treatment, use I-Dectin 0.4% w/v Oral Solution regularly as follows:

Sows: Treat preferably 14-21 days prior to farrowing to minimise infection of piglets.

Gilts: Treat 14-21 days prior to breeding. Treat 14-21 days prior to farrowing.

Boars: Treat at least 2 times per year.

Feeder Pigs (Finishers): All pigs received for finishing should be treated upon placement in clean quarters.

Contraindication:

None known.

Warning and Precautions:

Poisonous if swallowed. May irritate the eyes, avoid contact with eyes. Wash hands after use. If poisoning occurs contact a doctor immediately.

Interactions with Other Medicaments:

No information available.

Statement on Usage During Pregnancy and Lactation:

The product can be administered to sow during pregnancy and it can be used in breeding animals following the advice under RECOMMENDED TREATMENT PROGRAMS in this leaflet. No known information during lactation is available.

Side effects:

None known.

Overdose and Treatment:

No known information on overdose and treatment is available.

Withdrawal Period (s):

MEAT: Remove all medicated water 7 days before slaughter for human consumption.

Storage Condition:

Store below 30°C. Protect from heat and sunlight exposure.

Keep out of reach of children/ Jauhkan dari kanak-kanak.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Shelf life after first opening: 28 days.

Shelf life after reconstitution: 24 hours.

Packaging Available:

100 ml & 1L

Date of revision:

10-10-2017

Product Holder/ Manufacturer:

NAM PHARMA SDN. BHD. (768370-H)

5, Lebuhr Perusahaan Klebang 11,

Taman Perindustrian Antarabangsa IGB,

31200 Chemor, Perak, Malaysia.

Tel: +605-2931838, 1839 Fax: +605-2931833