

Veterinary Package Insert

TP-FLOR 250 MG/ML SOLUTION

PRODUCT DESCRIPTION

Light yellow to yellow colour liquid, which contains florfenicol 250mg in 1ml of solution (25% in w/v).

PHARMACODYNAMICS & PHARMACOKINETICS

Florfenicol is a broad-spectrum synthetic antibiotic in the phenicol group that is active against most Gram-positive and Gram-negative bacteria isolated from domestic animals. Florfenicol acts by inhibition of protein synthesis at the ribosomal level and is bacteriostatic.

After administration to pigs by gavage at 15 mg/kg under experimental conditions, absorption of florfenicol was variable but peak serum concentrations of approximately 5 µg/ml were reached approximately 2 hours after dosing. The terminal half-life was between 2 and 3 hours.

After absorption and distribution, florfenicol is extensively metabolised by pigs and rapidly eliminated, primarily in urine.

INDICATION

In pigs:

Treatment and prevention at the group level where clinical signs are present of swine respiratory disease associated with Actinobacillus pleuropneumoniae and Pasteurella multocida susceptible to florfenicol. The presence of the disease should be established in the herd before initiating preventive treatment.

RECOMMENDED DOSAGE AND ADMINISTRATION

To be administered in drinking water.

10 mg florfenicol per kg bodyweight per day in drinking water for 5 consecutive days.

Based on the recommended dose, and the number and weight of the animals to be treated, the exact daily amount of the veterinary product should be calculated according to the following formula:

X ml veterinary product / kg b.w./day

x

Mean body weight (kg) of animals to be treated

=

X

ml veterinary product per litre drinking water

Mean daily water consumption (litre) per animal

The appropriate quantity of medicated water should be prepared based on the daily water consumption. To ensure a correct dosage body weight should be determined as accurately as possible. In order to avoid under- and over-dosing, treated animals should be divided into groups of similar bodyweight and the dose should be calculated for each group individually.

For Bulk Tank:

To treat pigs drinking 10% of their bodyweight, at the dose of 10 mg/kg: add the florfenicol solution to the drinking water in the bulk tank. Use 200ml of the product for every 500 L of water, 400ml of product for every 1000 L of water or use 2L of product for every 5000 L of water and mix thoroughly.

CONTRAINDICATIONS

Do not use in boars intended for breeding purposes.

Do not use in case of hypersensitivity to the active substance or to the excipient.

WARNINGS & PRECAUTIONS.

Special warnings for each target species

The treated pigs should be placed under special observation. On each of the five days of treatment, unmedicated drinking water should not be given until the full daily amount of medicated drinking water has been ingested by pigs.

If there are no signs of improvement after three days of treatment, the diagnosis should be reviewed and, if necessary, the treatment changed.

Special precautions for use

Special precautions for use in animals

The veterinary medicinal product should be used in conjunction with susceptibility testing.

Use of the product deviating from the instructions given here may increase the prevalence of bacteria resistant to the florfenicol.

Official and local antimicrobial policies should be taken into account when the product is used.

Treatment should not exceed 5 days.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

People with known hypersensitivity to florfenicol or propylene glycol should avoid contact with the veterinary medicinal product.

In case of accidental spillage onto skin rinse with water. In case of contact with eyes, rinse immediately with copious amounts of water.

Personal protective goggles should be worn when handling the veterinary medicinal product.

Other precautions

In order to prevent any adverse effects on algae and possible contamination of groundwater, manure from treated pigs must not be spread onto land without dilution with manure from untreated pigs. Manure from treated pigs must be diluted with at least 5 times the weight of manure from untreated pigs before it can be spread onto arable land.

INTERACTIONS WITH OTHER MEDICATIONS

Data not available.

PREGNANCY AND LACTATION

Studies in laboratory animals have not revealed any evidence of potential embryotoxic or foetotoxic effect of florfenicol.

The safety of the veterinary medicinal product in sows has not been established during pregnancy and lactation.

The use is not recommended during pregnancy and lactation.

ADVERSE EFFECTS

A slight reduction of water consumption by the animals, dark brown faeces and constipation may be observed during treatment.

Commonly observed adverse effects are diarrhoea and/or peri-anal and rectal erythema/oedema which may affect approximately 40% of the animals. These effects are transient. In a few of the affected animals, prolapse of the rectum, that resolves without treatment may be observed.

OVERDOSE & TREATMENT

In case of overdosing, a decrease in weight gain, food and water consumption, peri-anal erythema and oedema and modification of some haematological and biochemical parameters indicative of dehydration may be observed.

WITHDRAWAL PERIOD

Meat and offal: 20 days

MAXIMUM RESIDUAL LIMIT (MRL)

Substance	Food	Maximum Residue Limits (MRLs) in µg/kg

Florfenicol	Pig, muscle	300
	Pig, skin & fat	500
	Pig, liver	2000
	Pig, kidney	500

STORAGE CONDITION

Store in a dry, cool place and avoid direct sunlight. It should be store at room temperature (30°C or below.)

SHELF LIFE

2 years.

After 1st opening, reconstitution or dilution, use within 24 hours.

PACKING

1 L

THYE PHARMA SDN. BHD.

(manufacturer & product registration holder)
No.2 Lorong Industri Ringan Permatang
Tinggi 8,
Kawasan Industri Ringan Permatang Tinggi,
14000 Bukit Mertajam,
Pulau Pinang.

Revised October 2017