

YH BACITRACIN ZINC 15%W/W GRANULE

ACTIVE CONSTITUENT: Bacitracin Zinc 750 mg/g (As bacitracin 150mg). 1mg bacitracin equivalent to 42iu.

PRODUCT DESCRIPTION: Light brown or tan granule.

ROUTE OF ADMINISTRATION: Oral administration.

INDICATIONS:

For the treatment of Necrotic Enteritis (*Clostridium perfringens* Types A and C) and as an aid in the prevention of Necrotic Enteritis in poultry.

RECOMMENDED DOSAGE:

Should be premixed with 10-20 kg of feed and then thoroughly mixed with the bulk feed.

Aid in the prevention of Necrotic Enteritis: Add YH Bacitracin Zinc 15% W/W Granule at the rate of 267 g/tonne of finished feed (corresponding to 40g bacitracin activity per tonne).

Treatment of Necrotic Enteritis: Add YH Bacitracin Zinc 15% W/W Granule at the rate of 1.33kg/tonne of finished feed (corresponding to 200g bacitracin activity per tonne) to be fed for 5-7 days.

PHARMACODYNAMICS & PHARMACOKINETIC:

Bacitracin is an antibiotic from the group of peptide antibacterial compounds. Bacitracin is produced by *Bacillus lincheniformis*, and is a mixture of several closely related polypeptides, mainly consisting of bacitracin A, B1, B2, C and a small portion of bacitracin F. Bacitracin is mainly active against gram-positive microorganisms. Bacitracin inhibits cell wall biosynthesis by inhibiting lipid pyrophosphatase which is involved in transmembrane transport of peptidoglycan precursors. After oral application to chickens and pigs, bacitracin is hardly absorbed from the gastrointestinal tract, and the distribution to organs and tissues is negligible. In chickens and pigs, approximately 95% of an oral dose is excreted via faeces, and only 3% or less via urine. Bacitracin is metabolized to amino acids and smaller peptides via the main metabolite desamido-bacitracin, which is microbiologically inactive.

WARNINGS AND PRECAUTIONS:

Any variation by prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may require extending the approved withholding period.

INTERACTION WITH OTHER MEDICINAL PRODUCT:

Bacitracin zinc can be used in combination with: (i) Amprolium and ethopabate; (ii) Clopidol; (iii) Decoquinat; (iv) Lasalocid; (v) Monensin; (vi) Naracin; (vii) Nitarsone; (viii) Robenidine; (ix) Salinomycin.

PREGNANCY AND LACTATION: None known

OVERDOSE AND TREATMENT: Stop medication and to seek for veterinarian advice if there is sign of over dosage.

STORAGE CONDITION: Store in room temperature (30°C) and dry place. Avoid light and close tightly.

CONTRAINDICATION: None Known

SIDE EFFECTS: None known.

WITHDRAWAL PERIOD: Meat: Nil, Eggs: Nil.

PACKAGING: 25kg/bag.

ADVERSE EFFECTS/UNDESIRABLE EFFECTS: None known.

SHELF LIFE AFTER RECONSTITUTION OR DILUTION: 6 months

DATE OF REVISION: August 2017

SHELF LIFE: 2 years

MANUFACTURED BY:

Lifecome Biochemistry Co., Ltd

Add.: No.1, Putan Avenue, Putan Industrial Park,
Pucheng, Fujian, China.

MARKETING AUTHORISATION HOLDER:

Yenher Agro-Products Sdn.Bhd.,

Add.: 1628, Jalan IKS Simpang Ampat1. Taman IKS Simpang Ampat,
14100 Simpang Ampat. Seberang Perai Selatan, Penang , Malaysia