

Pharmacotherapeutic group: Preparations with salicylic acid derivatives
ATC code : M02AC

Antiphlamine[®] S Lotion

- **ANTIPHLAMINE S LOTION** is containing Liquid lanolin, special base, so, it is absorbed into skin smoothly and rapidly.
- **ANTIPHLAMINE S LOTION** does not remain oily so, it does not easily stick to clothes.
- **ANTIPHLAMINE S LOTION** is a smooth lotion form so, it can be applied easily to curved body parts.
- **ANTIPHLAMINE S LOTION** can be used conveniently with one touch cap and also show strong massage effect by using acupressure container.

[COMPOSITION] 1g of this drug contains
Methyl salicylate ----- 200mg
L-Menthol ----- 60mg

[DESCRIPTION]
White homogenous lotion with menthol odor

[INDICATION]
For the temporary relief of minor aches and pains of muscles, simple backaches, strains and sprains.

[DOSAGE • ADMINISTRATION]
Apply a proper quantity to the affected area 1 ~ 4 times daily.
* Gently massage after this drug is sufficiently absorbed into skin and dry.
* Be cautious with using the acupressure container.
Hard handed massage with the acupressure container may easily damage skin and clothes.

[PHARMACOLOGICAL EFFECTS]
Menthol dilates the blood vessels causing a sensation of coldness, followed by an analgesic effect. Menthol also acts as a penetration enhancer, increasing the penetration of drugs when applied on the skin, to give a faster onset of action. Salicylates inhibit cyclooxygenase, thereby reducing the formation of prostaglandins, and cause platelet dysfunction. Methyl salicylate is used topically as a counter-irritant. Upon application, it is absorbed through the skin and is applied for the relief of pain in rheumatic conditions and painful muscle or joints.

[PHARMACOKINETICS]
The product is for topical administration. No absorption studies have been carried out with this product. Like other salicylates, methyl salicylate may be absorbed through intact skin.

- [PRECAUTION]**
1. Contraindication
 - Avoid use on eyes, near eyes and mucous membranes.
 - Dermatitis caused by eczema or rhus verniciflua, wounded area, sensitive skin
 - Hypersensitivity to Menthol / Methyl Salicylate is a contraindication
 2. Warnings
 - Patients with a history of allergy to drug preparation or cosmetics (example dermatitis caused by rash, itching etc)
 - Patients who or whose families have allergic constitutions.
 - Patients with severe wetting or erosion
 - Patients under the care of a doctor
 - Patients with diabetes or circulatory disorder

- Pediatric patients below 12 years of age.
- This product contains methyl salicylate and when applied or rub on the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding.
- Do not cover the applied area with a bandage. Do not expose to heat or hot water.
- This product should not be used in children under 2 years of age.
- Do not use during pregnancy or when breastfeeding.
- Use with caution in older children.
- If you are taking warfarin, please consult your doctor before using this product.

3. Side effect
If symptoms such as rash, flare, itching appear with this preparation, discontinue the use and notify a doctor or pharmacist.


4. Pregnancy and Lactation
There are no adequate studies done on menthol and methylsalicylate to determine safe and effective use in pregnant women. It is not known whether menthol and methylsalicylate enters breast milk; therefore, it is best to be cautious before using it in nursing mothers.

5. Driving and machines: None known.


6. Overdose: Not applicable

7. Drug interaction
Some medicines may interact with methyl salicylate/menthol lotion. Tell your health care provider if you are taking any other medicines, especially any of the following : Anticoagulants (eg, warfarin) because the risk of their side effects may be increased by methyl salicylate/menthol lotion. This may not be a complete list of all interactions that may occur. Ask your health care provider if methyl salicylate/menthol lotion may interact with other medicines that you take. Check with your health care provider before you start, stop, or change the dose of any medicine.


8. Caution on storage



1) Keep out of the reach of children.



2) Avoid direct sunlight and store in a tight container. If possible, store under dry and cool condition.



3) To prevent misuse and keep drug quality, don't change the original container

[STORAGE CONDITION]
Store below 30°C. Avoid direct sunlight and store in a tight container.

[SHELF LIFE] 3 years from the manufacturing date

[PACKAGE] 100mL/bottle

[MANUFACTURER]
Genuone Sciences Inc. / 245, Sandan-gil, Jeonui-myeon, Sejong-si, Korea For Yuhan Corporation

[PRODUCT REGISTRATION Holder]
Registration No.: MAL17106056XC
Eucogen Pharma Sdn Bhd, / 6A, Jalan Sungai Burung U 32/U, Bukit Rimau, 40460 Shah Alam, Selangor, Malaysia.

[SINGAPORE IMPORTER]
Euro Asia Medico Pte Ltd, 6 Tagore Drive, #04-13 Tagore Building, Singapore 787623.

[BRUNEI IMPORTER]
NOVA Borneo Company Sdn Bhd, Unit 4GF Bangunan Zainuddin & Azizah, Simpang 501 Jalan Tutong, Kampong Telanai, Brunei-Muara BA2312, Brunei Darussalam.

[Date of revision]
7 June 2022