Jewim Minoxidil Topical Solution

Product Name

Jewim Minoxidil Topical Solution 2% w/v.

Jewim Minoxidil Topical Solution 5% w/v.

Composition

Jewim Minoxidil Topical Solution 2% w/v contains minoxidil, at a concentration of 20mg minoxidil per mL in a solution composed of ethanol 96%, propylene glycol and purified water.

Jewim Minoxidil Topical Solution 5% w/v contains minoxidil, at a concentration of 50mg minoxidil per mL in a solution composed of ethanol 96%, propylene glycol and purified water.

Description

It is a colourless to yellow cutaneous solution of minoxidil in a suitable ethanolic solvent.

Pharmacodynamics/Pharmacokinetics

Pharmacodynamics

When applied topically, Jewim Minoxidil Topical Solution(minoxidil) has been shown to stimulate hair growth in males with alopecia androgenetica. The onset of hair growth stimulation occurs approximately after 4 (2% w/v)/2 (5% w/v) or more months of use and is variable among patients. Upon discontinuation of treatment with Jewim Minoxidil Topical Solution, new hair growth stops and restoration of pretreatment appearance may occur within 3-4 months.

The exact mechanism of action of Jewim Minoxidil Topical Solution in the treatment of alopecia androgenetica is not known. Topical application of Jewim Minoxidil Topical Solution showed no systemic effects related to absorption of minoxidil when tested in controlled clinical trials in both normotensive and untreated hypertensive patients.

In the treatment of hypertension, orally administered minoxidil has a direct peripheral vasodilator effect which reduces elevated systolic and diastolic blood pressure by decreasing peripheral vascular resistance. Reduction of peripheral arteriolar resistance and the associated fall in blood pressure induces sympathetic, vagal inhibitory, and renal homeostatic mechanisms, including an increase in renin secretion, which lead to increased heart rate and cardiac output, and salt and water retention. Minoxidil does not interfere with vasomotor reflexes and therefore does not produce orthostatic hypotension. In experimental animals, the drug does not enter the central nervous system (CNS) in significant amounts.

Pharmacokinetics

Following topical application, minoxidil is poorly absorbed from normal intact skin, with an average of approximately 1.4% (range 0.3-4.5%) of the total applied dose ultimately reaching the systemic circulation. This compares to essentially complete absorption from the gastrointestinal tract following oral administration of minoxidil tablets. Therefore, a 1-mL dose of Jewim Minoxidil Topical Solution, delivering 20 mg (2% w/v)/50 mg (5% w/v) minoxidil to the skin, would result in absorption of approximately 0.28 mg (2% w/v)/0.7 mg (5% w/v) minoxidil. The effects of concomitant dermal diseases on absorption are unknown.

Serum minoxidil levels resulting from topical administration of Jewim Minoxidil Topical Solution are governed by the drug's percutaneous absorption rate. Following cessation of topical dosing of Jewim Minoxidil Topical Solution approximately 95% of systemically absorbed minoxidil is eliminated within 4 days. The metabolic biotransformation of minoxidil absorbed following topical application has not been fully determined. The active form of the drug appears to be a sulfated metabolite, minoxidil sulfate. Orally administered minoxidil is metabolised predominantly by conjugation with glucuronic acid at the N-oxide position in the pyrimidine ring but also by conversion to more polar products. Known metabolites exert much less pharmacologic effect than minoxidil itself. Minoxidil does not bind to plasma proteins and its renal clearance corresponds to the glomerular filtration rate. Minoxidil does not cross the blood brain barrier. Minoxidil and its metabolites are haemodialysable, and are excreted principally in the urine.

Indication

Jewim Minoxidil Topical Solution is exclusively indicated for the treatment of alopecia androgenetica in males and females. Jewim Minoxidil Topical Solution stimulates hair growth and stabilizes hair loss in patients with alopecia androgenetica.

Dosage& Administration

For external use only. Use only as directed. Do not apply to any area of the body other than the scalp.

A total dose of 1mL should be applied twice per day to the scalp, beginning at the centre of the affected area. This dose should be used regardless of the size of the affected area. The total daily dose should not exceed 2mL. After application wash hands thoroughly. Apply when the hair and scalp are thoroughly dry. Do not use a hairdryer to speed the drying of the solution because blowing air on the scalp may decrease the effectiveness of Jewim Minoxidil Topical Solution.

The method of application:

Extended Spray-Tip Applicator: This applicator works best for applying Jewim Minoxidil Topical Solution to small areas of the scalp, or under hair. The pump spray applicator must be on the bottle to use the extended tip applicator. Remove large outer cap; remove small inner cap and discard it. Insert the pump spray applicator into bottle and screw on firmly. Remove small spray head from top of pump spray applicator; fit the extended spray tip applicator onto the spray shaft and push down firmly. Remove the small cap on the end of the extended tip. After aiming the applicator toward the center of the bald area of the scalp, press the pump once and spread Jewim Minoxidil Topical Solution with fingertips to cover all of the bald area. Repeat for a total of 6 times, to apply a dose of 1mL of solution. Avoid breathing spray mist. If desired, replace the small cap on end of extended tip when not in use.

Clinical experience indicates that twice daily application for ≥4 months may be required before evidence of hair regrowth stimulation can be expected. Onset and degree may be variable among patients. Relapse to pretreatment appearance following discontinuation of medication has been reported to occur within 3-4 months.

Contraindications

Topical minoxidil is contraindicated in patients with a history of hypersensitivity to minoxidil, propylene glycol or ethanol.

Warnings and Precautions

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Before starting on Jewim Minoxidil Topical Solution the patient should have a healthy, normal scalp.

Although the following effects have not been associated with topical use of Jewim Minoxidil Topical Solution, there is some absorption of minoxidil from the skin and the potential exists for systemic effects, eg, tachycardia, angina, edema or potentiation of the orthostatic hypotension produced by guanethidine. Patients should be observed periodically for any suggestion of systemic effects of minoxidil. In the event of systemic side effects discontinue administration of the drug. If necessary, fluid retention and edema can be managed with diuretic treatment. Tachycardia and angina can be controlled by administration of β-adrenergic blocking drugs or other sympathetic nervous system suppressants. Patients should discontinue use of Jewim Minoxidil Topical Solution and contact their physician in the event of systemic effects or severe dermatologic reactions.

Jewim Minoxidil Topical Solution will cause burning and irritation of the eye. In the event of accidental contact with sensitive surfaces (eye, abraded skin, mucous membranes), the area should be bathed with copious amounts of cool tap water.

Accidental ingestion of Jewim Minoxidil Topical Solution could lead to serious adverse effects. The effects in patients with concomitant skin diseases, or in those using topical corticosteroids or other dermatological preparations, are unknown. It has not been clearly determined whether occlusion (use of wigs) will increase the absorption of minoxidil after administration of Jewim Minoxidil Topical Solution.

Interactions

There are currently no known drug interactions associated with the use of Jewim Minoxidil Topical Solution. Although it has not been clinically demonstrated, there exists the theoretical possibility of absorbed minoxidil potentiating orthostatic hypotension in patients currently taking guanethidine.

Use in pregnancy & lactation

The effects of Jewim Minoxidil Topical Solution in pregnancy are unknown.

Systemically absorbed minoxidil is excreted in human milk.

Jewim Minoxidil Topical Solution should not be used by pregnant or nursing women.

Use in Children

Safety and efficacy of Jewim Minoxidil Topical Solution in patients under 18 years of age have not been established.

Use in elderly

Safety and efficacy of Jewim Minoxidil Topical Solution in patients over 65 years of age have not been established.

Adverse Effects

The most frequently reported adverse effects in clinical trials with Jewim Minoxidil Topical Solution were minor dermatological reactions. Local irritation was the most common adverse reaction reported, including scaling, erythema/flushing, dermatitis, dry skin, hypertrichosis (In areas other than where Jewim Minoxidil Topical Solution was applied), burning sensation and rash. Infrequently reported adverse reactions included allergic reactions (sensitivity, hives, generalized erythema and facial swelling); dizziness; tingling sensation; headache; weakness; neuritis; edema; eye irritation; altered taste; ear infection (particularly otitis externa) and visual disturbances. Rarely reported adverse reactions included alopecia, hair abnormalities, chest pain, blood pressure changes, pulse changes, hepatitis and kidney calculi.

Overdosage

Accidental ingestion may produce systemic effects related to the vasodilatory action of the drug (5 mL of Jewim Minoxidil Topical Solution 2% w/v contains 100 mg minoxidil, the maximum recommended adult oral dose for treatment of hypertension). Signs and symptoms of minoxidil overdosage would most likely be cardiovascular effects associated with fluid retention, lowered blood pressure and tachycardia. Fluid retention can be managed with appropriate diuretic therapy. Tachycardia can be controlled by administration of a β-adrenergic blocking agent. Hypotension should be treated by intravenous administration of normal saline. Sympathomimetic drugs, such as noradrenaline and adrenaline should be avoided because of their excessive cardiac stimulating activity.

Storage

Store below 30°C and keep it in airtight container.

Dosage Forms and packaging available

Jewim Minoxidil Topical Solution 2% w/v, containing 20 mg minoxidil per mL is available in a 60 mL bottle.

Jewim Minoxidil Topical Solution 5% w/v, containing 50 mg minoxidil per mL is available in a 60 mL bottle.

Name and Address of Manufacturer

Jewim Pharmaceutical (Shandong) Co., Ltd.
No. 5 Leigushi Street,
Taian Hi-Tech Industrial Development Zone,
Shandong Province 271000, PRC

Product Registration Holder

Medispec (M) Sdn Bhd
55 & 57, Lorong Sempadan 2,
11400 Ayer Itam, Penang,
Malaysia

Date of Revision

11/4/2022

10050623/02

10050623/02　　1. 尺寸：19×26CM　　2.材质：70g双胶纸，正反面印刷，黑色
3.颜色以20211231签注的鲁信天一色卡编号为“说明书：5%*60ml米诺地尔喷雾剂-MY0104”为准

物料编码/版本	10050623/02	物料说明	米诺地尔喷雾剂说明书(MY0104).5%×60ml		
尺　寸	19×26CM	材　质	70g双胶纸，正反面印刷，黑色	工艺	无
条码类型	无	颜色	颜色以20191016签注的鲁信天一色卡编号为“说明书：5%×60ml米诺地尔喷雾剂-MY0104”为准		
是否为新产品	否	打码空间要求	无		
变化点	Middle of Chuangye Main Street 改为No.5 leigushi Street China 改为PRC				
其它说明	无	设 计 人		变更管理员	

注：本设计图颜色仅供参考。