

hovid Ofloxacin Eye Drops 0,3% w/v

VIOFL02-0

DESCRIPTION

Clear, light yellow or yellowish green solution.

COMPOSITION

Each 8 mL contains Ofloxacin 24 mg.

ACTIONS AND PHARMACOLOGY

Ofloxacin is a synthetic fluorinated 4-quinolone antibacterial agent with activity against a broad spectrum of Gram negative and to a lesser degree Gram positive organisms.

Ofloxacin has been shown to be active against most strains of the following organisms both *in vitro* and clinically in ophthalmic infections.

Gram-negative bacteria: *Acinetobacter calcoaceticus* var. *anitratum*, and *A. calcoaceticus* var. *iwofii*; *Enterobacter* sp. including *E. cloacae*; *Haemophilus* sp. including *H. influenza* and *H. aegyptius*; *Klebsiella* sp., including *K. Pneumoniae*; *Moraxella* sp., *Morganella morganii*; *Proteus* sp., including *P. Mirabilis*; *Pseudomonas* sp.; including *P. Aeruginosa*, *P. cepacia*, and *P. fluorescens*; and *Serratia* sp., including *S. marcescens*.

Gram-positive bacteria: *Bacillus* sp.; *Corynebacterium* sp.; *Micrococcus* sp.; *Staphylococcus* sp., including *S. aureus* and *S. epidermidis*; *Streptococcus* sp., including *S. Pneumoniae*, *S. viridans* and Beta-haemolytic.

The primary mechanisms of action is through inhibition of bacterial DNA gyrase, the enzyme responsible for maintaining the structure of DNA.

Ofloxacin is not subject to degradation by beta-lactamase enzymes nor is it modified by enzymes such as aminoglycoside adenylases or phosphorylases, or chloramphenicol acetyltransferase.

PHARMACOKINETICS

After ophthalmic instillation, ofloxacin is well maintained in the tear-film.

In a healthy volunteer study, mean tear film concentrations of ofloxacin measured four hours after topical dosing (9.2 µg/g) were higher than the 2 µg/ml minimum concentration of ofloxacin necessary to inhibit 90% of most ocular bacterial strains (MIC90) *in-vitro*.

Maximum serum ofloxacin concentrations after ten days of topical dosing were about 1000 times lower than those reported after standard oral doses of ofloxacin, and no systemic side-effects attributable to topical ofloxacin were observed.

INDICATION

hovid Ofloxacin Eye Drops 0,3% w/v is indicated for the topical treatment of external ocular infections (such as conjunctivitis and keratoconjunctivitis) in adults and children caused by ofloxacin-sensitive organisms.

CONTRAINDICATIONS

hovid Ofloxacin Eye Drops 0,3% w/v is contra-indicated in individuals who have shown hypersensitivity to ofloxacin, any of its excipients or any other quinolones.

WARNINGS AND PRECAUTIONS

- Safety and effectiveness in infants below the age of one year have not been established.
- Serious and occasionally fatal hypersensitivity (anaphylactic/anaphylactoid) reactions, some following the first dose, have been reported in patients receiving systemic quinolones, including ofloxacin. Some reactions were accompanied by cardiovascular collapse, loss of consciousness, angioedema (including laryngeal, pharyngeal or facial oedema), airway obstruction, dyspnoea, urticaria, and itching.
- If an allergic reaction to Ofloxacin Eye Drops occurs, discontinue the drug. Use Ofloxacin Eye Drops with caution in patients who have exhibited sensitivities to other quinolones antibacterial agents.
- When using Ofloxacin Eye Drops, the risk of rhinopharyngeal passage which can contribute to the occurrence and the diffusion of bacterial resistance should be considered. As with other anti-infectives, prolonged use may result in overgrowth of non-susceptible organisms.
- If worsening infection occurs, or if clinical improvement is not noted within a reasonable period, discontinue use and institute alternative therapy.
- Stevens-Johnson syndrome has been reported in patients receiving topical ophthalmic ofloxacin, however, a causal relationship has not been established.
- **Cardiac disorders:** Caution should be taken when using fluoroquinolones, including Ofloxacin Eye Drops in patients with known risk factors for prolongation of the QT interval such as, for example:
 - congenital long QT syndrome
 - concomitant use of drugs that are known to prolong the QT interval (e.g. Class IA and III anti-arrhythmics, tricyclic antidepressants, macrolides, antipsychotics)
 - uncorrected electrolyte imbalance (e.g. hypokalaemia, hypomagnesaemia)
 - cardiac disease (e.g. heart failure, myocardial infarction, bradycardia)Elderly patients and women may be more sensitive to QTc-prolonging medications. Therefore, caution should be taken when using fluoroquinolones, including Ofloxacin Eye Drops in these populations.
- Use Ofloxacin Eye Drops with caution in patients who have exhibited sensitivities to other quinolone antibacterial agents.

- The use of Ofloxacin Eye Drops in neonates with ophthalmia neonatorum caused by *Neisseria gonorrhoeae* or *Chlamydia trachomatis* is not recommended as it has not been evaluated in such patients.
- It is necessary to advise caution regarding the risk of corneal perforation when using product to treat patients with corneal epithelial defects or corneal ulcers.
- Corneal precipitates have been reported during treatment with topical ophthalmic ofloxacin. However, a causal relationship has not been established.
- Ofloxacin Eye Drops contains the preservative benzalkonium bromide which may cause ocular irritation and discolour soft contact lenses.
- Sun or UV-exposition should be avoided during use of ofloxacin due to the potential for photosensitivity.
- Use of contact lenses is not recommended in patients receiving treatment for an eye infection.
- **Effects on ability to drive:** No studies on the effects on the ability to drive and use machines have been performed. Transient blurring of vision may occur on instillation of eye drops. Do not drive or operate hazardous machinery unless vision is clear.

PREGNANCY AND LACTATION

Pregnancy: There have been no adequate data, it is recommended that Ofloxacin Eye Drops not be used in pregnant women.

Lactation: Because ofloxacin and other quinolones taken systemically are excreted in breast milk, and there is potential for harm to nursing infants, a decision should be made whether to temporarily discontinue nursing or not to administer the drug, taking into account the importance of the drug to the mother.

DRUG INTERACTIONS

Metabolic clearance of caffeine and theophylline are not significantly affected by systemic ofloxacin.

It has not been reported that with the concomitant systemic use of NSAIDs and ofloxacin will cause an increased prevalence of CNS toxicity.

Ofloxacin Eye Drops, like other fluoroquinolones, should be used with caution in patients receiving drugs known to prolong the QT interval (e.g. Class IA and III anti-arrhythmics, tricyclic antidepressants, macrolides, antipsychotics).

MAIN SIDE/ADVERSE EFFECTS

General: Serious reactions after use of systemic ofloxacin are rare and most symptoms are reversible.

hovid Ofloxacin Eye Drops 0.3% w/v

Since a small amount of ofloxacin is systemically absorbed after topical administration, side-effects reported with systemic use could possibly occur.

Immune System Disorders

Not Known: Hypersensitivity reaction including signs or symptoms of Eye allergy (such as Eye pruritus and Eyelid pruritus) and Anaphylactic reactions (such as angioedema, dyspnea, anaphylactic shock, oropharyngeal swelling, facial oedema and tongue swollen)

Nervous System Disorders

Not known: Dizziness

Eye Disorders

Common: Eye irritation; Ocular discomfort
Not known: Keratitis; Conjunctivitis; Vision blurred; Photophobia; Eye oedema; Foreign body sensation in eyes; Lacrimation increased; Dry eye; Eye pain; Ocular hyperaemia; Periorbital oedema (including eyelid oedema)

Cardiac disorders

Not known: ventricular arrhythmia and torsades de pointes (reported predominantly in patients with risk factors for QT prolongation); ECG QT prolonged

Gastrointestinal Disorders

Not known: Nausea

Skin and Subcutaneous Tissue Disorders

Not Known: Stevens-Johnson syndrome; Toxic epidermal necrolysis

OVERDOSE AND TREATMENT

In the event of overdose, symptomatic treatment should be implemented, ECG monitoring should be undertaken, because of the possibility of QT interval prolongation.

DOSAGE AND ADMINISTRATION

Topical ocular instillation.

Usually, instill 1 drop a time 3 times daily. The dosage may be adjusted according to the patient's symptoms.

Note: The information given here is limited. For further information consult your doctor or pharmacist.

Storage: Store below 30°C. Protect from light. It is recommended to discard remaining contents one month after opening.

Presentation/Packing: Eye Drops of 8 mL, 1 bottle per box.

Manufactured for: **HOVID Bhd.**

121, Jalan Tunku Abdul Rahman, 30010 Ipoh, Malaysia.

Date of information: May 2017