

# PANFAST SALINOMYCIN 12% W/W GRANULE

## Composition:

Salinomycin Sodium.....120mg/g

Exp. Qs.....25kg

## Description

Pale brown, homogeneous granular

## Pharmacodynamic

The biological activity of Salinomycin is due to its ability to form complexes with cations, particularly its ability to transport potassium ions into and out of the cell. The resulting disturbance in the parasite's intercellular ion concentration explains the coccidiocidal activity.

During asexual development of the coccidian, Salinomycin has a coccidiocidal effect on sporozoties and merozoties. Subsequent development of schizonts is also severely impaired, resulting in a minimal damage of the intestine.

## Pharmacokinetic

The drug is rapidly and well (after administration into the crop) absorbed.

Salinomycin is distributed and eliminated quickly. The serum concentration kinetic most closely matches a two-compartment model with a distribution phase, which ends after one hour. Since the active ingredient is lipophilic and can readily diffuse across the cell membranes, the distribution of salinomycin in the tissue is greater than in the serum. The highest concentrations are in liver, followed by kidney, muscle, fat, heart and skin found. In the tissue, the active ingredient may after 48 h no longer be detected, except in the liver where traces to 72 h after application can be measured.

## Indication

For use in broiler chickens as an aid in the prevention of coccidiosis caused by *Eimeria acervulina*, *E. mitis*, *E. necatrix*, *E. maxima*, *E. tenella* and *E. brunetti*.

## Route of Administration

Orally, for use in complete feed only.

## Recommended dose

Broiler Chickens: Thoroughly mix 500 g of PANFAST SALINOMYCIN 12% W/W GRANULE with 999.5 kg of complete feed to obtain a medicated feed with a salinomycin sodium concentration of 0.0060% (60 mg/kg). The medicated feed can be prepared in the form of meal or pellets.

The medicated feed is to be fed continuously as the sole ration up to marketing.

## Contraindication

Do not allow turkeys, dogs or horses access to this medicated feed as it is known to be toxic to these species. Extra care should be taken to avoid contamination of feeds for these animals.

Do not administer tiamulin to birds receiving this medicated feed.

Do not use in feeds containing pellet binding agents.

## Interaction with other medicinal products

Do not administer tiamulin to birds receiving this medicated feed.

## Pregnancy and lactation

Do not feed to replacement, laying or breeding chickens

## Side effects

In the recommended dosage range, Salinomycin has no adverse effects.

## Symptoms and treatment of overdose

Administration of salinomycin sodium in feed at 2 times the recommended dose for 28 consecutive days is safe and nontoxic to quail

## Storage

Store in a cool dry place (<30°C). Strictly avoid light and heat exposure.

## Warning and Precaution

### Special precautions for use in animals

1. Do not feed to replacement, laying or breeding chickens.
2. Operators handling the concentrated product should wear a face mask, gloves and protective clothing.
3. Keep this, and all drugs, out of the reach of children.
4. A weight gain reduction may occur in cockerels fed this medicated feed in the absence of coccidial exposure.
5. Do not use this feed for treatment of outbreaks of coccidiosis.
6. Consult a veterinarian or poultry pathologist if losses exceed 0.5% in a two day period.

-Jauhkan ubat dari kanak-kanak/Keep out of reach of children.

## Shelf life

24 months

## Shelf life after opening

24 hours

## Shelf life after reconstitution or dilution

24 hours

## Withdrawal period

No withdrawal period is required for chickens when this drug product is used according to label directions.

## Packaging

25kg

## Revised date

18/9/2017

## Product Registration Holder:

### PANFAST MARKETING (M) SDN.BHD.

PENANG SCIENCE PARK 1102, MUKIM 13, LORONG PERINDUSTRIAN BUKIT  
MINYAK 18, KAWASAN PERINDUSTRIAN BUKIT MINYAK,  
14100 SIMPANG AMPAT, PENANG, MALAYSIA  
TEL: +604-5016868 FAX: +604-5016866

## Manufactured by:

Inner Mongolia Biok Biology Co., Ltd.  
Tuo Electricity Industrial Park, Tuoketuo County,  
Inner Mongolia, P.R.China  
Tel: 86-471-8661126 Fax : 86-471-8661477

text]