

PELANGI VET PHARMA SDN.BHD^(665097-K)
Lot 8, PERDA Industrial Park, Lorong IKS, Simpang Ampat B,
14100 Simpang Ampat, S.P.S, Pulau Pinang, Malaysia

PRODUCT LITERATURE

1. Product Name:

PVP-FENBENDAZOLE 10% W/V SUSPENSION

DESCRIPTION

2. Each 1 L solution contains:

Active Ingredient:

Fenbendazole BP	100.00g
-----------------	---------

Preservatives:

Sodium Propyl Paraben BP	200.00 mg
--------------------------	-----------

Sodium Methyl Paraben BP	2.00 g
--------------------------	--------

Sodium Benzoate BP	2.00 g
--------------------	--------

3. Product Description:

A light orange coloured suspension

4. Target Species:

Cattle, sheep, horses and other equines

5. Pharmacodynamics:

Fenbendazole is an anthelmintic belonging to the benzimidazolecarbamates group. It acts by interfering with the energy metabolism of the nematode. The anthelmintic efficacy is based on inhibition of the polymerisation of tubulin to microtubuli. The anthelmintic affects both adult and immature stages of gastro-intestinal and respiratory nematodes.

6. Pharmacokinetics:

Fenbendazole is only partly absorbed after oral administration and is then metabolised in the liver. The half life of fenbendazole in serum after oral application of the recommended dose in cattle is 10-18 hours and in sheep 21-33 hours. Fenbendazole and its metabolites are distributed throughout the body and high concentrations can be found in the liver. The elimination of fenbendazole and its metabolites occurs primarily via the faeces (>90%) and to a small extent in the urine and milk. Fenbendazole is metabolised to its sulfoxide then to sulfone and amines.

7. Environmental Properties:

Any unused product or waste material should be disposed of in accordance with national requirements. DANGEROUS to aquatic life. Do not contaminate ponds, waterways or ditches with the product or used container.

8. Indication:

Cattle and sheep

A broad spectrum anthelmintic for the treatment of sheep and cattle infected with mature and developing immature forms of nematodes of the gastro-intestinal and respiratory tracts.

Cattle: For the treatment of cattle infected with:

Ostertagia spp.	Cooperia spp.
Trichostrongylus spp.	Nematodirus spp.
Haemonchus spp.	Oesophagostomum spp.
Bunostomum spp.	Strongyloides spp.
Trichuris spp.	<i>Dictyocaulus viviparus</i>

The product is usually effective against inhibited larvae of *Ostertagia* spp. and against *Moniezia* spp. of tapeworm.

Sheep: For the treatment of sheep infected with benzimidazole susceptible:

Ostertagia spp.	Haemonchus spp.
Trichostrongylus spp.	Nematodirus spp.
Cooperia spp.	Oesophagostomum spp.

Chabertia spp.
Strongyloides spp.

Bunostomum spp.
Dictyocaulus filaria

The product is usually effective against *Moniezia* spp. of tapeworm and may have useful but variable efficacy against *Trichuris* spp.

Horses

For the treatment and control of adult and immature round worms of the gastrointestinal tract in horses and other equines. The product effectively treats and controls the following roundworm infections:

Large strongyles (adults and migrating larval stages of *S. vulgaris*; adults and tissue larval stages of *S. edentatus*). Benzimidazole susceptible adult and immature small strongyles (Cyathostomes). The product is also effective for the treatment and control of encysted mucosal 3rd and 4th stage small strongyle larvae and is also effective against encysted inhibited 3rd stage small strongyle larvae in the mucosa. Adult and immature *Oxyuris* spp., *Strongyloides* spp. and *Parascaris equorum*. Fenbendazole also has an ovicidal effect on nematode eggs.

9. Recommended Dose:

Shake container before use.

For oral administration only.

No dietary control is required before or after treatment.

To ensure administration of a correct dose, body weight should be determined as accurately as possible; accuracy of the dosing device should be checked.

Cattle and Horses: Administer orally 1 ml of the product per 13 kg body weight. (= 7.5 mg fenbendazole/kg body weight)

Practical dosage recommendations:

65 kg	5 ml
135 kg	10 ml
200 kg	15 ml
265 kg	20 ml
335 kg	25 ml
400 kg	30 ml

Above 400 kg, an extra 3.75 ml are required for each additional 50 kg body weight.

Sheep: Administer orally 0.5 ml per 10 kg body weight (= 5 mg fenbendazole/kg body weight)

Practical dosage recommendations:

Up to 10 kg	0.5 ml
11 to 20 kg	1.0 ml
21 to 30 kg	1.5 ml
31 to 40 kg	2.0 ml
41 to 50 kg	2.5 ml
51 to 60 kg	3.0 ml
61 to 70 kg	3.5 ml
71 to 80 kg	4.0 ml

Above 80 kg, an extra 0.5 ml is required for each additional 10 kg body weight.

For administration to horses, thoroughly mix the product with grain or concentrate feed and give the full dosage as one administration. Treatment should be repeated when natural re-infection of animals with parasitic worms occurs.

Horses:

Recommended dosage programs

All horses should be routinely wormed with the single dose regimen every 6-8 weeks.

Five day course

For the treatment and control of migrating and issue larval stages of large strongyles, encysted mucosal 3rd and 4th stage small strongyle larvae and encysted inhibited 3rd stage small strongyle larvae in the mucosa administer 5 ml of the product per 64 kg body weight daily for 5 days (= 7.5 mg fenbendazole/kg body weight daily for 5 days).

Single dose treatment

For the treatment and control of encysted mucosal stages of small strongyles administer 3 ml of the product per 10 kg body weight (= 30 mg fenbendazole / kg body weight). For the treatment and control of migrating and tissue stages of large strongyles administer 6 ml of the product per 10 kg body weight (= 60 mg fenbendazole / kg body weight). For the treatment of diarrhoea cause by Strongyloides westeri in two to three week old sucking foals administer 5 ml of the product per 10kg body weight (= 50mg fenbendazole / kg body weight). Do not mix with other products.

10. Mode of Administration:

Oral

11. Contraindications:

Do not use in horses and other equines intended for human consumption. Fenbendazole as a medicated liquid feed should not be used in the treatment of clinical infestations in cattle and sheep.

12. Warning & Precautions:

When administered by divided dosage in the form of liquid feed, the product may not be effective against *Strongyloides* and *Trichuris* spp. in cattle and *Strongyloides*, *Dictyocaulus* and *Bunostomum* spp. in sheep. Care should be taken to avoid the following practices because they increase the risk of development of resistance and could ultimately result in ineffective therapy:

Too frequent and repeated use of anthelmintics from the same class, over an extended period of time. Under dosing, which may be due to underestimation of body weight, misadministration of the product, or lack of calibration of the dosing device (if any).

Suspected clinical cases of resistance to anthelmintics should be further investigated using appropriate tests (e.g. Faecal Egg Count Reduction Test). Where the results of the test(s) strongly suggest resistance to a particular anthelmintic, an anthelmintic belonging to another pharmacological class and having a different mode of action should be used. Resistance to benzimidazoles (which include fenbendazole) has been reported in *Teladorsagia*, *Haemonchus*, *Cooperia* and *Trichostrongylus* species in small ruminants in a number of countries, including the EU. Therefore the use of this product should be based on local (regional, farm) epidemiological information about susceptibility of nematodes and recommendations on how to limit further selection for resistance to anthelmintics. Resistance to fenbendazole has been reported in cyathostomes in horses. Therefore the use of this product should be based on local (regional, farm) epidemiological information about susceptibility of nematodes and recommendations on how to limit further selection for resistance to anthelmintics.

Special precautions for use in animals

When incorporating this product into liquid feed, after thoroughly shaking the suspension, measure the required volume of the suspension and add it to approximately 10% of the liquid feed. Thoroughly mix this material and then add the remaining liquid feed and once again mix to produce a homogenous dispersion. Mix the medicated feed thoroughly prior to administration for example by rolling the drum or barrel. Assess body weight as accurately as possible before calculating the dosage. Intensive use or misuse of anthelmintic can give rise to resistance. To reduce this risk, dosing programmes should be discussed with your veterinary surgeon.

Special precautions to be taken by the person administering the medicinal product to the animals

Direct contact with the skin should be kept to a minimum. Wear suitable protective clothing including impermeable rubber gloves. Wash hands after use.

13. Drug Interactions:

None known

14. Pregnancy & Lactation:

The product can be administered to pregnant animals. Pregnant mares and young foals may also be safely treated with fenbendazole at the therapeutic dosage levels.

15. Side effects:

None

16. Symptoms & Treatment of Overdose:

Benzimidazoles have a high margin of safety. No specific overdose symptoms are known. No specific actions required.

17. Withdrawal Period:

Cattle – Meat: 12 Days

Milk: 5 Days

Sheep - Meat: 15 Days

Milk: 7 Days

Not to be used in horses intended for human consumption. Treated horses may never be slaughtered for human consumption. The horse must have been declared as not intended for human consumption under national horse passport legislation.

18. Storage:

Store below 30° C in dry place, protected from light.

19. Dosage Form:

Solution in the form of Suspension

20. Packing:

Packed in 1L in HDPE plastic container with caps.

21. Disposal of Container:

To be disposed as per local authority requirement.

22. Expiry date:

2 years from date of manufacturing.

Manufactured & Distributed by :

PelangiVet PharmaSdn Bhd. (665097-K)
Lot 8, PERDA Industrial Park,
Lorong IKS, SimpangAmpat B
14100 SimpangAmpat,
S.P.S Pulau Pinang, Malaysia

Last update on : 12/12/2016