

**PELANGI VET PHARMA SDN.BHD** (665097-K)  
LOT 8, PERDA Industrial Park, Lorong IKS, SimpangAmpat B,  
14100 SimpangAmpat, S.P.S, Pulau Pinang, Malaysia

## **PRODUCT LITERATURE**

### **1. Product Name:**

**PVP-TRICLABENDAZOLE 12%W/V SUSPENSION**

### **2. Each 1L suspension contains:**

#### **Active Ingredient:**

Triclabendazole 120.00g

#### **Preservatives:**

Sodium Propyl Paraben BP 200.00mg

Sodium Methyl Paraben BP 2.00g

Sodium Benzoate BP 2.00g

### **3. Product Description:**

Orange coloured suspension

### **4. Target Species:**

Cattle & Sheep

## **5. Pharmacodynamics:**

Triclabendazole differs from other benzimidazoles in that it is a narrow spectrum anthelmintic. The drug accumulates significantly in both immature and adult stages of *Fasciola hepatica* and stimulates the major routes of the parasite's energy generating system, as demonstrated by glucose derived acetate and propionate formation. However, under these conditions the parasite's motility decreased, indicating that the drug is not associated with inhibition of the energy generating pathways. Triclabendazole inhibits colchicine binding to microtubular proteins suggesting interference of the drug with microtubular structure and function. The drug strongly inhibits the release of proteolytic enzymes in immature and adult parasites, a process dependant on microtubular functions. The precise molecular mode of action of this fasciolicidal drug remains to be elucidated.

## **6. Pharmacokinetics:**

50-75% of the orally administered dose of triclabendazole is absorbed from the gastrointestinal tract. Very rapidly, absorbed triclabendazole is almost completely oxidised to its sulfoxide and sulfone. In cattle triclabendazole sulfoxide reaches peak concentrations approximately 27 hours after administration of the product and the sulfone reaches peak concentrations 64 to 72 hours after administration. In sheep triclabendazole sulfoxide reaches peak concentrations approximately 20 hours after administration of the product and the sulfone reaches peak concentrations 30 to 32 hours after administration. Both metabolites bind strongly to plasma proteins, particularly albumin. Metabolites are excreted via the bile mainly as conjugates. More than 90%- 95% of the total dose of triclabendazole is excreted in the faeces, about 2% in the urine and less than 1% in the milk. The elimination is virtually complete by 10 days after administration.

## **7. Environmental Properties**

*The product may have toxic effects on fish and aquatic invertebrates. Any unused product or waste material must not enter surface water and should be disposed of in accordance with national requirements.*

## **8. Indication:**

For the treatment of adult, immature and early immature stages of liver fluke (*Fasciola hepatica*) susceptible to triclabendazole.

## 9. Recommended Dose:

For single oral administration only using properly calibrated dosing equipment.

The product is suitable for most types of automatic drenching guns.

Shake the container before use.

Use unaltered from original container.

Clean drenching equipment before and after use.

PVP- Triclabendazole 12% w/v suspension is given as an oral drench and is suitable for most types of automatic drenching guns.

Recommended dose rate: 12 mg triclabendazole per kg bodyweight in cattle and 10mg triclabendazole per kg bodyweight in sheep.

**Practical Dosage Guide: Cattle: 5ml per 50kg bodyweight**

| Animal Weight            | Dose of product |
|--------------------------|-----------------|
| 50kg                     | 5ml             |
| 100kg                    | 10ml            |
| 150kg                    | 15ml            |
| 200kg                    | 20ml            |
| 250kg                    | 25ml            |
| 300kg                    | 30ml            |
| 350kg                    | 35ml            |
| 400kg                    | 40ml            |
| For each additional 50kg | 5ml             |

**Sheep: 0.8ml per 10kg bodyweight**

| Animal Weight            | Dose of product |
|--------------------------|-----------------|
| 10kg                     | 0.8ml           |
| 20kg                     | 1.6ml           |
| 30kg                     | 2.4ml           |
| 40kg                     | 3.2ml           |
| 50kg                     | 4ml             |
| 60kg                     | 4.8ml           |
| For each additional 10kg | 0.8ml           |

To ensure administration of a correct dose, bodyweight should be determined as accurately as possible; accuracy of the dosing device should be checked.

If animals are to be treated collectively rather than individually, they should be grouped according to their bodyweight and dosed accordingly, in order to avoid under- or overdosing.

The timing for re-treatment should be based on epidemiological risk patterns and should be customised for each individual farm.

To avoid the potential for the accumulation of residues following repeat administration of the product; animals should not be treated with a frequency of less than 10 weeks.

#### **10. Mode of Administration:**

Oral

#### **11. Contraindications:**

Do not use in animals known to be hypersensitive to the active substance.

#### **12. Warning & Precautions:**

Care should be taken to avoid the following practices, because they increase the risk of development of resistance and could ultimately result in ineffective therapy:

- Too frequent and repeated use of anthelmintics from the same class, over an extended period of time.
- Under dosing, which may be due to under estimation of bodyweight, misadministration of the product or lack of calibration of the dosing device (if any).

Suspected clinical cases of resistance to anthelmintics should be further investigated using appropriate tests (e.g. Faecal Egg Count Reduction Test). Where the results of the test(s) strongly suggest resistance to a particular anthelmintic, an anthelmintic belonging to another pharmacological class and having a different mode of action should be used.

Resistance to triclabendazole has been reported in *Fasciola hepatica* in cattle and sheep. Therefore, the use of this product should be based on local (regional/farm) epidemiological information about

susceptibility of the Fasciola Hepatica and recommendations on how to limit further selection for resistance to anthelmintics.

**Special precautions for use in animals**

Care should be taken when dosing animals to avoid causing injury to the mouth and pharynx.

**Special Precautions to be taken by the person administering the product to animals**

When using, do not eat, drink or smoke.

Wash splashes from eyes and skin immediately.

Take off immediately any contaminated clothing.

Wash hands and exposed skin before meals and after use.

**Other precautions**

The use of this product may have harmful effects on fish and aquatic invertebrates. Cattle and Sheep must not have any access to the surface water such as streams, ponds or ditches within days after treatment. When spreading manure from treated animals on arable lands, a safety distance of 10m to adjacent surface waters must be kept.

**13. Drug Interactions:**

None known

**14. Pregnancy & Lactation:**

The product is safe for use during pregnancy and lactation. However, the product is not permitted for use during lactation in animals producing milk for human consumption.

**15. Side effects:**

Occasionally, inflammation of the unpigmented skin, including the udder and the teats may occur after treatment in cattle exposed to intense sunshine.

**16. Symptoms & Treatment of Overdose:**

A single oral dose of 150-200mg Triclabendazole/kg of live bodyweight may lead to side effects such as unsteady gait, dullness and reduced appetite. These side effects are slight and last to 1 to 5 days. An antidote is not known.

**17. Withdrawal Period:**

Cattle ( meat and offal): 56 days

Milk:

Cattle (milk): Milk for human consumption may only be taken from 48 hours after calving. Not intended for use within 45 days of calving. Should a cow calve earlier than 45 days after the last treatment, milk for human consumption may only be taken from 45 days + 48 hours (47 days) after the last treatment.

Sheep (meat & offal): 56 days

Sheep (milk): *Not authorised for use in ewes producing milk for human consumption including during the dry period. Do not use within 1 year prior to the first lambing in ewes intended to produce milk for human consumption*

**18. Storage:**

Store below 30° C in dry place, protected from light.

**19. Dosage Form:**

Solution in the form of suspension.

**20. Packaging :**

Packed in 1L HDPE plastic container with caps.

**21. Disposal of Container :**

To be disposed as per local authority requirement.

**22. Expiry date :**

2 years from date of manufacturing

**Manufactured & Distributed by:**

**Pelangi Vet Pharma Sdn Bhd.** (665097-K)

Lot 8, PERDA Industrial Park,

Lorong IKS, SimpangAmpat B

14100 SimpangAmpat,

S.P.S Pulau Pinang, Malaysia

Last update on: 12 December 2016