

PRODUCT LITERATURE

1. Product Name: PVP- AMPROLIUM 20% W/W POWDER

2. Each 1 g powder contains:

Active Ingredient:

Amprolium Hydrochloride	226.2 mg
(Equivalent to Amprolium	200.0 mg)

3. Product Description:

White to off white powder.

4. Target Species:

Poultry (chickens, turkeys, ducks, guinea fowls, pheasants, partridges).

5. Pharmacodynamics

Amprolium is an anticoccidial agent belonging to the family of thiamine analogues. Amprolium acts by interfering as a competitive antagonist in thiamine transport mechanisms. It interferes in the metabolism of carbohydrates necessary for the multiplication and survival of coccidian.

6. Pharmacokinetics:

Amprolium is poorly absorbed after oral administration. The maximum plasma concentration is reached 4 hours later. The amprolium is eliminated mainly by faeces.

7. Indication:

In poultry (chickens, turkeys, ducks, guinea fowls, pheasants, partridges): - Treatment of coccidiosis due to coccidia sensitive to amprolium.

8. Recommended Dose:

Oral use.

20 mg amprolium per kg bodyweight per day, orally in the food, for 7 days.

For daily food intake of 100 g of food per kg of body weight, this dosage corresponds to 200 ppm of amprolium in the food, ie an incorporation rate of 1kg of PVP- Amprolium 20 % w/w Powder per tonne of food.

For a guinea fowl of 980 g of live weight consuming 80 g of food per day, this dosage corresponds to 245 ppm of amprolium in the food, ie an incorporation rate of 1.2 kg of PVP- Amprolium 20 % w/w Powder per tonne of food.

In order to respect the dosage and to take account of the effective intake of food, the incorporation rate may be increased or decreased, which leads to ppm levels in the diet.

9. Mode of Administration

For oral administration, mixed in feed.

10. Contraindications:

Not known.

11. Warning & Precautions:

Special warnings for each target species

Do not administer to laying birds whose eggs are intended for human consumption.

Special precautions for use in animals

This medicinal premix is intended for the preparation of solid medicated feed and cannot be used as its present state.

The incorporation rate of PVP- Amprolium 20 % w/w Powder cannot be less than 1 kg / tonne.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Avoid inhalation of dust.

Do not eat or drink while using.

Clean the skin thoroughly after work.

In the event of accidental spillage into the eyes, rinse thoroughly with water. If irritation persists, seek medical advice and show the label.

In case of accidental ingestion, consult a doctor.

Other precautions

None.

12. Drug Interactions:

Not known.

13. Pregnancy & Lactation:

Not applicable.

14. Side effects:

Not known.

15. Symptoms & Treatment of Overdose:

Prolonged use at high doses can cause thiamine deficiency. This deficit can be offset by a thiamine supply.

16. Withdrawal Period:

Meat and offal: zero days.

Egg: In the absence of determination of a waiting time for eggs, not to be used in egg-producing egg-producing species, 4 weeks before and during the start of egg production.

17. Storage Conditions:

Store below 30° C in dry place, protected from light.

18. Dosage Form:

Powder in the form of powder to be mixed in feed.

19. Packaging:

- 1) Pack of 1 kg in HDPE Plastic Container
- 2) Pack of 500 g in HDPE Plastic Container
- 3) Pack of 1 kg pack in Aluminium Sheet Lacquered
- 4) Pack of 500 g Pack in Aluminium Sheet Lacquered

20. Disposal of Container:

To be disposed as per local authority requirement.

21. Expiry Date:

2 years from date of manufacturing

22. Shelf life after reconstitution or dilution:

To be used within 24 hours after reconstitution.

Manufactured & Distributed by :

Pelangi Vet Pharma Sdn Bhd. (665097K)

Lot 8, PERDA Industrial Park,
Lorong IKS, Simpang Ampat B
14100 Simpang Ampat,
S.P.S Pulau Pinang, Malaysia

Date of revision: 06/04/2017