

**PELANGI VET PHARMA SDN.BHD**

Lot 8, PERDA Industrial Park, Lorong IKS, Simpang Ampat B,  
14100 Simpang Ampat, S.P.S, Pulau Pinang, Malaysia

**PRODUCT LITERATURE****1. Product Name:**

**PVP- CEFADROXIL 15% W/W SOLUBLE POWDER**

**2. Each 1 g powder contains:****Active Ingredient:**

Cefadroxil Monohydrate BP 165.7 mg  
(Equivalent to Cefadroxil 150 mg)

**3. Product Description:**

White to off white powder

**4. Target Species:**

Dogs and cats

**5. Pharmacodynamics:**

Cefadroxil, like other beta-lactam antibiotics, is a bactericidal agent that causes death of bacterial cells through a diversity of biological and biochemical effects on the cell wall. The spectrum of antibacterial activity includes many gram-negative organisms since cefadroxil, like other cephalosporins, has the ability to penetrate the outer envelope of gram-negative bacilli, thereby gaining access to cell wall target sites. Cefadroxil is generally not broken down by penicillinases such as those produced by penicillin-resistant staphylococci, although cephalosporinases have been identified that can inactivate the molecule.

**6. Pharmacokinetics:**

Cefadroxil is stable in gastric acid and only moderately bound to serum proteins (approximately 20%). Cefadroxil is well absorbed from the gastrointestinal tract even when administered with food. The drug is excreted largely unchanged by the kidney. The concurrent administration of probenecid retards the elimination rate.

In dogs, oral administration of cefadroxil at a dosage of 22 mg/kg results in mean peak serum concentration averaging 18.6 mcg/mL within 1 to 2 hours after treatment. The serum half-life ( $T_{1/2}$ ) following oral administration is approximately 2 hours. Over 50% of an orally administered dose is excreted unchanged in the urine of dogs within 24 hours.

In cats, oral administration of cefadroxil at a dosage of 22 mg/kg results in mean peak serum concentrations of 17.4 mcg/mL within 1 to 2 hours after treatment. The serum half-life ( $T_{1/2}$ ) following oral administration to cats is 2½ to 3.

## **7. Indication:**

**Dogs:** Genitourinary tract infections (cystitis) caused by susceptible strains of *Escherichia coli*, *Proteus mirabilis* and *Staphylococcus aureus*.

Skin and soft tissue infections including cellulitis, pyoderma, dermatitis, wound infections and abscesses caused by susceptible strains of *Staphylococcus aureus*.

**Cats:** Skin and soft tissue infections including abscesses, wound infections, cellulitis and dermatitis caused by susceptible strains of *Pasteurella multocida*, *Staphylococcus aureus*, *Staphylococcus epidermidis* and *Streptococcus* spp.

## **8. Recommended Dose:**

**Dogs:** PVP-CEFADROXIL 15% W/W SOLUBLE POWDER should be administered orally at a dosage of 22 mg/kg of body weight twice daily. Dogs with skin or soft tissue infections should be treated for a minimum of three days. Genitourinary tract infections should be treated for a minimum of seven days with cefadroxil. Maximum duration of therapy should not exceed 30 days.

**Cats:** PVP-CEFADROXIL 15% W/W SOLUBLE POWDER should be administered orally at a dosage of 22 mg/kg of body weight once daily. Maximum duration of therapy should not exceed 21 days.

In both species, drug treatment should continue for at least 48 hours after the animal is afebrile or asymptomatic. If no response is observed after three days of treatment, therapy should be discontinued and the case should be re-evaluated.

Tap bottle or aluminium pack lightly to loosen powder. For 1g of PVP-CEFADROXIL 15% W/W SOLUBLE POWDER, add 3ml of water in two portions. Mix well after each addition. Discard unused portion within 24 hours after reconstitution.

**9. Mode of Administration:**

Oral

**10. Contraindications:**

This product should not be administered to dogs or cats with a known allergy to cephalosporins. In penicillin-allergic animals, PVP-CEFADROXIL 15% W/W SOLUBLE POWDER should be used with caution.

**11. Warning & Precautions:**

For use in dogs and cats only. Do not use in animals intended for human consumption. Safety for use in pregnant female dogs and cats or in breeding males has not been determined.

**12. Drug Interactions:**

The concurrent administration of probenecid retards the elimination rate.

**13. Pregnancy & Lactation:**

Safety for use in pregnant female dogs and cats or in breeding males has not been determined.

**14. Side effects:**

Occasional nausea and vomiting have been reported following cefadroxil therapy. Administration with food appears to decrease nausea. Diarrhea and lethargy have been occasionally reported.

**15. Symptoms & Treatment of Overdose:**

Acute oral cephalosporin overdoses are unlikely to cause significant problems other than GI distress, but other effects are possible.

High doses or very prolonged use of cephalosporins have been associated with neurotoxicity, neutropenia, agranulocytosis, thrombocytopenia, hepatitis, positive Comb's test, interstitial nephritis, and tubular necrosis. Except for tubular necrosis and neurotoxicity, these effects have an immunologic component.

**16. Withdrawal Period:**

Not Applicable

**17. Storage Conditions:**

Store below 30°C in dry place, protected from light.

**18. Dosage Form:**

Powder in the form of powder to be dissolved in the water.

**19. Packaging:**

- i) Pack of 1 kg in HDPE Plastic Container
- ii) Pack of 500 g in HDPE Plastic Container
- iii) Pack of 1 kg pack in Aluminium Sheet Lacquered
- iv) Pack of 500 g pack in Aluminium Sheet Lacquered

**20. Disposal of Container:**

To be disposed as per local authority requirement.

**21. Expiry Date:**

2 years from date of manufacturing.

**22. Shelf life after reconstitution or dilution:**

To be used within 24 hours after reconstitution.



**Manufactured & Distributed by :**

**Pelangi Vet Pharma Sdn Bhd.** (665097K)

Lot 8, PERDA Industrial Park,  
Lorong IKS, Simpang Ampat B  
14100 Simpang Ampat,  
S.P.S Pulau Pinang, Malaysia

Date of revision: 05/06/2017