

OMS TYLOFAC 250MG/G PREMIX POWDER

Composition:

Each gm contains:

Tylosin (as tylosin phosphate).....250 mg

Product Description:

OMS Tylofac 250mg/g Premix Powder is a yellow or brownish yellow powder

Pharmacodynamics:

Tylosin is a macrolide antibiotic produced by a strain of *Streptomyces fradiae*. It exerts its antimicrobial effect by inhibiting protein synthesis of susceptible micro-organisms.

The tylosin spectrum of activity includes Gram-positive bacteria, some Gram-negative strains such as *Pasteurella*, and *Mycoplasma* spps at concentrations of 16µg/ml or less.

Pharmacokinetics:

Absorption: Tylosin reaches maximal blood levels between 1 and 3 hours after an oral dose. Minimal or no blood levels remain 24 hours after an oral dose.

Distribution: After oral doses were given to pigs, tylosin was found in all tissues, between 30 minutes and two hours after administration, except for the brain and spinal cord.

Biotransformation and Elimination: It has been shown that most of the material which is excreted is to be found in the faeces and consists of tylosin (factor A), relomycin (factor D) and dihydrodesmycosin.

Indications:

Pigs:

For the prevention and control of enzootic pneumonia.

For the treatment and control of *Lawsonia intracellularis*, the organism associated with Porcine Intestinal Adenomatosis (Ileitis) and Porcine Haemorrhagic Enteropathy.

Broilers and pullets:

Treatment and prevention of respiratory infections caused by *Mycoplasma gallisepticum* and *Mycoplasma synoviae*, when the disease has been diagnosed in the flock.

Treatment and prevention of necrotic enteritis caused by *Clostridium perfringens*, when the disease has been diagnosed in the flock.

Mode of Administration:

Orally. Mixed into feed.

Recommended Dosage:

For oral administration.

Pigs:

Prevention and control of enzootic pneumonia:

Treatment and control of Lawsonia intracellularis:

3-6 mg tylosin activity/kg bodyweight, which may normally be achieved by adding the product at the rate of 400 g per tonne, giving 100 g tylosin base per tonne. Feed as the only ration for 21 days.

Broilers and pullets:

For the treatment and prevention of respiratory infections:

127 mg tylosin per kg bodyweight (corresponding to 508 mg OMS Tylofac 250mg/g Premix Powder per kg bodyweight) for the first 5 days of life. It is strongly recommended to repeat the treatment of the birds at the age of 3-4 weeks.

For the treatment and prevention of necrotic enteritis:

10 – 20 mg tylosin per kg bodyweight (corresponding to 40 – 80 mg OMS Tylofac 250mg/g Premix Powder per kg bodyweight) for 7 days.

Contraindications:

Do not use in known cases of hypersensitivity.

Do not use in animals vaccinated with tylosin-sensitive vaccines either at the same time or within one week previously.

Warnings and Precautions:

Special warnings for each target species

None.

Special precautions for use

Official, national and regional antimicrobial policies should be taken into account when the product is used.

- i. Special precautions for use in animals
For incorporation into medicated feedingstuffs only.

To ensure thorough dispersion of the product it should first be mixed with a small quantity of feed ingredients before incorporation into the final mix.

- i. Special precautions to be taken by the person administering the veterinary medicinal product to animals

Tylosin may induce irritation. Macrolides, such as tylosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tylosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided.

To avoid exposure during preparation of the medicated feed, wear overalls, safety glasses, impervious gloves and wear either a disposable half mask respirator. Wash hands after use.

In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water.

Do not handle the product if you are allergic to ingredients in the product.

If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

Only to be incorporated by a manufacturer who is approved to mix at a rate of below 2 kg per tonne final feed.

Interactions with Other Medicaments:

Lincosamides and aminoglycoside antibiotics antagonise the activity of tylosin.

Pregnancy and Lactation:

No studies have been conducted in the target species population. Use only according to the benefit risk assessment by the responsible veterinarian.

Adverse Effects/Undesirable Effects:

None known.

Overdose and Treatment:

No adverse effects when fed to pigs at 600 ppm in the feed (six times the recommended maximum level) for 28 days.

Withdrawal Period(s):

Meat:

Pigs: Zero days.

Broilers and pullets: 1 day

Do not use in laying hens producing eggs for human consumption.

Storage conditions:

Store in a cool dry place, protected from light. Do not store above 30°C.

Keep out of reach of children/Jauhkan dari kanak-kanak.

Shelf life:

Shelf life as packaged for sale: 12 months

Shelf life after reconstitution into feed: 24 hours

Pack size(s):

Paper bag of 25kg.

Product Registration Holder/ Manufacturer

Osmosis Nutrition Sdn. Bhd. (594876A)

Lot 16014(P.T. 24341), Jalan Nilam 3,

Bandar Nilai Utama,

71800 Nilai, Negeri Sembilan, Malaysia.

Tel: +606-799 9090 Fax: +606-799 9155

Date of revision of Package Insert:

12/11/2017