

Veterinary Package Insert

TP-FLOR 50MG/G Powder (5% Florfenicol)

PRODUCT DESCRIPTION

White or almost white colour powder, which contains 50mg of florfenicol per gram in 1kg sachet.

PHARMACODYNAMICS & PHARMACOKINETICS

Florfenicol is a broad-spectrum synthetic antibiotic in the phenicol group that is active against most Gram-positive and Gram-negative bacteria isolated from domestic animals. Florfenicol acts by inhibition of protein synthesis at the ribosomal level and is bacteriostatic. However, bactericidal activity has been demonstrated in-vitro against *Pasteurella multocida* when florfenicol is present at concentrations above the MIC for 4 to 12 hours.

In-vitro testing has shown that florfenicol is active against the bacterial pathogens most commonly isolated in respiratory diseases in pigs, including *Pasteurella multocida*.

The only mechanisms of chloramphenicol resistance that are known to have significant clinical relevance are CAT (Chloramphenicol Acetyl Transferase)-mediated inactivation and efflux-pump resistance. Of these, only some of the efflux mediated resistance would also confer resistance to florfenicol and thus have the potential to be affected by florfenicol use in animals.

Florfenicol is well absorbed when administered orally and following distribution it is rapidly eliminated in the urine and faeces in a ratio of 3:1. A fraction is excreted unchanged and the rest is metabolised into 5 major metabolites.

INDICATION

For the treatment and prevention of swine respiratory disease caused by *Pasteurella multocida* susceptible to florfenicol in infected herds. The presence of the disease should be established in the herd before initiating preventive treatment.

Targeted species: Pigs (Fattening pigs)

RECOMMENDED DOSAGE AND ADMINISTRATION

To be administered orally, in medicated feeding stuff.

Dosage:

10 mg of florfenicol per kg body weight (bw) (equivalent to 200 mg TP-Flor 50mg/g Powder) per day administered for 5 consecutive days.

Administration:

For a daily feed intake of 50g/kg body weight, this dosage corresponds to a rate of incorporation of 4 kg of premix per ton of feed, i.e. 200 ppm of florfenicol.

The rate of incorporation of the medicated premix in the feed may be increased in order to achieve the required dosage on a mg/kg body weight basis and to take into account the actual feed intake. Thus, the inclusion level may need adjusting as follows to give the correct dose.

200 mg of PRODUCT per kg body weight and day	x	Average pig body weight (kg)	= mg PRODUCT per kg of feed
--	---	---------------------------------	--------------------------------

Average daily feed intake (kg/animal)

The maximum rate of incorporation is 10 kg/ton (500 ppm of florfenicol), higher rates of inclusion may lead to poor palatability and decreased food consumption.

Under no circumstances should the incorporation rate of the premix be below 4kg/ton of feed.

In all cases the recommended dose of 10mg of florfenicol per kg of body weight per day, for 5 consecutive days has to be respected.

CONTRAINDICATIONS

Do not administer to boars intended for breeding.

Do not use in case of hypersensitivity to the active substance or any of the excipients.

WARNINGS & PRECAUTIONS.

Special warnings for each target species:

Animals showing a decreased appetite and/or a poor general condition should be treated by the parenteral route.

Special precautions for use in animals:

The product should be used in conjunction with susceptibility testing and take into account official and local policy relating to the use of antimicrobials.

This premix contains calcium carbonate. Care should therefore be taken to consider the calcium content of the final medicated feed.

Treatment should not exceed 5 days.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Skin sensitisation may occur.

Avoid skin contact.

Do not handle this product in case of known sensitization to active substance or any of the excipients.

Handle this product with care to avoid exposure during incorporation of premix into feed and administration of feed to animals, taking all recommended precautions.

Wear either a disposable half-mask respirator or a non-disposable respirator with a filter, chemically resistant gloves, protective coveralls and goggles while incorporating the premix into feed.

Wear gloves and do not smoke, eat, or drink when handling the product or medicated feed.

Wash hands thoroughly with soap and water after use of the product or medicated feed

Rinse thoroughly with water in case of exposure.

If you develop symptoms following exposure such as skin rash, you should seek medical advice and take the package leaflet or the label with you.

Other precautions:

Manure from treated swine must be stored for a minimum of one month before being spread and incorporated in fields.

INTERACTIONS WITH OTHER MEDICATIONS

No data available.

PREGNANCY AND LACTATION

The safety of the product during pregnancy and lactation has not been studied in sows. Use of the product during pregnancy and lactation is therefore not recommended.

ADVERSE EFFECTS

Commonly observed adverse effects are diarrhoea perianal inflammation and rectal eversion. Increased serum calcium may also be observed. These effects are transient, resolving on cessation of treatment.

OVERDOSE & TREATMENT

In the event of overdose, a reduction in food and water consumption, together with a decrease in bodyweight may be observed. There may be an increase in refused feed and an increase in serum calcium.

WITHDRAWAL PERIOD

Meat and offal: 14 days

MAXIMUM RESIDUAL LIMIT (MRL)

Substance	Food	Maximum Residue Limits (MRLs) in µg/kg
Florfenicol	Pig, muscle	300
	Pig, skin & fat	500
	Pig, liver	2000
	Pig, kidney	500

STORAGE CONDITION

Store below 30°C. Protect from direct sunlight.

SHELF LIFE

2 years.

After 1st opening, reconstitution or dilution, use within 24 hours.

PACKING

500g

1kg

THYE PHARMA SDN. BHD.

(manufacturer & product registration holder)

No.2 Lorong Industri Ringan Permatang
Tinggi 8,

Kawasan Industri Ringan Permatang Tinggi,
14000 Bukit Mertajam,
Pulau Pinang.

Revised Aug 2017