

EASIFLAM MOUTHWASH 0.15 % w/v

VIEASxx-01 (MY)

DESCRIPTION

Clear green solution with an odour of peppermint.

COMPOSITION

Each 15 ml dose contains:
Benzydamine Hydrochloride 22.5mg

ACTIONS AND PHARMACOLOGY

The indazole analogue benzydamine has physicochemical properties and pharmacological activities which differ from those of the aspirin-like NSAIDs. Unlike aspirin-like NSAIDs which are acids or metabolised to acids, benzydamine is a weak base. In further contrast, benzydamine is a weak inhibitor of the prostaglandin synthesis. Only at concentration of 1mM and above benzydamine effectively inhibits cyclooxygenase and lipoxygenase enzyme activity. It mostly exerts its effects through inhibition of the synthesis of proinflammatory cytokines including tumour necrosis factor-alpha (TNF- α) and Interleukin-1 β (IL-1 β) without significantly affecting other pro-inflammatory (IL-6 and 8) or anti-inflammatory cytokines (IL-10, IL-1 receptor antagonist). Further mechanisms of action are hypothesised including the inhibition of the oxidative burst of neutrophils as well as membrane stabilisation as demonstrated by the inhibition of granule release from neutrophils and the stabilization of lysosomes. The local anaesthetic activity of the compound has been related to an interaction with cationic channels.

PHARMACOKINETICS

Oral doses of benzydamine are well absorbed and plasma drug concentrations reach a peak fairly rapidly and then decline with a half-life of about 13 hours. Less than 20% of the drug is bound to plasma proteins.

Although local drug concentrations are relatively large, the systemic absorption of mouthwash-gargle doses of benzydamine is relatively low compared to oral doses. This low absorption should greatly diminish the potential for any systemic drug side-effects when benzydamine is administered by this route. Benzydamine is metabolized primarily by oxidation, conjugation and dealkylation.

INDICATION

For the temporary relief of painful conditions of the oral cavity including: tonsillitis, sore throat, radiation mucositis, aphthous ulcers, post-oral surgical and periodontal procedures.

CONTRAINDICATIONS

Easiflam Mouthwash is contra-indicated in patients with known hypersensitivity to the active substance benzydamine hydrochloride or to any of the excipients.

PREGNANCY AND LACTATION

Pregnancy

Easiflam Mouthwash should not be used in pregnancy unless considered essential by the physician.

Breast-feeding

Easiflam Mouthwash should not be used in lactation unless considered essential by the physician.

PRECAUTIONS

- Benzydamine use is not advisable in patients with hypersensitivity to acetylsalicylic acid or other NSAIDs.
- Easiflam Mouthwash should generally be used undiluted, but if 'stinging' occurs the rinse may be diluted with water.
- Avoid contact with eyes.

INTERACTIONS WITH OTHER MEDICAMENTS

None known.

MAIN SIDE/ ADVERSE EFFECTS

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

The following rate values have been used: Very common, Common, Uncommon, Rare, Very rare and not known.

The most common side effects are numbness and a stinging feeling in the mouth.

Respiratory, thoracic and mediastinal disorders

Very rare: Laryngospasm or bronchospasm

Gastrointestinal disorders

Uncommon: Oral numbness (hypoesthesia) and a stinging feeling in the mouth (oral pain)

Skin and subcutaneous tissue disorders

Very rare: Hypersensitivity reactions which may be associated with pruritus, urticaria, photosensitivity reaction and rash

Frequency not known: Angioedema

Immune system disorders

Frequency not known: Anaphylactic reaction which can be potentially life-threatening.

OVERDOSE AND TREATMENT

Intoxication is only to be expected if large quantities of Easiflam Mouthwash are swallowed.

Symptoms associated with ingested overdose of benzydamine are mainly gastrointestinal symptoms and symptoms of the central nervous system. Most frequent gastrointestinal symptoms are nausea, vomiting, abdominal pain, and esophageal irritation. Symptoms of the central nervous system include dizziness, hallucinations, agitation, anxiety, and irritability.

In acute overdose only symptomatic treatment is possible. Patients should be kept under close observation and supportive treatment should be given. Adequate hydration must be maintained.

DOSAGE AND ADMINISTRATION

Easiflam Mouthwash should generally be used undiluted, but if stinging occurs it may be diluted with water. The solution should be expelled from the mouth after use. Uninterrupted treatment should not exceed seven days.

Dosage in Adults

When used as a gargle, the usual dose of Easiflam Mouthwash is 15ml (approximately one tablespoon) which should be gargled for at least 30 seconds at 1 to 3 hourly intervals, as needed.

When used as a rinse for oral lesions, the usual dose is again 15ml (approximately one tablespoon) which should be held in the mouth and swirled around for at least 30 seconds, with repeat use every 1 to 3 hours throughout the day.

Dosage in Children

5-15ml as a gargle if able to do so, or as an oral rinse.

With Impaired Renal Function

Since absorbed benzydamine and its metabolites are excreted in the urine, the possibility of systemic effects should be considered in patients with severe renal impairment.

With Impaired Liver Function

Since absorbed benzydamine is highly metabolised in the liver the possibility of systemic effects should be considered in patients with severe hepatic impairment.

Storage: Store below 30°C. Protect from light.

Presentation/ Packing: 100ml per bottle

Product Owner/ Product Registration Holder
(Malaysia) / Manufactured By:

HOVID Berhad
121, Jalan Tunku Abdul Rahman
(Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia

Information date: August 2023