

CONTROLLED MEDICINE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY
LIFECOME BACITRACIN METHYLENE DISALICYLATE
50% W/W SOLUBLE POWDER

ACTIVE CONSTITUENT: Bacitracin (As bacitracin methylene disalicylate) 500mg/g.

PRODUCT DESCRIPTION: Light brown to greyish-white powder.

PHARMACODYNAMICS/PHARMACOKINETICS:

Bacitracin is an antibiotic from the group of peptide antibacterial compounds. Bacitracin is produced by *Bacillus licheniformis* and *Bacillus subtilis*, and is a mixture of several closely related polypeptides, mainly consisting of bacitracin A, B1, B2, C and a small portion of bacitracin F. Bacitracin is mainly active against gram-positive microorganisms. It inhibits cell wall biosynthesis by inhibiting lipid pyrophosphatase which is involved in transmembrane transport of peptidoglycan precursors. After oral application to rats, chickens and pigs, bacitracin is hardly absorbed from the gastrointestinal tract, and the distribution to organs and tissues is negligible. In rats, chickens and pigs, approximately 95% of an oral dose is excreted via faeces, and only 3% or less via urine. Bacitracin is metabolized to amino acids and smaller peptides via the main metabolite desamidobacitracin, which is microbiologically inactive.

INDICATIONS:

Broiler and replacement chickens: An aid in the prevention and control of necrotic enteritis caused by *Clostridium perfringens* susceptible to bacitracin methylene disalicylate.

Swine: For use in the treatment of swine dysentery associated with *Brachyspira hyodysenteriae*.

RECOMMENDED DOSAGE:

Target Animal	Condition	Prevention (mg bacitracin/L)	Prevention (mg product/L)	Control (mg bacitracin/L)	Control (mg product/L)
Broiler and replacement chickens	Necrotic enteritis caused by <i>Clostridium perfringens</i> susceptible to bacitracin methylene disalicylate. Administer continuously 5 to 7 days or as long as clinical signs persist, then reduce to prevention levels.	26	52	53-106	106-212
Swine	Swine dysentery (bloody scours) associated with <i>Brachyspira hyodysenteriae</i> . Treatment not to exceed 14 days. If symptoms persist after 4 to 5 days consult a veterinarian. Not to be given to swine that weight more than 113.5 kg.	-	-	264	528

MODE OF ADMINISTRATION: Oral.

CONTRAINDICATIONS: Do not use in known cases of hypersensitivity to active ingredient.

WARNINGS AND PRECAUTIONS: Will irritate the eyes, nose and throat. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Do not inhale dust. When opening the container and mixing into water, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, goggles and disposable dust mask. If product in eyes, wash it out immediately with water. After each day's use, wash gloves, goggles and contaminated clothing.

INTERACTIONS WITH OTHER MEDICAMENTS: In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other products.

PREGNANCY AND LACTATION: In the absence of toxicology study, this veterinary medicinal product must not be used during pregnancy and lactation.

UNDESIRABLE EFFECTS: If suspected adverse reaction occur, treatment should be discontinued and to inform veterinarian/local authority.

OVERDOSE AND TREATMENT: Bacitracin methylene disalicylate has a wide margin of safety. It has been administered to broiler and replacement chickens and swine at rates 20 times the highest approved use levels with no adverse effects.

STORAGE CONDITIONS: Keep it sealed, stored in room temperature (30°C) and dry place.

DOSAGE FORM: Soluble powder.

PACKAGING: 25kg/Drum and 10×100g/Bag.

WITHDRAWAL PERIODS: Chicken: zero days before slaughter. Swine: Not stated.

SHELF LIFE (after reconstitution in drinking water): 75 hours.

DATE OF REVISION: December 2021

MANUFACTURED BY:

Lifecome Biochemistry Co., Ltd

No.6, Industrial Park Avenue, Pucheng, Fujian, China

MARKETING AUTHORISATION HOLDER:

Age D'or Sdn Bhd

15, Jalan 31/54

Kota Kemuning

40460 Shah Alam

Selangor