



Important information. Please read carefully.

MucoKIDS™

125MG/5ML

Oral Solution

COMPOSITION

Each oral solution contains 125mg/5ml of carbocisteine.

PHARMACODYNAMICS

Carbocisteine is a mucolytic that alters the structure of mucus to decrease its viscosity thereby facilitating its removal by ciliary action or expectoration.

PHARMACOKINETICS

Absorption

Carbocisteine is rapidly and well absorbed from the gastrointestinal tract with peak plasma concentrations occurring about 2 hours after an oral dose.

Distribution

It appears to penetrate into lung tissue and respiratory mucus.

Metabolism

Acetylation, decarboxylation, and sulfoxidation have been identified as the major metabolic pathways.

Elimination

Carbocisteine is excreted in the urine as unchanged drug and metabolites.

INDICATIONS

MucoKids is used for the adjunctive therapy of respiratory tract disorders characterised by excessive, viscous mucus, including chronic obstructive airways disease.

DOSAGE AND ADMINISTRATION

For oral use only.

Children 2-5 years: 2.5ml - 5ml four times daily

Children 5-12 years: 10ml three times daily

CONTRAINDICATIONS

Contraindicated in children who:

- are under 2 years of age
- known hypersensitivity to carbocisteine
- have stomach or duodenum ulcers

WARNING AND PRECAUTIONS

Cough suppressant medicines or medicines that dry up bronchial secretions should not be taken while this medication is being used.

Carbocisteine should be used with caution in patients:

- with a history of peptic ulcer disease because of the risk that mucolytics may disrupt the gastric mucosal barrier.
- have chronic (long-term) illness of the bronchi or lungs
- with diabetes or on a low-sugar diet

DRUG INTERACTIONS

None stated. In order to prevent possible interactions between several medicines, the patient should inform doctor or pharmacist if they are taking any other medicines.

PREGNANCY AND LACTATION

Recommended to avoid in first-trimester pregnancy.

As a general rule, if you are pregnant or breast-feeding, you should always ask the advice of your doctor or pharmacist before taking any medicine.

SIDE EFFECTS

Nausea and gastric discomfort, and gastrointestinal bleeding have occasionally occurred with carbocisteine.

Immune System Disorders:

Anaphylactic/ anaphylactoid reaction

Skin and Subcutaneous Tissue Disorders:

Severe cutaneous adverse reactions (SCAR) e.g. erythema multiforme, Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN). In most of these cases reported at least one other drug was

administered at the same time, which may possibly enhanced the described mucocutaneous effects.

SYMPTOMS AND TREATMENT FOR OVERDOSAGE

Gastric lavage may be beneficial, followed by observation. Gastrointestinal disturbance is the most likely symptom of overdosage.

SHELF-LIFE

The expiry date is indicated on the packaging.

STORAGE CONDITION

Store below 30°C.

PRODUCT DESCRIPTION

Each bottle contains red syrup with vanilla and mixed berries flavour.

Available in pack size of 90ml and 120ml.

KEEP OUT OF REACH OF CHILDREN/

JAUHI DARI KANAK-KANAK

For further information, please consult your pharmacist or physician.

Revision Date: 27-Sept-2024

Manufactured and Marketed by

Xepa-Soul Pattinson (Malaysia) Sdn. Bhd. (8504-M)

1-5 Cheng Industrial Estate, 75250 Melaka, Malaysia.

KLF0060L1



001