


148 mm

75 mm

Pharmaco-
code
Read Direction

For dogs of more than 25 kg:

For easy application the dog should be standing. The entire contents of the pipette should be applied evenly as 3 or 4 spots along the top of the back, from between the shoulders to the base of the tail. At each spot, part the coat until the skin is visible. Wherever possible apply to undamaged skin. Place the tip of the pipette on the skin and gently squeeze the pipette to expel a portion of its contents directly onto the skin. Do not apply an excessive amount of solution at any one spot, as that could cause some of the veterinary medicinal product to run down the animal's side.



Contraindications

Do not use in puppies under 7 weeks of age.
Do not use in cases of hypersensitivity to the active substances or to any of the excipients.
Do not use in dogs classified as Class 4 for heartworm disease as the safety of the product has not been evaluated in this animal group.
Do not use in cats. Instead, the corresponding "Advocate Spot-on Solution for Cats" product (0.4 or 0.8 ml), which contains 100 mg/ml imidacloprid and 10 mg/ml moxidectin, must be used for cats.
Do not use in ferrets. Only "Advocate Spot-on Solution for Small Cats and Ferrets" (0.4 ml) must be used for ferrets.
Do not use on canaries.

Warnings and Precautions

Brief contact of the animal with water on one or two occasions between monthly treatments is unlikely to significantly reduce the efficacy of the veterinary medicinal product. However, frequent shampooing or immersion of the animal in water after treatment may reduce the efficacy of the veterinary medicinal product.
The possibility that other animals in the same household can be a source of re-infection with fleas, mites, gastrointestinal nematodes, heartworm and/or lungworm should be considered, and these should be treated as necessary with an appropriate product.
Unnecessary use of antiparasitics or use deviating from the instructions given in the SPC may increase the resistance selection pressure and lead to reduced efficacy. The decision to use the product should be based on confirmation of the parasitic species and burden, or of the risk of infestation based on its epidemiological features, for each individual animal.
Efficacy against adult *Dirofilaria repens* has not been tested under field conditions.
Special precautions for safe use in the target species:
The treatment of animals weighing less than 1 kg should be based on a benefit-risk assessment.
There is limited experience on the use of the veterinary medicinal product in sick and debilitated animals, thus the veterinary medicinal product should only be used based on a benefit-risk assessment for these animals.
Do not apply in the mouth, in the eyes or the ears of the animal.
Care should be taken that the veterinary medicinal product is not ingested by animals and does not come into contact with the eyes or mouth of the recipient and/or other animals.
Consider carefully the correct application method described in section Method of Administration, especially that the veterinary medicinal product should be applied to the site specified in order to minimise the risk for the animal to lick the veterinary medicinal product.
Do not allow recently treated animals to groom each other. Do not allow treated animals to come into contact with untreated animals until the application site is dry.
When the veterinary medicinal product is applied in 3 to 4 separate spots (see section Method of Administration), specific care should be taken to prevent the animal licking the application sites.
This veterinary medicinal product contains moxidectin (a macrocyclic lactone), therefore special care should be taken with Collie or Old English Sheep dogs and related breeds or crossbreeds, to correctly administer the veterinary medicinal product as described under section Method of Administration; in particular, oral uptake by Collie or Old English Sheep dogs and related breeds or crossbreeds should be prevented.
The safety of the veterinary medicinal product has only been evaluated in dogs classified as either Class 1 or 2 for heartworm disease in laboratory studies and in a few Class 3 dogs in a field study. Therefore, the use in dogs with obvious or severe symptoms of the disease should be based on a careful benefit-risk assessment by the treating veterinarian.
Although experimental overdosage studies have shown that the veterinary medicinal product may be safely administered to dogs infected with adult heartworms, it has no therapeutic effect against adult *Dirofilaria immitis*. It is therefore recommended that all dogs 6 months of age or more, living in areas endemic for heartworm, should be tested for existing adult heartworm infection before being treated with the veterinary medicinal product. At the discretion of the veterinarian, infected dogs should be treated with an adulticide to remove adult heartworms. The safety of the veterinary medicinal product has not been evaluated when administered on the same day as an adulticide.
Imidacloprid is toxic for birds, especially canaries.
Special precautions to be taken by the person administering the veterinary medicinal product to animals:
This veterinary medicinal product can cause skin, eye, or mouth irritation. In very rare cases the veterinary medicinal product may cause skin sensitisation or transient skin reactions (for example numbness, irritation or burning/tingling sensation).
In very rare cases the veterinary medicinal product may cause respiratory irritation in sensitive individuals.
People with a known hypersensitivity to benzyl alcohol, imidacloprid or moxidectin should administer the veterinary medicinal product with caution.
Avoid contact with skin, eyes or mouth.
Do not eat, drink or smoke during application.
Wash hands thoroughly after use.
After application do not stroke or groom animals until the application site is dry.
In case of accidental spillage onto skin, wash off immediately with soap and water.
If the veterinary medicinal product accidentally gets into eyes, they should be thoroughly flushed with water.
If skin or eye symptoms persist, or the veterinary medicinal product is accidentally swallowed, seek medical advice immediately and show the package leaflet or label to the physician.
Special precautions for the protection of the environment:
The veterinary medicinal product should not enter water courses as imidacloprid and moxidectin may be dangerous for fish and other aquatic organisms. Dogs should not be allowed to swim in surface waters for 4 days after treatment.

Other precautions:

The solvent in the veterinary medicinal product may stain or damage certain materials including leather, fabrics, plastics and finished surfaces. Allow the application site to dry before permitting contact with such materials.

Interaction with Other Medicinal Products and Other Forms of Interaction

During treatment with the veterinary medicinal product no other antiparasitic macrocyclic lactone should be administered.
No interactions between the veterinary medicinal product and routinely used veterinary medicinal products or medical or surgical procedures have been observed.
Safety of the veterinary medicinal product when administered on the same day as an adulticide to remove adult heartworms has not been evaluated.

Use During Pregnancy, Lactation or Lay

The safety of the veterinary medicinal product has not been established during pregnancy and lactation in the target species.

Pregnancy and lactation:

The use is not recommended during pregnancy and lactation.

Fertility:

Do not use in breeding animals.

Adverse Events

Dogs:

Common (1 to 10 animals / 100 animals treated):	Diarrhoea ¹ , Vomiting ¹ Cough ¹ , Dyspnoea ¹ , Tachypnoea ¹ Inappetence ¹ , Lethargy ¹
Rare (1 to 10 animals / 10,000 animals treated):	Vomiting
Very rare (< 1 animal / 10,000 animals treated, including isolated reports):	Application site greasy fur ² , Application site hair loss ² , Application site itching ² , Application site reddening ² Behavioural disorder (e.g. agitation) ³ Hypersalivation ⁴ Neurological signs (e.g. ataxia, muscle tremor) ⁵ Pruritus Inappetence ³ , Lethargy ³

- ¹ These signs are common in heartworm positive dogs with microfilaraemia, and there is a risk of gastrointestinal signs and severe respiratory signs that may require prompt veterinary treatment.
² These signs disappear without further treatment.
³ Transiently noted and related to sensation at application site.
⁴ This is not a sign of intoxication and disappears within minutes without treatment. Correct application will minimise licking of the application site.
⁵ Most neurological signs occur transiently.

Overdose

Up to 10 times the recommended dose was tolerated in adult dogs with no evidence of adverse effects or undesirable clinical signs. Five times the recommended minimum dose applied at weekly intervals for 17 weeks was investigated in dogs aged over 6 months and tolerated with no evidence of adverse effects or undesirable clinical signs.
The veterinary medicinal product was administered to puppies at up to 5 times the recommended dose, every 2 weeks for 6 treatments, and there were no serious safety concerns. Transient mydriasis, salivation, vomiting and transient rapid respiration were observed.
After accidental oral ingestion or overdose, neurological signs (most of which are transient) such as ataxia, generalised tremors, ocular signs (dilated pupils, little pupillary reflex, nystagmus), abnormal respiration, salivation and vomiting may occur in very rare cases.
Ivermectin-sensitive Collie dogs tolerated up to 5 times the recommended dose repeated at monthly intervals without any adverse effects, but the safety of application at weekly intervals has not been investigated in ivermectin-sensitive Collie dogs. When 40% of the unit dose was given orally, severe neurological signs were observed. Oral administration of 10% of the recommended dose produced no adverse effects.
Dogs infected with adult heartworms tolerated up to 5 times the recommended dose, every 2 weeks for 3 treatments, without any adverse effects.
In case of accidental oral uptake, symptomatic treatment should be administered. There is no known specific antidote. The use of activated charcoal may be beneficial.

Withdrawal Period

Not applicable.

Storage Condition

Do not store above 30°C.

Dosage forms and packaging available

Advocate Spot-on Solution for Dogs is available in triple pipettes per box for the following:
Advocate Spot-on Solution for Small Dogs
Advocate Spot-on Solution for Medium Dogs
Advocate Spot-on Solution for Large Dogs
Advocate Spot-on Solution for Extra-Large Dogs

Marketing Authorisation Holder:

Elanco Malaysia Sdn Bhd
Unit 5.04, Level 5 & 6, Tower Block, The Bousteador,
No. 10, Jalan PJU 7/6, Mutiara Damansara,
47800 Petaling Jaya, Selangor, Malaysia

Manufactured by:

KVP Pharma + Veterinär Produkte GmbH
Projensdorfer Straße 324, D-24106 Kiel, Germany.
Date of revision: Oct 2025

Advocate, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.
© 2025 Elanco or its affiliates

Elanco

297 mm