

Important Information. Please read carefully.

Aveflon

Tablet 500 mg

COMPOSITION

Each film-coated tablet contains Micronized Purified Flavonoid Fraction 500mg.
- Diosmin 90% (450mg)
- Hesperidin 10% (50mg)

PHARMACODYNAMICS

Micronized purified flavonoid fraction is a venotonic and vascular protector. It exerts an action on the vascular return system by reducing the venous distensibility and stasis (at the venous level) and normalises capillary permeability and increases capillary resistance (at the microcirculatory level).

PHARMACOKINETICS

Diosmin is synthesized from Hesperidin. Diosmin is highly metabolized and the elimination half-life is 11 hours. The excretion of Diosmin is mainly faecal, and about 14% of the dose administered is excreted in the urine.

INDICATIONS

Aveflon is used for the treatment of:

- Symptoms related to venolymphatic insufficiency, such as heavy legs, pain, early morning restless leg.
- Functional symptoms related to acute hemorrhoidal attack.

DOSAGE AND ADMINISTRATION

For oral use only.

Usual Dosage

2 tablets daily: 1 tablet at mid-day and 1 tablet in the evening at meal times.

Acute Hemorrhoidal Attack

First 4 days- 6 tablets daily

Subsequent 3 days- 4 tablets daily

CONTRAINDICATIONS

Contraindicated in patients with known hypersensitivity to micronized purified flavonoid fraction.

WARNINGS AND PRECAUTIONS

Acute Hemorrhoidal Attack

The administration of this product is not substituting the specific treatment of other anal disorders. The treatment must be short-term. If the symptoms do not disappear rapidly, proctological examination should be performed and the treatment should be reviewed.

DRUG INTERACTIONS

No clinically significant interactions is available with the use of micronized purified flavonoid fraction.

PREGNANCY AND LACTATION

Pregnancy

Experimental studies available have not shown any teratogenic effect in animals. No harmful effects have been reported to date in humans.

Lactation

In the absence of studies concerning excretion into breast milk, breastfeeding is not recommended during treatment.

SIDE EFFECTS

May cause minor gastrointestinal effects such as diarrhoea, abdominal pain, dyspepsia, nausea and vomiting. Rare adverse effects such as dizziness, headache, malaise, rash, pruritus and urticaria may also occur. Neurovegetative disorders have been reported with the use of micronized purified flavonoid fraction.



SYMPTOMS AND TREATMENT FOR OVERDOSAGE

No specific over dosage cases have been reported with the use of micronized purified flavonoid fraction. Overdose may lead to undesirable effects.

STORAGE CONDITION

Store below 30°C.

SHELF LIFE

The expiry date is indicated on the packaging.

PRODUCT DESCRIPTION, DOSAGE FORM AND PACKAGING AVAILABLE

Brown, oval shape, film coated tablet, with "AV" marking on one side and plain on the other side.

Available in packing of 10 film-coated tablets per blister, 3 blisters in a pack.

For further information, please consult your physician or pharmacist.

Revision Date: 03-Aug-2021

Manufacturer

Xepa-Soul Pattinson (Malaysia) Sdn Bhd

1-5 Cheng Industrial Estate 75250 Melaka, Malaysia.

Product Registration Holder

Apex Pharmacy Marketing Sdn Bhd

No.2, Jalan SS13/5, 47500 Subang Jaya, Selangor, Malaysia.

AV0900-003

08.21