

NEOFEED 490,000iu/g POWDER

Description and Composition

NEOFEED 490,000iu/g POWDER is a white to yellow white powder. Each gram contains 490,000iu Neomycin Sulphate.

Pharmacodynamics

Neomycin is an antibiotic of the aminoglycoside family, obtained from *Streptomyces fradiae*. Its spectrum of activity covers Gram-positive bacteria, particularly Staphylococci and less actively Streptococci, and Gram-negative bacteria, especially *Escherichia coli*. Neomycin binds to the 30S subunit of the bacterial ribosome, which disrupts the reading of the constitutive code of the messenger RNA, and finally the bacterial protein synthesis. At high concentrations, aminoglycosides have been shown to damage the bacterial wall, which adds bactericidal properties to bacteriostatic properties.

Pharmacokinetics

Neomycin is absorbed very little in the digestive tract. In plasma and tissues, neomycin concentrations are very low. It is therefore present in large quantities and persistently in the various sections of the digestive tract. Neomycin is almost exclusively eliminated in faeces after oral administration.

Indication

For treatment and prevention of gastrointestinal infections due to *Escherichia coli* in pigs, chickens, turkeys and ducks.

Recommended Dose

20,000iu neomycin per kg body weight per day, orally, for 10 days.

Incorporation rate

The amount of medicated feed consumed by animals depends on their physiological and clinical status. In order to comply with the recommended doses of neomycin and take into account the actual dietary intake, the incorporation rate may be increased or possibly decreased to the minimum incorporation limit of 200iu neomycin per gram of feed. This may lead to levels of neomycin in the medicated feed different from those mentioned below for information purposes.

Pigs: For daily feed intake of 50g of feed per kg body weight, this dosage corresponds to 400iu neomycin per gram of feed.

Chickens, turkeys and ducks: For daily feed intake of 100g of feed per kg body weight, this dosage corresponds to 200iu neomycin per gram of feed.

Route of Administration

For oral administration via feed.

Contraindication

Do not administer in case of known hypersensitivity to neomycin.

Warning and Precautions

Special warnings for each target species

This medicinal premix is intended exclusively for the target species.

Special precautions for use

Special precautions for use in animals

This medicinal premix is intended for the manufacture of solid medicinal feed and the rate of incorporation of the medicated premix into the solid feed cannot be less than 200iu neomycin per gram of feed.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Aminoglycosides may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Allergic reactions to these substances can be severe.

In case of known hypersensitivity to aminoglycosides, do not handle this product.

Handle this product with the recommended precautions to avoid exposure.

Wearing the mask, goggles and protective gloves is recommended when preparing the food. Avoid breathing dust and avoid contact with skin and eyes.

If symptoms such as rash appear, consult a physician and show the package leaflet.

Swelling in the face, lips or eyelids or difficulty in breathing are more serious symptoms and require medical attention.

Other precautions

None.

Interaction with Other Medicaments

Not known.

Pregnancy and Lactation

Safety in pregnant females has not been studied. However, due to very low absorption, the use of the product during gestation and lactation does not appear to pose any particular problem.

Side Effects

Not known.

Symptoms and Treatment of Overdose

Not known.

Storage Condition

Store below 30°C.

Shelf Life

3 years.

To be used within 24 hours after mixing into feed.

Withdrawal Periods

Pigs:

Meat and offal: 14 days

Chickens, turkeys and ducks:

Meat and offal: 14 days

Eggs: Zero days

Packing

10kg, 25kg.

Revised date: 5.3.2026

	Manufacturer & Product Registration Holder:
	Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 E-mail: info@rangepharma.com