

Avosil Lozenge

2mg

Cetylpyridinium Chloride 2mg

Composition

Each tablet (lozenge) contains:

Cetylpyridinium Chloride 2 mg

PRODUCT DESCRIPTION

A yellowish round tablet, debossed with "SP104" on one side.

INDICATION

Temporary relief of throat pain and inflammation.

PHARMACOLOGICAL PROPERTIES

ATC code: R02AA06

Pharmacotherapeutic group: Throat preparation, Antiseptic cetylpyridinium

Pharmacodynamic properties

Mechanism of action:

Cetylpyridinium Chloride is a quaternary ammonium compound with broad spectrum anti-bacterial activity. It is a cationic surfactant that absorbs readily to oral surfaces. Thus, it is of value in treating superficial infections of the mouth and throat.

The action of sucking the lozenge allows the active ingredient to work in the discomfort area and, also helps lubricate and soothe the painful area. This helps relieve the soreness and discomfort of mouth and throat infections

Pharmacokinetic properties

Not applicable.

RECOMMENDED DOSAGE

Route of administration: For oral use only.

Adult: 1 tablet 3 to 4 times daily

Children (above 6 years old): ½ tablet 3 to 4 times daily

Allow Avosil troche to dissolve slowly in the mouth. Suck the Avosil troche in the mouth, and avoid chewing or swallowing.

Taking Avosil before going to bed, the active ingredients can be highly maintained in saliva and gives better efficacy

CONTRAINDICATIONS

Avosil is contraindicated in those patients with a history of hypersensitivity to cetylpyridium or any of the other ingredients.

WARNING AND PRECAUTIONS

- If any form of allergies such as rash appears, immediately discontinue use of Avosil.
- Avosil has been shown to stimulate oral cavity and pharynx.
- Care to be taken for patient with diabetic; having intolerance to sucrose (sugar) due to the content of sucrose in Avosil; or taking or using other medicines
- Not to be used for children under 6 years old
- User to consult doctor or pharmacist if symptoms persist for more than 3 days

DRUG INTERACTIONS

None known.

PREGNANCY & LACTATION

The effects of Avosil in pregnancy and breast-feeding mothers are unknown.

SIDE EFFECTS

Medicines and their possible side effects can affect individual people in different ways. The following are some of the side effects that are known to be associated with this medicine. Just because a side effect is stated here, it does not mean that all people using this medicine will experience that or any side effect.

- Itchy rash (urticaria).
- Burning sensation in the mouth.

SIGNS AND SYMPTOMS OF OVERDOSE

There is no experience of overdosage but normal procedures of gastric lavage and maintenance of respiration and circulation (using vasopressor drugs if necessary) should apply.

STORAGE:

Store below 30°C. Protect from light and moisture.

Keep in a tightly closed container, protect from light. Keep out of reach of children.

PACKING:

20, 480 and 1000 tablets packed in a box of 2, 48, or 100 blisters of Aluminium/PVC blister; each blister containing 10 tablets. 2 blister pack into an aluminium bag

Product Registration Holder:

Nano Medic Care Sdn. Bhd
No. 1-6 & 1-12, 1st Floor, Jalan Indah 2, Taman Indah, 84000 Muar Johor Malaysia.

Manufactured by

Sinphar Pharmaceutical Co. Ltd. (Taiwan),
No. 84, Chung Shan Rd., Chung Shan Village, Tung-Shan Township, Yilan County 269, Taiwan R.O.C..

Distributed by:

Apex Pharmacy Marketing Sdn. Bhd
No. 2, Jalan SS 13/5, 47500 Subang Jaya, Selangor Malaysia.

Malaysia Registration No.: MAL18046131XC

Revision date: 2024.05.09