

PF DICLAZURIL PREMIX 0.5% W/W POWDER

Composition:

Diclazuril.....5mg

Exp. Qs.....1g

Description

White Powder

Pharmacodynamics

Diclazuril has potent anticoccidial activity even at low levels. It is effective against second-generation schizonts and sexual stages of *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, and *E. mivati*. Diclazuril is effective against *E. maxima* later in its life cycle. It also inhibits sporulation of oocyst. It is effective against *E. tenella* that has become resistant to other anticoccidial drugs.

Pharmacokinetics

The drug is poorly absorbed. Peak plasma concentration is attained within 6 hours in chicken

Indication

As an aid in the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mitis*, *E. necatrix* and *E. tenella* in broiler chickens.

Route of Administration

Oral

Recommended dose

Broiler Chickens: 1 mg/kg (milligram per kilogram) (0.0001%) of diclazuril in the complete feed. Feed this medicated feed as the sole ration from one day of age up until slaughter.

To obtain the recommended level of 0.0001% of diclazuril in the complete feed, PF Diclazuril Premix should be mixed thoroughly with the feed at the rate of 200 grams per metric tonne. It is recommended that a premix be made from PF Diclazuril Premix before final mixing. It is suggested to premix 200 grams of PF Diclazuril Premix with 20-50 kg of a feed ingredient (or vitamin-mineral mixture) before incorporating in the feed.

Contraindication

Hypersensitivity to diclazuril.

Interaction with other medicinal products

PF Diclazuril Premix should not be mixed or used simultaneously with any other anticoccidial.

Pregnancy and lactation

Not for use in hens producing eggs for human food.

Side effects

No reports of adverse effects at the recommended dose of 1 ppm in feed.

Symptoms and treatment of overdose

Chickens for fattening tolerated a 25-fold overdose of diclazuril for 37 days without any observable negative effects on performance, haematology, necropsy and histopathology

Storage

Store in a cool dry place (<30°C). Strictly avoid light and heat exposure.

Warnings and Precautions

Keep out of reach of children / Jauhi ubat dari kanak-kanak

1. Do not use diclazuril medicated feed for the treatment of outbreaks of coccidiosis.

2. If unexpected deaths occur, obtain an accurate diagnosis and follow veterinarian's or poultry pathologist's recommendations.

3. Do not use with pellet binding agents except Pel-Aid or Lignin Sulfonate.

Shelf life

36 months

Shelf life after opening

24 hours

Shelf life after dilution or constitution

24 hours

Withdrawal period

Chicken (broiler): 0 days

Not for use in hens producing eggs for human food.

Packaging

25kg

Revised date

22/5/2018

Product Registration Holder:

PANFAST MARKETING (M) SDN.BHD.

PENANG SCIENCE PARK 1102, MUKIM 13, LORONG PERINDUSTRIAN BUKIT MINYAK 18, KAWASAN PERINDUSTRIAN BUKIT MINYAK, 14100 SIMPANG AMPAT, PENANG, MALAYSIA
TEL: +604-5016868 FAX: +604-5016866

Manufactured by:

SHENNONG ANIMAL HEALTH (M) SDN.BHD.

No.6, 8, 10, Jalan Industri USJ 1/19, Taman Perindustrian USJ 1
47610 SUBANG JAYA, Selangor Darul Ehsan, Malaysia.
Tel: 03-8011 4646 Fax : 03-8024 1166