

Metaflor 20% w/v

For Animal Use Only

Oral Solution

CONTENT : Each ml contains:

Florfenicol.....200mg

PRODUCT DESCRIPTION: A yellow color, clear solution.

PHARMACODYNAMICS:

Florfenicol is a broad-spectrum synthetic antibiotics in the phenicol group that is active against most Gram-positive and Gram-negative bacteria isolated from domestic animals. Florfenicol acts by inhibition of protein synthesis at the ribosomal level and is bacteriostatic.

PHARMACOKINETICS:

After absorption and distribution, florfenicol is extensively metabolized by pigs and rapidly eliminated, primarily in urine. After parenteral dosing of florfenicol to pigs, it has been shown that lung concentrations are similar to serum concentrations.

INDICATION:

Broiler chickens:

For the treatment and control of air sacculitis associated with *E.coli* susceptible to florfenicol.

Swine:

For the treatment of swine respiratory disease associated with *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*.

DOSAGE & ADMINISTRATION:

For Broiler chickens: 100ppm (100mg/L) for 3 consecutive days in drinking water. This dosage rate is equivalent to 35mg/kg body weight of birds or 0.175ml Metaflor 20% w/v Oral Solution per kg body weight.

For Swine: 100ppm (100mg/L) for 5 consecutive days in drinking water. This dosage rate is equivalent to 10mg/kg body weight for pigs that are drinking 10% of their body weight or 0.05ml Metaflor 20% w/v Oral Solution per kg body weight.

CONTRAINDICATIONS:

Do not use in boars intended for breeding purposes.

Do not use in case of hypersensitivity to the active substance or to the excipients.

WARNINGS / PRECAUTIONS:

Special warnings for each target species

The treated pigs should be placed under special observation. On each of the five days of treatment, unmedicated drinking water should not be given until the full daily amount of medicated drinking water has been ingested by pigs.

If there are no signs of improvement after three days of treatment, the diagnosis should be reviewed and, if necessary, the treatment changed.

Special precautions for use in animals

The veterinary medicinal product should be used in conjunction with susceptibility testing.

Official and local antimicrobial policies should be taken into account when the product is used.

Treatment should not exceed 5 days.

Special precautions to be taken by the person administering the veterinary medicinal product to animals.

People with known hypersensitivity to florfenicol or polyethylene glycols should avoid contact with the veterinary medicinal product.

In case of accidental spillage onto skin rinse with water. In case of contact with eyes, rinse immediately with copious amounts of water.

Personal protective goggles should be worn when handling the veterinary medicinal product.

Other precautions

In order to prevent any adverse effects on algae and possible contamination of groundwater, manure from treated pigs must not be spread onto land without dilution with manure from untreated pigs. Manure from treated pigs must be diluted with at least 5 times the weight of manure from untreated pigs before it can be spread onto arable land.

To limit the development of antimicrobial resistance, the choice of this drug as the most appropriate treatment should be confirmed by clinical experience supported where possible by pathogen culture and drug susceptibility testing

DRUGS INTERACTIONS:

No data available

PREGNANCY AND LACTATION:

Studies in laboratory animals have not revealed any evidence of potential embryotoxin or foetotoxin effect of florfenicol. The safety of the veterinary medicinal product in sows has not been established during pregnancy and lactation. The use is not recommended during pregnancy and lactation.

Do not feed to birds intended for breeding purposes. The effects of product on the reproductive function of treated chickens have not been determined.

ADVERSE EFFECTS:

For swine: A slight reduction of water consumption by the animals, dark brown faeces and constipation may be observed during treatment.

Commonly observed adverse effects are diarrhea and/or peri-anal and rectal erythema/oedema which affect approximately 40% of the animal. These effects are transient. In a few of the affected animals, prolapse of the rectum that resolves without treatment may be observed.

Broiler chickens: None known.

OVERDOSE AND TREATMENT:

In case of overdosing, a decrease in weight gain, food and water consumption, peri-anal erythema and oedema and modification of some haematologi and biochemical parameter indicative of dehydration may be observed.

STORAGE CONDITIONS: Store at room temperature below 30°C. Protect from heat and sunlight exposure.

WITHDRAWAL PERIOD(S):

Broiler chickens: 5 days.

Swine: 21 days.

DOSAGE FORMS AND PACKAGING AVAILABLE:

1 Liter Oral Solution packed into HDPE bottle.

DATE OF REVISION OF PACKAGE INSERT:

8 May 2018.



Product Registration Holder:

NUTRI PHARMAX SDN. BHD.

(Co. No. 802189-M)

6, Lorong Serunai 1, Valdor Sungai Bakap,
14200 Sungai Jawi, Penang, Malaysia.
Tel : +604-582 9377

Manufactured by:

Nam Pharma Sdn Bhd

5, Lebuhr Perumahan Klebang 11
Taman Perindustrian Antarabangsa IGB
31200 Chemor, Perak