

PF TIAMULIN PREMIX 10% W/W POWDER

Composition:

Tiamulin Hydrogen Fumarate.....100mg
Exp. Qs.....1g

Description

White colour powder

Pharmacodynamic

Tiamulin is a bacteriostatic semi-synthetic antibiotic belonging to the pleuromutilin group of antibiotics and acts at the ribosomal level to inhibit bacterial protein synthesis.

Tiamulin has shown a in-vitro activity against a wide range of bacteria such as *Brachyspira hyodysenteriae*, *Brachyspira pilosicoli*, *Lawsonia intracellularis* and *Mycoplasma* spp.

Tiamulin is bacteriostatic at therapeutic concentrations and has been shown to act at the 70S ribosome level and the primary binding site is on the 50S subunit and possibly a secondary site where the 50S and 30S subunits join. It appears to inhibit microbial protein production by producing biochemical inactive initiation complexes, which prevent elongation of the polypeptide chain.

Mechanisms responsible for resistance development in *Brachyspira* spp to the pleuromutilin class of antibiotics are considered to be based on mutations at the ribosomal target site. Clinically relevant resistance to tiamulin requires combinations of mutations around the tiamulin binding site.

Resistance to tiamulin may be associated with decreased susceptibility to other pleuromutilins.

Pharmacokinetic

Pig

Tiamulin is well absorbed in the pig (over 90%) following oral administration and widely distributed through the body. Tiamulin has been shown to concentrate in the lung, a target tissue, and also in liver, where it is metabolised and excreted (70-85%) in the bile, the remainder is excreted via the kidney (15-30%). Tiamulin which has not been absorbed or metabolised, passes down the intestines to the colon and concentrates there.

Indication

Pig: For the treatment and prevention of swine dysentery caused by *Brachyspira hyodysenteriae*.

For the treatment of colitis caused by *Brachyspira pilosicoli*.

For the treatment of ileitis caused by *Lawsonia intracellularis*.

For the treatment of enzootic pneumonia caused by *Mycoplasma hyopneumoniae*.

Route of Administration

Oral

Recommended dose

Calculations to achieve the correct dose rate and achieve the correct inclusion rate should be based on: - Inclusion rate (ppm) = dose rate (mg/kg bwt) x bodyweight (kg) / daily feed intake (kg).

To ensure a correct dosage, bodyweight should be determined as accurately as possible to avoid underdosing.

The intake of medicated feed depends on the clinical condition of the animals. In order to obtain the correct dosage the concentration of tiamulin hydrogen fumarate has to be adjusted accordingly.

Pigs

Treatment of Swine Dysentery caused by *B. hyodysenteriae*, treatment of Porcine Colonic Spirochaetosis (colitis) caused by *B. pilosicoli*

Dosage: 5 - 10 mg tiamulin hydrogen fumarate/kg bodyweight daily administered for 7 to 10 consecutive days. The dosage will normally be achieved by an inclusion level of 100 – 200 ppm tiamulin hydrogen fumarate in the finished feed provided that feed intake is unaffected.

Amount of THF (mg/g) per premix formulation	Amount of premix formulation per one tonne of feed
100.0	1.0 – 2.0 kg

Prevention of Swine Dysentery caused by *B. hyodysenteriae*

Dosage: 2.0 mg tiamulin hydrogen fumarate/kg bodyweight daily. The dosage will normally be achieved by an inclusion level of 40 ppm tiamulin hydrogen fumarate in the finished feed, providing feed intake is unaffected. Preventive medication with tiamulin should be given for 2-4 weeks.

Preventive treatment with tiamulin should only be initiated after confirmed infection with *B. hyodysenteriae* and then as part of a program including measures aiming to eradicate or control the infection in the herd.

Amount of THF (mg/g) per premix formulation	Amount of premix formulation per one tonne of feed
100.0	0.4 kg

Treatment of Porcine Proliferative Enteropathy (ileitis) caused by *L. intracellularis*

Dosage: 7.5 mg tiamulin hydrogen fumarate/kg bodyweight daily administered for 10 to 14 consecutive days. The dosage will normally be achieved by an inclusion level of 150 ppm tiamulin hydrogen fumarate in the finished feed providing that feed intake is unaffected.

Amount of THF (mg/g) per premix formulation	Amount of premix formulation per one tonne of feed
100.0	1.5 kg

Treatment of Enzootic Pneumonia caused by *M. hyopneumoniae*

Dosage: 5.0 – 10.0 mg tiamulin hydrogen fumarate/kg bodyweight daily administered for 7 to 10 consecutive days. The dosage will normally be achieved by an inclusion level of 100 -200 ppm tiamulin hydrogen fumarate in the finished feed provided that feed intake is unaffected.

Secondary infection by organisms such as *Pasteurella multocida* and *Actinobacillus pleuropneumoniae* may complicate enzootic pneumonia and require specific medication.

Amount of THF (mg/g) per premix formulation	Amount of premix formulation per one tonne of feed
100.0	1.0 – 2.0 kg

Contraindication

Animals should not receive products containing ionophores (monensin, narasin or salinomycin) during or for at least seven days before or after treatment with tiamulin. Severe growth depression or death may result.

Refer to Interaction with other medicinal products section for information regarding interaction between tiamulin and ionophores

Interaction with other medicinal products

Tiamulin has been shown to interact with ionophores such as monensin, salinomycin and narasin and may result in signs indistinguishable from an ionophore toxicosis. Animals should not receive products containing monensin, salinomycin or narasin during or at least 7 days before or after treatment with tiamulin. Severe growth depression, ataxia, paralysis or death may result.

If signs of an interaction do occur, administration of contaminated feed should be stopped immediately. The feed should be removed and replaced with fresh feed not containing the anticoccidials monensin, salinomycin or narasin.

Pregnancy and lactation

Can be used in pigs during pregnancy and lactation.

Side effects

On rare occasions erythema or mild oedema of the skin may occur in pigs following the use of tiamulin.

Symptoms and treatment of overdose

Pigs: Single oral doses of 100 mg/kg bodyweight in pigs caused hyperpnoea and abdominal discomfort. At 150 mg/kg no CNS effects were noted except for sedation. At 55 mg/kg given for 14 days a transient salivation and slight gastric irritation occurred. A minimum lethal dose has not been established in the pig.

Storage

Store in a cool dry place (<30°C). Strictly avoid light and heat exposure.

Warning and Precaution

Keep out of reach of children / Jauhi ubat dari kanak-kanak

Special warnings for each target species

In case of reduced feed intake, the inclusion levels in feed may need to be increased to achieve target dosage. Acute cases and severely diseased animals with reduced feed intake should be treated with a product of suitable formulation such as an injectable or water solution.

Special precautions for use in animals

It is sound clinical practice to base treatment on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of target bacteria.

Refer to Interaction with other medicinal products for information regarding interaction between tiamulin and ionophores

Special precautions to be taken by the person administering the veterinary medicinal product to animals.

When mixing the veterinary medicinal product and handling the medicated feed, direct contact with eyes, skin and mucous membranes should be avoided. Personal protective equipment should be worn when mixing the veterinary medicinal product or handling the medicated feed: overalls, impervious gloves and either a disposable half-mask respirator or a non-disposable respirator. Wash contaminated skin.

In case of accidental ingestion, seek medical advice immediately and show the package insert or label to the physician.

People with known hypersensitivity to tiamulin should administer the product with caution

Shelf life

36 months

Shelf life after dilution or constitution

24 hours

Shelf life after opening

24 hours

Packaging

25kg

Withdrawal period

Pigs

Revised date

25/2/2018

Prevention (at 2.0 mg/kg bwt): 1 day

Treatment (at 5-10 mg/kg bwt): 6 days

Product registration holder:

PANFAST MARKETING (M) SDN.BHD.

PENANG SCIENCE PARK 1102, MUKIM 13, LORONG PERINDUSTRIAN BUKIT

MINYAK 18, KAWASAN PERINDUSTRIAN BUKIT MINYAK,

14100 SIMPANG AMPAT, PENANG, MALAYSIA

TEL: +604-5016868 FAX: +604-5016866

Manufactured by:

SHENNONG ANIMAL HEALTH (M) SDN.BHD.

No.6, 8, 10, Jalan Industri USJ 1/19, Taman Perindustrian USJ 1

47610 SUBANG JAYA, Selangor Darul Ehsan, Malaysia.

Tel: 03-8011 4646 Fax : 03-8024 1166