

PF FLORFENICOL PREMIX 20% W/W POWDER

Composition:

Florfenicol.....200mg

Exp. Qs.....1g

Description

White colour powder

Pharmacodynamic

Florfenicol is a broad-spectrum synthetic antibiotic in the phenicol group that is active against most Gram-positive and Gram-negative bacteria isolated from domestic animals. Florfenicol acts by inhibition of protein synthesis at the ribosomal level and is bacteriostatic.

In-vitro testing has shown that florfenicol is active against the bacterial pathogens most commonly isolated in respiratory diseases in pigs, including *Pasteurella multocida*.

Pharmacokinetic

Florfenicol is well absorbed when administered orally and following distribution it is rapidly eliminated in the urine and faeces in a ratio of 3:1. A fraction is excreted unchanged and the rest is metabolised into 5 major metabolites.

After parenteral dosing of florfenicol to pigs, it has been shown that lung concentrations are similar to serum concentrations.

Indication

For the treatment and prevention of swine respiratory disease caused by *Pasteurella multocida* susceptible to florfenicol in infected herds. The presence of the disease should be established in the herd before initiating preventive treatment.

Route of Administration

Oral

Recommended dose

To be administered orally, in medicated feeding stuff.

Dosage:

10 mg of florfenicol per kg body weight (bw) (equivalent to 50 mg PF Florfenicol Premix) per day administered for 5 consecutive days.

Administration:

For a daily feed intake of 50g/kg body weight, this dosage corresponds to a rate of incorporation of 1 kg of premix per ton of feed, i.e. 200 ppm of florfenicol.

The rate of incorporation of the medicated premix in the feed may be increased in order to achieve the required dosage on a mg/kg body weight basis and to take into account the actual feed intake. Thus, the inclusion level may need adjusting as follows to give the correct dose.

$$\frac{50 \text{ mg PF Florfenicol Premix per kg body weight and day} \times \text{Average pig body weight (kg)}}{\text{Average daily feed intake (kg/animal)}} = \text{mg PF Florfenicol premix per kg of feed}$$

In all cases the recommended dose of 10mg of florfenicol per kg of body weight per day, for 5 consecutive days has to be respected.

To ensure a correct dosage body weight should be determined as accurately as possible to avoid underdosing. The required doses should be measured by suitably calibrated weighing equipment.

A horizontal ribbon mixer should be used to incorporate the product into the feeding stuff. It is recommended that the product is added to the mixer containing the feeding stuff ingredients and mixed thoroughly to produce a homogeneous medicated feeding stuff. Medicated feed may also then be pelleted. Pelleting conditions include a pre-conditioning step with steam and then the mixture is passed through a pelleter or extruder under normal conditions.

Contraindication

Do not administer to boars intended for breeding.

Do not use in case of hypersensitivity to the active substance or any of the excipients.

Interaction with other medicinal products

None known.

Pregnancy and lactation

The safety of the product during pregnancy and lactation has not been studied in sows. Use of the product during pregnancy and lactation is therefore not recommended.

Side effects

Commonly observed adverse effects are diarrhoea perianal inflammation and rectal eversion. Increased serum calcium may also be observed. These effects are transient, resolving on cessation of treatment.

Symptoms and treatment of overdose

In the event of overdose, a reduction in food and water consumption, together with a decrease in bodyweight may be observed. There may be an increase in refused feed and an increase in serum calcium.

Storage

Store in a cool dry place (<30°C). Strictly avoid light and heat exposure.

Warning and Precaution

Keep out of reach of children / Jauhi ubat dari kanak-kanak

Special warnings for each target species

Animals showing a decreased appetite and/or a poor general condition should be treated by the parenteral route.

Special precautions for use in animals

The product should be used in conjunction with susceptibility testing and take into account official and local policy relating to the use of antimicrobials.

Treatment should not exceed 5 days.

In a field clinical study, within a week after the administration of the last dose, the incidence of pigs presenting either mild depression and/or mild dyspnea and/or pyrexia (40°C) was approx. 20 % in the initially severely ill animals.

Special precautions to be taken by the person administering the medicinal product to animals

Skin sensitisation may occur.

Avoid skin contact.

Do not handle this product in case of known sensitization to active substance or any of the excipients.

Handle this product with care to avoid exposure during incorporation of premix into feed and administration of feed to animals, taking all recommended precautions.

Wear either a disposable half-mask respirator conforming to European standard EN 149 or a non-disposable respirator to European Standard EN 140 with a filter to EN 143, chemically resistant gloves, protective coveralls and goggles while incorporating the premix into feed.

Wear gloves and do not smoke, eat, or drink when handling the product or medicated feed.

Wash hands thoroughly with soap and water after use of the product or medicated feed

Rinse thoroughly with water in case of exposure.

If you develop symptoms following exposure such as skin rash, you should seek medical advice and take the package leaflet or the label with you.

Other precautions

Manure from treated swine must be stored for a minimum of one month before being spread and incorporated in fields.

Shelf life

36 months

Shelf life after opening

24 hours

Shelf life after dilution or constitution

24 hours

Withdrawal period

Meat and offal: 14 days

Packaging

25kg

Revised date

02/04/2018

Distributed by:

PANFAST MARKETING (M) SDN.BHD.

PENANG SCIENCE PARK 1102, MUKIM 13, LORONG PERINDUSTRIAN BUKIT
MINYAK 18, KAWASAN PERINDUSTRIAN BUKIT MINYAK,
14100 SIMPANG AMPAT, PENANG, MALAYSIA
TEL: +604-5016868 FAX: +604-5016866

Manufactured by:

SHENNONG ANIMAL HEALTH (M) SDN.BHD.

No.6, 8, 10, Jalan Industri USJ 1/19, Taman Perindustrian USJ 1
47610 SUBANG JAYA, Selangor Darul Ehsan, Malaysia.
Tel: 03-8011 4646 Fax : 03-8024 1166