

For Animal Use Only

Metaenro 20% w/v

Oral Solution

CONTENT: Each Liter contains:
Enrofloxacin.....200gm

PRODUCT DESCRIPTION: A yellow colour, clear solution.

PHARMACODYNAMICS:

Two enzymes essential in DNA replication and transcription, DNA gyrase and topoisomerase IV, have been identified as the molecular targets of fluoroquinolones. They modulate the topological state of DNA through cleaving and resealing reactions. Initially, both strands of the DNA double helix are cleaved. Then, a distant segment of DNA is passed through this break before the strands are resealed. Target inhibition is caused by non-covalent binding of fluoroquinolone molecules to an intermediate state in this sequence of reactions, in which DNA is cleaved, but both strands are retained covalently attached to the enzymes. Replication forks and translational complexes cannot proceed beyond such enzyme-DNA-fluoroquinolone complexes, and inhibition of DNA and mRNA synthesis triggers events resulting in a rapid, drug concentration-dependant killing of pathogenic bacteria.

PHARMACOKINETICS:

Enrofloxacin administered via drinking water to poultry is rapidly and very well absorbed with a bioavailability of approx. 90 %. Maximum plasma concentrations of 2 mg/L are reached within 1.5 hours after a single bolus dose rate of 10 mg/kg body weight. Concentrations in target tissues like lungs, liver, kidney, intestine and muscle tissue, exceed plasma concentrations by far. In poultry enrofloxacin is poorly metabolised to its active metabolite ciprofloxacin (approximately 5%). Enrofloxacin is eliminated from the body at a half-life of 6 hours. Protein binding in poultry is approximately 25%.

INDICATION:

Treatment of infections caused by the following bacteria susceptible to enrofloxacin:

Chickens

Mycoplasma gallisepticum,
Mycoplasma synoviae,
Avibacterium paragallinarum,
Pasteurella multocida,
Escherichia coli.

Turkey

Mycoplasma gallisepticum,
Mycoplasma synoviae,
Pasteurella multocida,
Escherichia coli.

Enrofloxacin should be used where clinical experience, supported where possible by sensitivity testing of the causal organism, indicates enrofloxacin as the active substance of choice.

DOSAGE & ADMINISTRATION:

Chicken and Turkey:

10 mg enrofloxacin/kg bodyweight per day for 3-5 consecutive days.

Treatment for 3-5 consecutive days; for 5 consecutive days in mixed infections and chronic progressive forms. If no clinical improvement is achieved within 2-3 days, alternative antimicrobial therapy should be considered based on susceptibility testing. Via the drinking water. Always make sure that the entire dose offered has been consumed. The medicated water should be made up fresh each day just before it is offered to the animals. The drinking water must be medicated throughout the treatment period, and no other water source should be available. Determine the bodyweight of the birds as accurately as possible in order to avoid underdosing. Use only fresh pre-solutions, prepared every day before start of treatment. Pumping systems should be checked constantly to assure proper medication. Empty the water system and fill it with medicated water before starting the treatment.

Calculate the daily quantity (ml) of Metaenro 20% w/v Oral Solution required for treatment period as follows:

Total number of birds x Average body weight in kg x 0.05 = Total volume (ml) per day

Metaenro 20% w/v Oral Solution may be put directly into the header tank or introduced via a water proportioner pump.

CONTRAINDICATIONS:

Do not use for prophylaxis.

Do not use when resistance/ cross-resistance to (fluoro)quinolones is known to occur in the flock intended for treatment

Do not use in the case of known hypersensitivity to the active substance, other (fluoro)quinolones or to any of the excipients.

WARNINGS / PRECAUTIONS:

- Treatment of *Mycoplasma* spp infections may not eradicate the organism.

Special precautions for use

Special precautions for use in animals.

- Official and local antimicrobial policies should be taken into account when the product is used.

- Fluoroquinolones should be reserved for the treatment of clinical conditions which have responded poorly, or are expected to respond poorly, to other classes of antimicrobials.

Special precautions to be taken by the person administering the medicinal product to animals.

- Those with known hypersensitivity to (fluoro)quinolones should avoid contact with this product.

- Avoid contact with skin and eyes.

- Rinse any splashes from skin or eyes immediately with water.

- Wash hands and exposed skin after use.

- Do not eat, drink or smoke while using the product.

- TO BE PRESCRIBED BY REGISTERED VETERINARY SURGEONS ONLY

DRUGS INTERACTIONS:

In vitro, an antagonism was shown, when combining fluoroquinolones with bacteriostatic antimicrobial agents such as macrolides or tetracyclines and phenicols. The simultaneous application of substances containing aluminium or magnesium can impair the absorption of enrofloxacin.

PREGNANCY AND LACTATION:

Do not use in laying hens producing eggs for human consumption.

Do not administer to layer replacement birds within 14 days of coming into lay.

ADVERSE EFFECTS:

None known.

OVERDOSE AND TREATMENT:

No adverse clinical symptoms were observed in chickens and turkeys treated respectively with doses up to 10 and 6 times higher than the therapy dose.

The use of fluoroquinolones during the growth phase combined with a marked and prolonged increase in the intake of drinking water, and hence active ingredient, possibly due to high temperatures, may potentially be associated with damage of the articular cartilage.

STORAGE CONDITIONS:

Store at room temperature below 30°C. Protect from heat and sunlight exposure.

WITHDRAWAL PERIOD(S):

Chickens : Meat and offal : 7 days.

Turkeys : Meat and offal : 13 days.

Not authorised for use in birds producing eggs for human consumption.

Do not administer to layer replacement birds within 14 days of coming into lay.

DOSAGE FORMS AND PACKAGING AVAILABLE: 1 Liter Oral Solution packed into HDPE bottle.

DATE OF REVISION OF PACKAGE INSERT: 20 February 2018

TO BE PRESCRIBED BY REGISTERED VETERINARY SURGEONS ONLY



Product Registration Holder:
NUTRI PHARMAS SDN. BHD.
(Co. No. 802189-M)
6, Lorong Serunai 1, Valdor Sungai Bakap,
14200 Sungai Jawi, Penang, Malaysia.
Tel : +604-582 9377

Manufactured by:
Nam Pharma Sdn Bhd
5, Lebuhr Perumahan Klebang II
Taman Perindustrian Antarabangsa IGB
31200 Chemor, Perak