

# Nicarmix 25

## Granules

### MEDICATED PREMIX

#### FORMULA:

Each kg of premix contains

Nicarbazin .....250g

Excipients up to .....1kg

#### PRODUCT DESCRIPTION:

It is a granulated product, yellow to dark yellow color, with occasional presence of white particles, free from visible impurities and/or agglomerated material.

#### PHARMACODYNAMICS:

Nicarbazin's mechanism of action is unknown; however, it is thought to be via inhibition of succinate-linked NAD (nicotinamide adenine dinucleotide) reduction and the energy-dependent transhydrogenase, and the accumulation of  $\text{Ca}^{2+}$  in the presence of ATP (Adenosine triphosphate).

#### PHARMACOKINETICS:

Nicarbazin is a mixture of 4,4'-dinitrocarbanilide (DNC) and 2-hydroxy-4,6-dimethylpyrimidine (HOP). DNC and HOP are absorbed separately from the digestive tract. DNC is absorbed more rapidly but disappears more slowly from the tissues than HOP.

2-hydroxy-4,6-dimethylpyrimidine (HOP) also known as 4,6-dimethyl-2-pyrimidinol (DHP or HDP).

#### INDICATION:

As an aid in preventing outbreaks of cecal (*Eimeria tenella*) and intestinal (*E. acervulina*, *E. maxima*, *E. necatrix* and *E. brunetti*) coccidiosis in broilers.

#### RECOMMENDED DOSE:

Uniformly mix 500 grams of NICARMIX 25 per M.T of feed to provide 125 ppm of Nicarbazin in the finished feed.

Feed continuously as sole ration from time chicks are placed on litter until past the time when coccidiosis is ordinarily a hazard.

#### MODE OF ADMINISTRATION:

Route of Administration: ORAL

Mix thoroughly in feed.

#### CONTRAINDICATIONS:

Do not use as a treatment for outbreaks of coccidiosis.

#### WARNINGS AND PRECAUTIONS:

Do not use as a treatment for coccidiosis.

Do not use in flushing mashers.

Do not feed to laying hens.

#### INTERACTIONS WITH OTHER MEDICAMENTS:

Not known.

#### STATEMENT ON USAGE DURING PREGNANCY AND LACTATION:

Do not feed to laying hens.

**ADVERSE EFFECTS / UNDESIRABLE EFFECTS:**

- (a) Nicarbazin may bleach brown-shelled eggs, cause mottled egg yolks and poor hatchability, and impair egg production.
- (b) Medicated broilers may be more susceptible to heat stress.

**OVERDOSE AND TREATMENT:**

Not known. If overdose is confirmed, remove the feed containing nicarbazin and provide proper access to drinking water to the birds.

**WITHDRAWAL PERIOD(S):**

The withdrawal period in broilers is 4 (FOUR) days.

**STORAGE CONDITIONS:**

Keep product in dry and aired environment below 30°C, protected against moisture and sunlight.

**SHELF LIFE:**

3 years.

**SHELF LIFE AFTER RECONSTITUTION OF PRODUCT INTO FEED:**

24 hours.

**DOSAGE FORMS AND PACKAGING AVAILABLE:**

Medicated granules premix  
25 kg bag

Manufactured by  
Phibro Saúde e Nutrição Animal Ltda.  
Rua Professor Gilberto Venturini, 555,  
Cidade Planejada II, CEP 12922-735,  
Bragança Paulista/SP, Brasil

Registration holder:  
Phibro Corporation (M) Sdn Bhd  
311, Level 3, Menara Axis,  
2, Jalan 51A/223, Seksyen 51A,  
46100 Petaling Jaya, Selangor, Malaysia

Revision Date of Package Insert: Aug 15, 2025