

YH BACITRACIN ZINC 15%W/W POWDER

ACTIVE CONSTITUENT: Bacitracin (As bacitracin zinc) 150g/kg. 1mg bacitracin equivalent to 42iu.

PRODUCT DESCRIPTION: Light brown or tan powder.

ROUTE OF ADMINISTRATION: Oral administration.

INDICATIONS&RECOMMENDED DOSAGE:

Indications: For the treatment of Necrotic Enteritis (*Clostridium perfringens* Types A and C) and as an aid in the prevention of Necrotic Enteritis in poultry.

Dosage and Administration: Should be premixed with 10-20 kg of feed and then thoroughly mixed with the bulk feed.

Aid in the prevention of Necrotic Enteritis: Add YH Bacitracin Zinc 15% W/W Powder at the rate of 267 g/tonne of finished feed (corresponding to 40g bacitracin activity per tonne). Medicated feed should be fed to poultry during the period when they are at most risk of the disease, which is usually between 2 and 12 weeks.

Treatment of Necrotic Enteritis: Add YH Bacitracin Zinc 15% W/W Powder at the rate of 1.33kg/tonne of finished feed (corresponding to 200g bacitracin activity per tonne) to be fed for 5-7 days.

PHARMACODYNAMICS &PHARMACOKINETIC:

Bacitracin is an antibiotic from the group of peptide antibacterial compounds. Bacitracin is produced by *Bacillus licheniformis* and *Bacillus subtilis*, and is a mixture of several closely related polypeptides, mainly consisting of bacitracin A, B1, B2, C and a small portion of bacitracin F. Bacitracin is mainly active against gram-positive microorganisms. The activities of each of bacitracins B1, B2 and C constitute about 90% of the antibacterial activity of bacitracin A. Bacitracin F is almost inactive. After oral application to chickens, bacitracin is hardly absorbed from the gastrointestinal tract, and the distribution to organs and tissues is negligible. In chickens, approximately 95% of an oral dose is excreted via faeces, and only 3% or less via urine. Bacitracin is metabolised to amino acids and smaller peptides via the main metabolite desamido-bacitracin, which is microbiologically inactive. Main metabolites in faeces are bacitracin A, B1, B2, F, desamido-bacitracin and catabolic peptides. In urine and bile only hydrolytic cleavage products (di- and tripeptides) are present.

WARNINGS AND PRECAUTIONS:

Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may require extending the approved withholding period.

INTERACTION WITH OTHER MEDICINAL PRODUCT: None Known

PREGNANCY AND LACTATION: None known

OVERDOSE AND TREATMENT: Stop medication and to seek for veterinarian advice if there is sign of over dosage.

STORAGE CONDITION: Store in room temperature (30°C) and dry place. Avoid light and close tightly.

CONTRAINDICATION: Do not use in known cases of hypersensitivity to active ingredient.

SIDE EFFECTS: If suspected adverse reactions occur, treatment should be discontinued and to inform veterinarian/local authority.

WITHDRAWAL PERIOD: Meat: Nil; Eggs: Nil

PACKAGING: 25kg/bag.

SHELF LIFE AFTER RECONSTITUTION OR DILUTION: 6 months

SHELF LIFE: 2 years

DATE OF REVISION: February 2026

MANUFACTURED BY:

Lifecome Biochemistry Co., Ltd

Add.: No.1, Putan Avenue, Putan Industrial Park,
Pucheng, Fujian, China.

MARKETING AUTHORISATION HOLDER:

Yenher Agro-Products Sdn.Bhd.,

Add.: 1628, Jalan IKS Simpang Ampat1. Taman IKS Simpang Ampat,
14100 Simpang Ampat. Seberang Perai Selatan, Penang , Malaysia