

Package Inserts

I.	Brand or Product Name: FLOR-LD 200MG/ML Oral Solution		Do not use in boars intended for breeding purpose. Studies in rats have revealed evidence of potential adverse effects on the male reproductive system.	XIII.	Side Effects: A slight reduction of water intake by the animals, dark brown faeces and constipation may be observed during treatment.
II.	Name and Strength of Active Substance: Florfenicol, 200mg in 1mL	X.	Warning and Precautions: <u>Special warning for each target species</u> The treated pigs should be placed under special observation. On each of the five days of treatment, unmedicated drinking water should not be given until the full daily amount of medicated drinking water has been ingested by pigs. If there are no signs of improvement after three days of treatment, the diagnosis should be reviewed and if necessary, the treatment changed. In case of insufficient water intake, animals should be treated parentally. <u>Special precautions for use in animals:</u> Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. <u>Special precautions to be taken by the person administering the veterinary medicinal product to animals.</u> This product may cause hypersensitivity (allergy). People with known hypersensitivity to florfenicol, polyethylene glycol 400, polysorbate 80 or N-methylpyrrolidone Pregnant women and women of child-bearing age should avoid working with this product. Contact of the product or the medicated drinking water with skin and eyes should be avoided. Personal protective equipment consisting of protective gloves, coverall and safety glasses should be worn when handling and mixing the product. Do not smoke, eat or drink when handling the product or mixing the medicated drinking water. In case of accidental spillage into eyes, wash them immediately with water. In case of contact with skin, wash the affected area immediately and remove any contaminated clothing. If you develop symptoms following exposure such as skin rash, seek medical advice and take the package leaflet or the label with you. Wash hands after use.	XIV.	Symptoms and Treatment of Overdose: In case of overdosing, a decrease in weight gain, water intake, peri-anal erythema and oedema and modification of some haematological and biochemical parameters indicative of dehydration may be observed.
III.	Product Description: A yellow clear solution.			XV.	Withdrawal Period: Meat and Offal is 23 days.
IV.	Target Species: Pig			XVI.	Storage Condition: Stored at ≤ 30°C / store below 30°C.
V.	Pharmacodynamics: Florfenicol is a derivative of thiamphenicol, in which the hydroxyl group has been replaced with fluorine. This makes it effective against chloramphenicol resistant, acetyl transferase producing bacteria. Laboratory tests have confirmed the activity of florfenicol against <i>Actinobacillus pleuropneumoniae</i> and <i>Pasteurella multocida</i> in swine.			XVII.	Proposed Shelf Life: As packaged for sale: 2 years After first opening of the immediate packaging: 28 days After dilution or reconstitution: 24 hours
VI.	Pharmacokinetics: Florfenicol is well distributed to most body tissues. The maximum concentration is reached in kidney, liver, bladder, lung and in intestines. Approximately 50% of florfenicol is excreted from the organism unchanged. The remaining part is excreted as a metabolite (mainly florfenicol amine).			XVIII.	Dosage Forms and packaging available: In liquid form, packed in 1L container.
VII.	Indication: Treatment and metaphylaxis of swine respiratory disease associated with <i>Actinobacillus pleuropneumoniae</i> and <i>Pasteurella multocida</i> susceptible to florfenicol. The presence of the disease in the herd should be established before initiating metaphylaxis.			XIX.	Name and Address of manufacturer NAM PHARMA SDN. BHD. 5, Lebuhr Perusahaan Klebang 11, Taman Perindustrian Antarabangsa IGB, 31200 Chemor, Perak, Malaysia.
VIII.	Recommended Dose and Administration: For in drinking water use. The recommended dose is 10 mg florfenicol per kg body weight daily (corresponding to 5 ml of the product / 100 kg b.w.) given for 5 consecutive days. The oral suspension must be shaken before use.	XI.	Interaction with other medicinal product: No data available	XX.	Name and Address Product holder KTY CHEMICALS SDN BHD 65, Jalan Cenderawasih 3, Taman Cenderawasih Indah, 14300 Nibong Tebal, SPS, Penang, Malaysia.
IX.	Contraindication: Do not use in case of hypersensitivity to the active substance or to any of the excipients.	XII.	Pregnancy and Lactation: As no studies have been performed in pigs during pregnancy and lactation, do not use the product in pregnant or lactating animals.	XXI.	Date of Revision of Package Insert: 04/04/2018