

Product:	Esberitox Compact	Fontsize:	24 pt; 8 pt	5. proof:	17.02.2017
Country:	MALAYSIA (MAL)	Used Font:	ITC_Eras Std; Frutiger LT Std.	6. proof:	14.03.2017
Packaging:	patient information leaflet	Colours:	Black	7. proof:	02.06.2017
Manufacturer:	Schaper & Brümmer GmbH & Co. KG			8. proof:	29.08.2017
Drawing No:	9908/019/00	1. proof:	22.11.2016	10. proof:	15.01.2018
Pharmacode:	99080	2. proof:	01.12.2016	11. proof:	15.01.2018
Dimension:	148 x 210 mm	3. proof:	12.01.2017	12. proof:	31.01.2018
Page:	1 of 2	4. proof:	26.01.2017	13. proof:	02.02.2018
		9. proof:	07.11.2017	14. proof:	27.04.2018
☐ Correction required		☐ Approval		15. proof:	03.05.2018
☐ Final Version/Date:				16. proof:	09.01.2019
				17. proof:	04.02.2019

Package leaflet: Information for the user

Esberitox[®] COMPACT

T A B L E T S

for application in adults and adolescents from the age of 12 years

dry extract from a mixture of wild indigo rootstock, purple coneflower root, pale coneflower root and thuja herba

Read all of this package leaflet carefully before you start taking this medicine because it contains important information for you.

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist have told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- You must talk to your doctor or pharmacist if you do not feel better or if you feel worse after 7 days.

What is in this leaflet

1. What Esberitox[®] COMPACT is and what it is used for
2. What you need to know before you take Esberitox[®] COMPACT
3. How to take Esberitox[®] COMPACT
4. Possible side effects
5. How to store Esberitox[®] COMPACT
6. Contents of the pack and other information

1. What Esberitox[®] COMPACT is and what it is used for

Esberitox[®] COMPACT is a traditional medicine used for the relief of cold.

2. What you need to know before you take Esberitox[®] COMPACT

2.1 Do not take Esberitox[®] COMPACT

- if you are allergic to the pharmacologically active ingredients (wild indigo rootstock, purple coneflower root, cone flower root, thuja herba) or any of the other ingredients of Esberitox[®] COMPACT or to composites.

2.2 Warnings and precautions

The use is not recommended in cases of progressive systemic disorders, autoimmune diseases, immunodeficiencies, immunosuppression and diseases of the white blood cell system.

2.3 Children:

This medicine is not suitable for children under the age of 12 years.

2.4 Other medicines and Esberitox[®] COMPACT

Interactions are not known.

Please nevertheless inform your doctor or pharmacist, if you take or have recently taken other medicines, including medicines obtained without a prescription.

2.5 Pregnancy and breast-feeding:

Contraindicated in pregnant women. Insufficient reliable data in breastfeeding women.

2.6 Driving and using machines:

No special precautions are required.

3. How to take Esberitox[®] COMPACT

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

3.1 Unless otherwise prescribed by the doctor, the recommended dose is:

Adults and adolescents from the age of 12 years: Take **1 tablet** each time after meal. To be taken **3 times daily**.

Product:	Esberitox Compact	Fontsize:	24 pt; 8 pt	5. proof:	17.02.2017
Country:	MALAYSIA (MAL)	Used Font:	ITC_Eras Std; Frutiger LT Std.	6. proof:	14.03.2017
Packaging:	patient information leaflet	Colours:	Black	7. proof:	02.06.2017
Manufacturer:	Schaper & Brümmer GmbH & Co. KG			8. proof:	29.08.2017
Drawing No:	9908/019/00	1. proof:	22.11.2016	10. proof:	15.01.2018
Pharmacode:	99080	2. proof:	01.12.2016	11. proof:	15.01.2018
Dimension:	148 x 210 mm	3. proof:	12.01.2017	12. proof:	31.01.2018
Page:	2 of 2	4. proof:	26.01.2017	13. proof:	02.02.2018
		9. proof:	07.11.2017	14. proof:	27.04.2018
☐ Correction required		☐ Approval		15. proof:	03.05.2018
☐ Final Version/Date:				16. proof:	09.01.2019
				17. proof:	04.02.2019

Please take Esberitox® COMPACT in the morning, in the middle of the day and in the evening with sufficient liquid, preferably 1 glass of water. The tablets may also be chewed.

3.2 Duration of use:

Start the treatment as early as possible after the first symptoms appear and continue taking Esberitox® COMPACT until the symptoms disappear. However, take note of the details on precautionary measures of section 2.2 and the details on side effects of section 4 in this package leaflet.

Do not take Esberitox® COMPACT longer than 10 days without medical advice.

3.3 If you take more Esberitox® COMPACT than you should:

Toxications with Esberitox® COMPACT have not been known until now. If you inadvertently have taken one or two tablets more than you should, this has normally no adverse effects. If you have taken a considerably larger quantity of Esberitox® COMPACT than you should, please consult a doctor. He/she can decide on required measures to be taken, if necessary.

3.4 If you forget to take Esberitox® COMPACT:

Do not take a double dose, but continue the intake as described in this package leaflet or as prescribed by your doctor.

4. Possible side effects

- Hypersensitivity reactions, e. g. skin rash, itching, facial swelling, shortness of breath, blood pressure decrease
- Abdominal pain, vomiting, diarrhoea
- Dizziness

5. How to store Esberitox® COMPACT

Store below 30 degree C. Protect from light and moisture.

Keep away from children. Jauhkan dari kanak-kanak.

6. Contents of the pack and other information

What Esberitox® COMPACT contains:

The active substance is:
Each 550 mg tablet contains:
16 mg extract (as dry extract) from Baptisia tinctoria L. (wild indigo), radix corresponding to approx. 53 mg wild indigo rootstock; Echinacea purpurea Moench (purple coneflower), radix corresponding to approx. 20 mg dried root from purple coneflower; Echinacea pallida Nutt. (pale coneflower), radix corresponding to approx. 20 mg dried root from pale coneflower and Thuja occidentalis L. (thuja), herba corresponding to approx. 11 mg dried herba from thuja.

The other ingredients are:
Mannitol (Ph. Eur.), betadex, lemon flavouring, glycerol dibehenate (Ph. Eur.), magnesium stearate (Ph. Eur.), citric acid monohydrate, saccharin sodium.

What Esberitox® COMPACT looks like and contents of the pack:

Round, beige-coloured tablets

Esberitox® COMPACT is available in packs with 20, 40 and 60 tablets.

Marketing authorisation holder

Quanstar Biotech Sdn Bhd, B-10-2,
Northpoint Office
Mid Valley City, No. 1, Medan Syed Putra Utara,
59200 Kuala Lumpur
Malaysia

Manufacturer

Schaper & Brummer GmbH & Co. KG
Bahnhofstr. 35
38259 Salzgitter
Lower Saxony, Germany
Tel. +49 53 41 3 07-0
Fax +49 53 41 3 07-124
info@schaper-bruemmer.de

This product information leaflet was last revised in January 2018