



Cimalgex 8 mg chewable tablets for dogs
Cimalgex 30 mg chewable tablets for dogs
Cimalgex 80 mg chewable tablets for dogs
Cimicoxib

STATEMENT OF THE ACTIVE SUBSTANCE AND OTHER INGREDIENTS

Each tablet contains:
Cimicoxib 8 mg
Cimicoxib 30 mg
Cimicoxib 80 mg
Cimalgex 8 mg tablets: oblong tablets (10x4mm), white to beige, brown spotted, with one break-line on both sides.
Cimalgex 30 mg tablets: Oblong tablets (15x6.5mm), white to beige, brown spotted, with 2 parallel breaking lines on both sides.
Cimalgex 80 mg tablets: Oblong tablets (21.5x8.5mm), white to beige, brown spotted, with 3 parallel breaking lines on both sides.

PHARMACODYNAMICS
Cimicoxib is a non-narcotic, non-steroidal anti-inflammatory drug (NSAID). It selectively inhibits the cyclooxygenase 2 enzyme (COX-2), which is responsible for pain, inflammation or fever. The cyclooxygenase 1 enzyme (COX-1) which has protective functions, for example, in the digestive tract and kidneys, is not inhibited by cimicoxib.

PHARMACOKINETICS
After oral administration in dogs at the recommended doses, cimicoxib is rapidly absorbed. Metabolism of cimicoxib is extensive. The major metabolite, demethylated cimicoxib is mainly eliminated in faeces by the biliary route and, to a lesser extent, in urine. The other metabolite, glucuronide conjugate of the demethylated cimicoxib, is eliminated in urine.

In an artificially induced pain model in dogs it was shown that the pain and inflammation reducing effects of cimicoxib lasted for approximately 10-14 hours.

DISPOSAL OF CONTAINERS
Medicines should not be disposed of via wastewater or household waste.

Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

DIRECTION FOR USE
For the treatment of pain and inflammation associated with osteoarthritis, and the management of peri-operative pain due to orthopaedic or soft tissue surgery, in dogs.

DOSAGE AND ADMINISTRATION
Oral use.
The recommended dose of cimicoxib is 2 mg per kg bodyweight, once daily.

The following table is presented as an example of how the tablets and tablet parts could be used in order to reach the recommended dose.

Bodyweight / kg	8 mg	30 mg	80 mg
2	1/2		
3	1		
4	1		
5		1/3	
6	1+1/2		
7-8	2		
9-11	2+1/2		
12	3		
13-17		1	
18-22			1/2
23-28		1+2/3	
29-33		2	
34-38		2+1/3	
39-43			1
45-48		3	
49-54			1+1/4
55-68			1+1/2

The choice of the most suitable tablet type or tablet parts is left to the discretion of the veterinarian based on the circumstances in each case, without leading to important over- or underdosing.

Treatment duration:
• Management of peri-operative pain due to orthopaedic or soft tissue surgeries: one dose 2 hours prior to surgery, followed by 3 to 7 days of treatment, based on the judgment of your veterinary surgeon.
• Relief of pain and inflammation associated with osteoarthritis: 6 months. For longer-term treatment, regular monitoring should be undertaken by your veterinary surgeon.

Cimalgex tablets can be given to dogs with or without food. The tablets are flavoured and studies (in healthy Beagle dogs) show they are likely to be taken voluntarily by most dogs.

CONTRAINDICATIONS
Do not use in dogs less than 10 weeks of age.
Do not use in dogs suffering from stomach or digestive system disorders or in dogs with bleeding problems.
Do not use at the same time as corticosteroids or other non-steroidal anti-inflammatory drugs (NSAIDs).
Do not use if the dog is hypersensitive to cimicoxib or any of the other ingredients in the product.
Do not use in breeding, pregnant or lactating animals (See "Special Warnings")

WARNINGS AND PRECAUTIONS
Special precautions for use in animals
The safety of this veterinary medicinal product has not been established in young dogs, so careful monitoring by your veterinary surgeon is recommended if the dog is less than 6 months of age.
Use in animals suffering from impaired cardiac, renal or hepatic function, may involve additional risk.
If such use cannot be avoided, these animals require careful veterinary monitoring. Avoid using this product in any animals which are dehydrated, hypovolaemic or hypotensive, as it may increase the risk of renal toxicity.
Use this veterinary medicinal product under strict veterinary monitoring in dogs with a risk of stomach ulcers or if the animal previously displayed intolerance to other NSAIDs.

Special precautions for people
Cimicoxib may cause skin sensitisation. Wash hands after use of the veterinary medicinal product.
In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.
People with a known hypersensitivity to cimicoxib should avoid contact with the veterinary medicinal product.

Statement on usage during pregnancy and lactation
Do not use in breeding, pregnant or lactating bitches. Although no data are available in dogs, studies with laboratory animals have shown effects on their fertility and foetal development.

Interactions with other medicaments
Cimalgex should not be administered in conjunction with corticosteroids or other NSAIDs. Pre-treatment with other anti-inflammatory substances may result in additional or increased adverse effects and accordingly a treatment-free period with such drugs should be observed before starting treatment with Cimalgex. The treatment-free period should take into account the pharmacokinetic properties of the veterinary medicinal products used previously.

ADVERSE EFFECTS/UNDESIRABLE EFFECTS
Mild gastro-intestinal disorders (vomiting and/or diarrhoea) were very commonly reported but these only lasted for a short time.
On rare occasions, serious gastrointestinal problems such as bleeding and ulcer formation have occurred. Other adverse reactions including loss of appetite or lethargy may also be observed on rare occasions.
In very rare cases, increases in kidney function results (renal biochemistry parameters) were noted. Furthermore, in very rare cases, renal failure has been reported. As for any long term NSAID treatment, kidney function should be monitored.

If any observed adverse effect persists after stopping treatment, the advice of your veterinary surgeon should be sought.
If adverse reactions such as persistent vomiting, repeated diarrhoea, blood in the stools, sudden weight loss, loss of appetite, lethargy or worsening of liver or kidney function results occur, use of the product should be discontinued and the advice of your veterinary surgeon should be sought immediately.

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

The frequency of adverse reactions is defined using the following convention:
- very common (more than 1 in 10 animals displaying adverse reactions during the course of one treatment)
- common (more than 1 but less than 10 animals in 100 animals)
- uncommon (more than 1 but less than 10 animals in 1,000 animals)
- rare (more than 1 but less than 10 animals in 10,000 animals)
- very rare (less than 1 animal in 10,000 animals, including isolated reports).

OVERDOSE AND TREATMENT
In an overdose study where 3 times (5.8 to 11.8 mg/kg body weight) and 5 times (9.7 to 19.5 mg/kg body weight) the recommended dose was administered to dogs for a period of 6 months, a dose related increase in gastrointestinal disturbances, which affected all dogs in the highest dose group, was noted. Similar dose related changes to haematology and white blood cell counts, as well as renal integrity, were also noted.
As with any NSAID, overdose may cause gastrointestinal, kidney, or liver toxicity in sensitive or compromised dogs.
There is no specific antidote to this product. Symptomatic, supportive therapy is recommended consisting of administration of gastrointestinal protective agents and infusion of isotonic saline.

WITHDRAWAL PERIOD: Not applicable.

STORAGE CONDITION
Keep out of reach of children.
Jauhi daripada kanak-kanak.
For animal use only.
Controlled medicines/Ubat terkawal.
Do not store above 25°C.
Blister packs - Any remaining divided tablets should be stored in the blisters below 25°C but discarded if not used within 24 hours.
Do not use this veterinary medicinal product after the expiry date which is stated on the label.

PACK SIZES
All strengths of Cimalgex tablets are available in the following pack sizes and types:
• Aluminium blisters (each strip containing 8 tablets) packaged into an outer cardboard box. Pack sizes of 32 or 144 tablets.

DATE OF REVISION: May 2018.

NAME AND ADDRESS OF THE MANUFACTURER
Vétoquinol SA
Magny Vernois
F-70200 Lure, France

NAME AND ADDRESS OF THE REGISTRATION HOLDER
GLADRON CHEMICALS SDN. BHD.
No. 7, Jalan TP 7, UEP Industrial Park,
40400 Shah Alam, Selangor D.E., Malaysia

REG. NUMBER MALXXXXXXXXXX

VETOQUINOL	Nom du produit : Cimalgex MY	
Créé par: P.GRANDVOINNET/P.JARDEL/A.STIFANI	Référence article : 443246 0517 N L	
le :	Dimensions : 210x500	
Visa :	1 couleur	
281 U		