

Tebonin® Film-Coated Tablet 40mg

EGb 761® (Each 266mg tablet contains Standardised Ginkgo biloba Extract 40mg equivalent to dry leaf 2,000mg)

Description & Composition

Tebonin® Film-Coated Tablet 40mg is presented as yellow and round film-coated tablets in blister strip packing.

One film-coated tablet contains:

Active component:

EGb 761®, a dry extract from Ginkgo biloba leaves (35 – 67:1) 40mg. EGb 761® is quantified to 24% flavone glycosides, 6% terpene lactones & less than 5ppm ginkgolic acids as required by the European Pharmacopoeia (Ph. Eur.)

Indications

Traditionally used to improve blood circulation and for general well-being.

General Warnings

As the use of Ginkgo may increase the tendency of bleeding, please consult your physician/pharmacist if you are on or intend to start any other medicines, and before you undergo any surgery/dental procedure.

(Memandangkan Ginkgo boleh meningkatkan kemungkinan pendarahan, sila rujuk kepada doctor/ahli farmasi sekiranya anda sedang atau sedang akan menggunakan ubat lain dan sebelum prosedur pembedahan/dental dijalankan)

Should you suffer from epileptic seizures, please consult with your doctor before starting treatment with Tebonin® Film-Coated Tablet 40mg. It cannot be excluded that the application of Ginkgo preparations promotes the occurrence of seizures in epileptic patients.

Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take Tebonin® Film-Coated Tablet 40mg.

This is a Traditional Medicine. Please consult a pharmacist/doctor before taking this product.

Contraindications

Tebonin® Film-Coated Tablet 40mg should not be used by patients with known hypersensitivity to Ginkgo extracts or to any of the excipients.

Children and adolescents

Do not give this medicine to children under 18 years because there is no relevant indication for this group.

Pregnancy, breast-feeding and fertility

Contraindicated in pregnant women. Insufficient reliable data in breastfeed-
ing women. It is unknown whether the extract components are excreted in
human milk. Animal studies do not reveal harmful effects of Tebonin® Film-Coated
Tablet 40mg on fertility.

Adverse effects

There are no verified data on the frequency of the undesirable effects observed
during treatment with Ginkgo biloba-containing preparations, since these un-
desirable effects have become known through single reports from patients,
physicians or pharmacists. According to these reports, the following side effects
may occur during treatment with Tebonin® Film-Coated Tablet 40mg: headache,
bleeding from single organs, mild gastrointestinal disturbances, allergic skin
reactions (reddening, swelling, itching).

Drug Interactions

No clinically relevant interactions have become known up to now.

Dosage and Administration

Adult: Take 1 tablet of Tebonin® Film-Coated Tablet 40mg each time, 3 times
daily. The intake of Tebonin® Film-Coated Tablet 40mg is independent of meals.

Expiry: 3 years from the date of manufacture

Pack size: 30's

Keep in a dry place below 30°C, protect from light and moisture

Keep out of reach of children.

Jauhkan daripada capaian kanak-kanak.

Revision date: July 2022

Manufactured by



Dr. Willmar Schwabe

From Nature. For Health.

Dr. Willmar Schwabe GmbH & Co. KG

Willmar-Schwabe-Str. 4, D-76227 Karlsruhe, Germany

Product Registration Holder:

A. Menarini Singapore Pte. Ltd (994247-T)

B-18-2, Level 18, The Ascent Paradigm,
No. 1 Jalan SS7/26A, Kelana Jaya,
47301 Petaling Jaya, Selangor

Product Registration.: MAL18076053T