

FOLDING BOX SANBEFLAM
(30 MEI '23) - Exp. MALAYSIA

Controlled
Medicine
SANBEFLAM®
Ketoprofen Gel 2.5% w/w

Active Ingredient
Each gram contains:
Ketoprofen 25 mg
Route of Administration: Topical

Product Description
Colorless gel, characteristic odor (mixed fragrance of Lavender oil and Neroli oil)

Pharmacodynamics
Pharmacotherapeutic category: non-steroid anti-inflammatory drug for topical use.
ATC Code: MO2AA10

Mechanism of Action
Ketoprofen is an inhibitor of both the cyclo-oxygenase and lipoxygenase pathways. Inhibition of prostaglandin synthesis provides for potent anti-inflammatory, analgesic and antipyretic effects. Lipoxygenase inhibitors appear to attenuate cell-mediated inflammation and thus retard the progression of tissue destruction in inflamed joints. In addition, Ketoprofen is a powerful inhibitor of bradykinin (a chemical mediator of pain and inflammation), it stabilises lysosomal membranes against osmotic damage and prevents the release of lysosomal enzymes that mediate tissue destruction in inflammatory reactions.

Pharmacokinetics
Absorption
By cutaneous route, absorption is very low. The percutaneous application of 50-150 mg of Ketoprofen produces plasma levels of the active ingredient of 0,08 - 0,15 µg/mL approx. 5 - 8 hours after application.

SANBEFLAM® gel allows the site specific topical delivery of Ketoprofen with very low plasma concentrations of drug. Therapeutic levels in the affected tissues provide relief from pain and inflammation, yet will satisfactorily overcome the problem of significant systemic unwanted effects.

Distribution
After oral administration of a single dose, maximum blood concentrations are achieved within 2 hours. Ketoprofen plasma half-life ranges from 1 to 3 hours. Plasma protein binding is 60% - 90%.

Elimination
Elimination is mainly by urinary route and in glucuronated form; approximately 90% of the amount administered is excreted within 24 hours.

Indication
Local treatment of painful disorders of the osteo-articular and muscular system of rheumatic or traumatic origin: contusions, distortions, muscle strains, stiff neck, lumbago.

Recommended Dose
Apply a thin layer of gel 1 to 3 times daily onto the area of affected skin, gently massaging to help absorption.

Contraindication
- Hypersensitivity to the active substance or to any of the excipients of the product.
- History of any photosensitivity reaction.
- Known hypersensitivity reactions, such as symptoms of asthma, allergic rhinitis or urticaria to fenofibrate, tiaprofenic acid, acetylsalicylic acid, or to other NSAIDs.
- History of skin allergy to ketoprofen, tiaprofenic acid, fenofibrate or UV blocker or perfumes.
- Sun exposure, even in case of hazy sun, including UV light from solarium, during the treatment and 2 weeks after its discontinuation.
- Ketoprofen gel should not be applied to open or infected wounds or lesions of the skin, such as occurs, for example, with eczema, or acne, or near the eyes.
- Third trimester of pregnancy.

Warning and Precautions
- The gel should be used with caution in patients with reduced heart, liver or renal function: isolated cases of systemic adverse reactions affecting renal function have been reported.
- The topical use of large amounts of product may give rise to systemic effects such as hypersensitivity and asthma.
- The treatment should be interrupted if rash appears.
- The recommended length of treatment should not be exceeded due to the risk of developing contact dermatitis and photosensitivity reactions increases over time.
- Hands should be washed thoroughly after each application of the product.
- Treatment should be discontinued immediately upon development of any skin reaction including cutaneous reactions after co-application of octocrylene containing products.
- It is recommended to avoid exposure of treated skin to direct sunlight including solarium (sunbeds), and to protect treated areas by wearing clothing during treatment with the product and for two weeks following its discontinuation to avoid the risk of photosensitisation.
- Do not use with occlusive dressings.
- The gel must not come into contact with mucous membranes.

- Patients with asthma combined with chronic rhinitis, chronic sinusitis, and/or nasal polyposis have a higher risk of allergy to aspirin and/or NSAIDs than the rest of the population.
- The use of topical products, especially if it is prolonged, may give rise to phenomena of sensitisation or local irritation.

Interactions with Other Medicaments
Interactions are unlikely as serum concentrations following topical administration are low. Serious interactions have been recorded after the use of high dose methotrexate with non-steroidal anti-inflammatory agents, including ketoprofen, when administered by the systemic route.

Pregnancy and Lactation
No embryopathic effects have been demonstrated in animals and there is epidemiological evidence of the safety of ketoprofen in human pregnancy. Nevertheless, it is recommended that ketoprofen should be avoided during pregnancy. Non-steroidal anti-inflammatory drugs may also delay labour. Trace amounts of ketoprofen are excreted in breast milk, therefore SANBEFLAM® Gel should not be used during breastfeeding.

Side Effects
The most common adverse reactions are photosensitive reactions (phototoxic and photosensitivity allergic reactions), the majority of which occurs after an incorrect use of the product (exposure of the skin to sunlight or solarium before 15 days from the last application, see contraindication and warning and precautions). There have been reports of localised skin reactions due to photosensitivity, including erythema, pruritus and burning sensations, which might spread beyond the area of application. Cases of more severe reactions such as bullous or phlyctenular eczema which may spread or become generalized have occurred rarely.
Other systemic effects of anti-inflammatory drugs: hypersensitivity, gastrointestinal and renal disorders (these depend on the transdermic spreading of the active ingredient, hence on the amount of gel applied, on the surface involved, on the degree of intactness of the skin, on the duration of the treatment and on the use of occlusive bandages).

Table with 5 columns: System Organ Class, Uncommon, Rare, Very Rare, Not known. Rows include Infections and Infestations, Blood and lymphatic system disorders, Immune system disorders, Eye disorders, Vascular disorders, Gastrointestinal disorders, Skin and subcutaneous tissue disorders, Renal and urinary disorders, General disorders and administration site conditions, Injury, poisoning and procedural complications.

Elderly patients are particularly susceptible to the adverse effects of non-steroidal anti-inflammatory drugs.

Symptoms and Treatment of Overdose
Overdose is unlikely by topical administration.
If accidentally ingested, the gel may cause systemic adverse effects depending on the amount ingested. However, if they occur, treatment should be supportive and symptomatic.

Storage Condition
Store below 30°C.

KEEP MEDICINES OUT OF REACH OF CHILDREN
JAUHKAN DARIPADA KANAK-KANAK

Presentations
Tube with net contents of 20 grams.
MAL, No.: MAL18081006AZ

Manufactured by: PT CAPRIFARMINDO LABS.
Jl. Industri Cimoreme No.8, Desa Cipeundeuy, Kecamatan Padalarang, Kabupaten Bandung Barat - Indonesia
For : PT SANBE FARMA
Jl, Taman Sari No,10
Bandung, Indonesia

Date of revision:
30.05.2023
* BF 197 - EMY2