

Front Label — 120x70mm

Ensure there is 5mm space
around the Pharma Code

Read Direction: →
Height: 12mm inc bleed

Pharma code:

Position: Centred 31.5mm
from right-hand edge of page

Elanco™

AH0470

Pulmotil™ AC

240 ml

Aqueous Concentrate 250 mg/ml

TILMICOSIN

Active Ingredient:

250 mg/ml of Tilmicosin (as the phosphate)

STORAGE CONDITIONS: Do not store above 30°C. Protect from direct sunlight. Store in a dry place and protect from humidity

Warning and Precautions: Refer inner label
FOR ORAL ANIMAL USE ONLY

KEEP OUT OF THE REACH OF CHILDREN

JAUHI UBAT DARI KANAK-KANAK

CONTROLLED MEDICINE

MAL18086143HA



Pull tab
for more
information



DOSAGE AND ADMINISTRATION:

Pigs: To be included in the drinking water to provide a daily dose of 15-20 mg/kg bodyweight for 5 days, which may be achieved by the inclusion of 200 mg tilmicosin per litre (80 ml Pulmotil AC per 100 litres).

Chickens and Turkeys (except hens producing eggs for human consumption): To be included in the drinking water at a daily dose of 15-20 mg/kg bodyweight in chickens and 10-27 mg/kg bodyweight in turkeys for 3 days, which may be achieved by the inclusion of 75 mg tilmicosin per litre (30 ml Pulmotil AC per 100 litres).

Calves: To be included in milk replacer only, at a dose of 12.5 mg/kg bodyweight and given twice daily for 3-5 consecutive days, which may be achieved by the inclusion of 1 ml of product every 20 kg bodyweight.

WITHDRAWAL PERIOD:

Pigs - 14 days; Chickens - 12 days; Turkeys - 19 days; Calves - 42 days
Not authorized for use in laying birds producing eggs for human consumption. Do not use within 14 days of onset of the laying.
Not for use in lactating animals.

Disposal of Containers: Refer inner label

Manufactured by: Elanco UK AH Limited
Fleming Road
Liverpool
L24 9LN
UK

For: Elanco Malaysia Sdn Bhd
Unit 5.04, Level 5 & 6,
Tower Block, The Bousteador,
No. 10, Jalan PJU 7/6,
Mutiara Damansara,
47800 Petaling Jaya,
Selangor, Malaysia

Pulmotil, Elanco and the diagonal bar logo are trademarks
of Elanco or its affiliates.

© 2025 Elanco or its affiliates

TAKE TIME



OBSERVE LABEL
DIRECTIONS

YL243464X

Inside Front Label — 120x70mm — Use only for text if necessary

Elanco™

AH0470

240 ml

Pulmotil™ AC *Aqueous Concentrate 250 mg/ml*

MAL18086143HA

TILMICOSIN

ACTIVE INGREDIENT:

250 mg/ml Tilmicosin (as the phosphate).

PRODUCT DESCRIPTION:

Concentrate for Oral Solution for use in drinking water or milk replacer. Clear yellow to amber coloured solution.

PHARMACODYNAMICS:

Tilmicosin is a semi-synthetic antibiotic of the macrolide group and is believed to affect protein synthesis. It has bacteriostatic action but at high concentrations it may be bactericidal.

This antibacterial activity is predominantly against Gram-positive microorganism with activity against certain gram-negative ones and *Mycoplasma* of a bovine, porcine, ovine and avian origin. In particular, its activity has been demonstrated against the following microorganism:

Pigs: *Mycoplasma hyopneumoniae*, *Pasteurella multocida* and *Actinobacillus pleuropneumoniae*.

Chickens and turkeys: *Mycoplasma gallisepticum* and *Mycoplasma synoviae*.

Calves: *Mannheimia haemolytica*, *Pasteurella multocida*, *Mycoplasma bovis* and *M. dispar*.

Scientific evidence suggests that macrolides act synergistically with the host immune system. Macrolides appear to enhance phagocyte killing of bacteria. Tilmicosin has been shown to inhibit *in vitro* the replication of the Porcine Reproductive and Respiratory Syndrome virus in alveolar macrophages in a dose dependent fashion. Cross-resistance between tilmicosin and other macrolides and lincomycin has been observed.

PHARMACOKINETICS:

Whilst blood concentrations of tilmicosin are low, there is pH-dependent macrophage accumulation of tilmicosin in inflamed tissues.

Pigs: After oral administration of 200 mg tilmicosin/l drinking water, the average active substance concentrations detected in lung tissue, alveolar macrophages and bronchial epithelium 5 days after the start of treatment were 1.44 µg/ml, 3.8 µg/ml and 7.4 µg/g respectively.

Chickens: As early as 6 hours after oral administration of 75 mg tilmosin/l drinking water, the average active substance concentrations detected in lung and alveolar tissue were 0.63 µg/g and 0.30 µg/g respectively. 48 hours after the start of treatment, the tilmosin concentrations in lung and alveolar tissue were 2.3 µg/g and 3.29 µg/g respectively.

Calves: As early as 6 hours after oral administration of 25 mg tilmosin/kg body weight/day in milk replacer, an average active substance concentration of 3.1 µg/g was detected in lung tissue. 78 hours after the start of treatment, the tilmosin concentration in lung tissue was 42.7 µg/g. Therapeutically effective concentrations of tilmosin were measured up to 60 hours after treatment.

Turkeys: After oral administration of 75 mg tilmosin/l drinking water, the average active substance concentrations detected in lung tissue, air sac tissue and plasma 5 days after the start of treatment were 1.89 µg/ml, 3.71 µg/ml and 0.02 µg/g respectively. The highest mean tilmosin concentration detected for lung tissues was 2.19 µg/g at 6 days; for air sac tissue it was 4.18 µg/g at 2 days and in the plasma it was 0.172 µg/g at 3 days.

TARGETED SPECIES:

Chickens (except hens producing eggs for human consumption), Turkeys, Pigs. Calves (non ruminant)

INDICATION:

Pigs: For the treatment and prevention of respiratory disease in pig herds, associated with *Mycoplasma hyopneumoniae*, *Pasteurella multocida*, *Actinobacillus pleuropneumoniae* and other organisms susceptible to tilmosin.

Chickens: For the treatment and prevention of respiratory disease in chicken flocks, associated with *Mycoplasma gallisepticum* and *M. synoviae*.

Turkeys: For the treatment and prevention of respiratory disease in turkey flocks, associated with *Mycoplasma gallisepticum* and *M. synoviae*.

Calves: For the treatment and prevention of bovine respiratory disease, associated with *Mannheimia haemolytica*, *Pasteurella multocida*, *Mycoplasma bovis*, *M. dispar* and other organisms susceptible to tilmosin.

DOSAGE AND ADMINISTRATION:

Pigs: To be included in the drinking water to provide a daily dose of 15-20 mg/kg bodyweight for 5 days, which may be achieved by the inclusion of 200 mg tilmosin per litre (80 ml Pulmotil AC per 100 litres).

Chickens and Turkeys (except hens producing eggs for human consumption): To be included in the drinking water at a daily dose of 15-20 mg/kg bodyweight in chickens and 10-27 mg/kg bodyweight in turkeys for 3 days, which may be achieved by the inclusion of 75 mg tilmosin per litre (30 ml Pulmotil AC per 100 litres).

Calves: To be included in milk replacer only, at a dose of 12.5 mg/kg bodyweight and given twice daily for 3-5 consecutive days, which may be achieved by the inclusion of 1 ml of product every 20 kg bodyweight.

One 240 ml bottle of Pulmotil AC is sufficient to medicate 300 litres of drinking water for pigs or 800 litres of drinking water for chickens or turkeys.

One 240 ml bottle of Pulmotil AC is sufficient to medicate in milk replacer 12 to 20 veal calves each of 40 kg bodyweight depending on the duration of treatment.

The uptake of medicated water/milk replacer depends on the clinical condition of the animals. In order to obtain the correct dosage, the concentration of Pulmotil AC should be adjusted accordingly.

CONTRAINDICATION:

Do not allow horses and other equines access to drinking water containing tilimicosin. Do not use in case of hypersensitivity to tilimicosin or to any of the excipients.

WARNINGS AND PRECAUTIONS:

Special warnings for each target species

Important: Must be diluted before administration to animals.

Pigs, chickens and turkeys: Water consumption should be monitored in order to guarantee adequate dosing. In case water consumption does not match quantities for which recommended concentrations were calculated, the concentration of Pulmotil AC has to be adapted in a way that the recommended dosage will be taken up by the animals or different medication should be considered.

Special precautions for use

i. Special precautions for use in animals

For oral use only. Contains disodium edetate; do not inject.

Severely ill individuals tend to drink less and may need simultaneous treatment, preferably by parenteral medication.

Inappropriate use of the product may increase the prevalence of bacteria resistant to tilimicosin and may decrease the effectiveness of treatment with tilimicosin-related substances. The use of the product should be based on susceptibility tests.

Medicated drinking water or milk replacer should be prepared fresh every 24 hours.

Official, national and regional antimicrobial policies should be taken into account when the product is used.

ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals

- Tilimicosin may induce irritation. Macrolides, such as tilimicosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tilimicosin may lead to cross reactions to other macrolides and *vice versa*.

Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided.

- To avoid exposure during preparation of the medicated drinking water, wear overalls, safety glasses, and impervious gloves.

- Do not eat, drink or smoke when handling this product. Wash hands after use.

- In the case of accidental ingestion, wash out mouth immediately with water and seek medical advice.

- In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water.

- Do not handle the product if you are allergic to ingredients in the product.

- If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning.

- Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

PREGNANCY AND LACTATION:

The safety of tilimicosin has not been established in animals used for breeding purposes.

INTERACTION WITH OTHER MEDICINAL PRODUCTS:

None known

SIDE EFFECTS:

In very rare cases, a decrease in water intake has been observed.

SYMPTOMS AND TREATMENT OF OVERDOSE:

When pigs are offered drinking water containing 300 or 400 mg/litre (equivalent to 22.5-40 mg/kg bodyweight or 1.5-2 times the recommended concentration) commonly animals exhibit a reduced water intake. Although this has a self-limiting effect on tilmicodin intake, it could, in extreme circumstances, result in dehydration. This can be corrected by the removal of the medicated drinking water and replacement with fresh unmedicated water.

No symptoms of overdose have been seen in chickens given drinking water containing levels of tilmicodin up to 375 mg/litre (equivalent to 75-100 mg/kg bodyweight or 5 times the recommended dose) for 5 days; daily treatment with 75 mg/litre (equivalent to the maximum recommended dose) for 10 days resulted in a reduction in faecal consistency.

No symptoms of overdose have been seen in turkeys given drinking water containing levels of tilmicodin up to 375 mg/litre (equivalent to 50-135 mg/kg bodyweight or 5 times the recommended dose) for 3 days; daily treatment with 75 mg/litre (equivalent to the maximum recommended dose) for 6 days also produced no symptoms of overdose.

No symptoms of overdose, with exception of a slight decrease in the milk consumption, have been seen in calves given twice daily doses 5 times the maximum recommended dose or for twice the maximum recommended duration of treatment.

WITHDRAWAL PERIOD:

Pigs – 14 days; Chickens – 12 days; Turkeys – 19 days; Calves – 42 days

Not authorised for use in laying birds producing eggs for human consumption. Do not use within 14 days of onset of the laying. Not for use in lactating animals.

STORAGE CONDITIONS:

Do not store above 30°C. Protect from direct sunlight. Store in dry place and protect from humidity.

DISPOSAL OF CONTAINER:

No special package disposal required.

SHELF LIFE:

Shelf life as packaged for sale: 3 years.

After opening the bottle: 24 hours.

After reconstitution / dilution: 24 hours.

Manufactured by: Elanco UK AH Limited
Fleming Road
Liverpool
L24 9LN
UK

For: Elanco Malaysia Sdn Bhd
Unit 5.04, Level 5 & 6,
Tower Block, The Bousteador,
No. 10, Jalan PJU 7/6,
Mutiar Damansara,
47800 Petaling Jaya,
Selangor, Malaysia

REVISED: 11 September 2025

Elanco™

AH0470

Pulmotil™ AC

240 ml

Aqueous Concentrate 250 mg/ml

TILMICOSIN

Active Ingredient:

250 mg/ml of Tilmicosin (as the phosphate)

STORAGE CONDITIONS: Do not store above 30°C. Protect from direct sunlight. Store in a dry place and protect from humidity

Warning and Precautions: Refer inner label

FOR ORAL ANIMAL USE ONLY

KEEP OUT OF THE REACH OF CHILDREN

JAUHI UBAT DARI KANAK-KANAK

CONTROLLED MEDICINE

MAL18086143HA

DOSAGE AND ADMINISTRATION:

Pigs: To be included in the drinking water to provide a daily dose of 15-20 mg/kg bodyweight for 5 days, which may be achieved by the inclusion of 200 mg tilmicosin per litre (80 ml Pulmotil AC per 100 litres).

Chickens and Turkeys (except hens producing eggs for human consumption): To be included in the drinking water at a daily dose of 15-20 mg/kg bodyweight in chickens and 10-27 mg/kg bodyweight in turkeys for 3 days, which may be achieved by the inclusion of 75 mg tilmicosin per litre (30 ml Pulmotil AC per 100 litres).

Calves: To be included in milk replacer only, at a dose of 12.5 mg/kg bodyweight and given twice daily for 3-5 consecutive days, which may be achieved by the inclusion of 1 ml of product every 20 kg bodyweight.

WITHDRAWAL PERIOD:

Pigs - 14 days; Chickens - 12 days; Turkeys - 19 days; Calves - 42 days
Not authorized for use in laying birds producing eggs for human consumption. Do not use within 14 days of onset of the laying.
Not for use in lactating animals.

Disposal of Containers: Refer inner label

Manufactured by: Elanco UK AH Limited
Fleming Road
Liverpool
L24 9LN
UK

For: Elanco Malaysia Sdn Bhd
Unit 5.04, Level 5 & 6,
Tower Block, The Bousteador,
No. 10, Jalan PJU 7/6,
Mutiar Damansara,
47800 Petaling Jaya,
Selangor, Malaysia

Pulmotil, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

© 2025 Elanco or its affiliates

YL243464X



35mm

29mm

Overprint Page — 175x70mm

Batch Number:

Mfg Date:

Expiry Date:

Legend — 175x70mm

LEGEND KEY

Template: YL_120x70mm_VD35mm_v4a

Text Free Area

Print Free Area

Variable Data
and Code area

50 mm Rule Bar

Elanco Template Key v2

PRODUCT INFO

BLUE #:	243464AP	Product Name:	Pulmotil AC
Item Code:	YL243464X	Component:	Label
Product Code:	AH0470	Pack Size:	240ml
Previous Item Code	YL213320A		

ARTWORK INFO

Template:	YL_120x70mm_VD35mm_v4a	Packaging Spec(s):	Base Spec: GPG365 Primary Spec: DPG650	
Barcodes/Type:	Pharmacode: 652	Add. Info:	N/A	
		Minimum Core Data Point Size:	4.5pt	
GTIN:	GTIN Not Required	Proof #	P1a	By/Date RA 08-Oct-2025
	BLACK PMS 2395 C NON PRINTING			

ELANCO ARTWORK LEGEND v16