

COXIDIN® 200 g/kg MicroGranulate Premix for medicated feeding stuff

STATEMENT OF THE ACTIVE SUBSTANCE:

Monensin sodium 200 g/kg

PRODUCT DESCRIPTION:

Coxidin® 200 g/kg MicroGranulate represents granules of beige to brown colour intended for administration through feed of poultry.

PHARMACODYNAMICS:

Monensin sodium acts as an ionophore, i.e. a chemical substance that complexes monovalent cations with the lipophilic structure that facilitates the transport of the bound ion through biological membranes. Coccidial sporozoites exposed to monensin sodium in the intestinal lumen exhibit considerable swelling, large vacuoles, pitting and holes in the surface suggesting extreme osmotic damage which is potentially lethal. Development of a sporozoite that successfully invades a host cell is inhibited as the ionophore continues its destructive process. Monensin sodium selectively destroys intracellular sporozoites while remaining relatively non-injurious to the host cell. It has been shown that monensin sodium has an effect on second generation merozoites but not upon developing gametocytes.

PHARMACOKINETICS:

In the chicken monensin sodium is absorbed at a limited extent. It is metabolized extensively and gives rise to demethylated, oxidized and decarboxylated metabolites, found in the excreta and tissues where they represent each less than 10% of the total monensin derivatives. The proportion of unchanged monensin in the excreta measured during a relatively short initial excretion period (8 hrs) is 95% and 63% for the chicken and turkey, respectively. The studies on monensin residues in chicken and turkey show that the highest contents are found in the skin plus fat then the liver for both species and that their remanence is very limited in all tissues (less than 24 hours). Monensin sodium is excreted rapidly mainly through the faeces with considerable biliary excretion.

INDICATIONS FOR USE:

As an aid in the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mivati*, *E. necatrix* and *E. tenella* in poultry (broilers and replacement layers).

RECOMMENDED DOSAGE:

Monensin 100-120 mg/kg (ppm) in ration (equivalent to 0.5 – 0.6 kg Coxidin® 200/t of final feed).

MODE OF ADMINISTRATION:

For oral administration through the feed.

CONTRAINDICATIONS:

Dogs, horses or other equines must not be allowed access to feeds containing COXIDIN® 200. Ingestion by these species may be fatal.

WARNINGS AND PRECAUTIONS:

Recommended levels of COXIDIN® 200 must not be exceeded. High levels can cause reduced feed intake, poor growth, diarrhea and death. Responses to COXIDIN® 200 are unlikely if the nutrient intake is significantly restricted.

Poultry consuming monensin should not be treated with products containing tiamulin. Severe growth depression may occur.

INTERACTIONS WITH OTHER

MEDICAMENTS:

This coccidiostat should not be mixed or used simultaneously with any other medicinal product having a similar effect.

STATEMENT ON USAGE DURING

PREGNANCY AND LACTATION:

In the absence of toxicology study, this veterinary medicinal product must not be used during pregnancy and lactation.

ADVERSE EFFECTS/UNDESIRABLE

EFFECTS:

If suspected adverse reactions occur, treatment should be discounted and to inform veterinarian/local authority.

OVERDOSE AND TREATMENT:

Excessively high levels due to mixing errors can cause reduced feed intake, poor growth, diarrhoea and death.

WITHDRAWAL PERIODS:

Poultry (meat): 0 day.

Poultry (eggs): Do not use in birds producing eggs for human consumption.

STORAGE CONDITIONS:

Keep in cool and dry place away from direct sunlight. Store below 30°C.

Keep out of reach of children.

PACKAGING AVAILABLE:

Polyethylene-lined paper bags of 20 kg.

MANUFACTURER:

BIOVET Joint Stock Company
39 Petar Rakov Street, 4550 Peshtera, Bulgaria

MARKETING AUTHORIZATION HOLDER:

Pahang Pharmacy Sdn. Bhd. (10826-K)
Lot 5979 Jalan Teratai, 5½ Mile Off Jalan Meru,
41050 Klang, Selangor Darul Ehsan, Malaysia

DATE OF REVISION:

February 2018.