

For Veterinary Use Only*Broad Spectrum Anticoccidials***Tolcoxin 5% w/v suspension****CONTENT**

Each Liter contains 50g of Toltrazuril
Sodium Benzoate(Preservative)- - - - - 2.1g/1L

PRODUCT DESCRIPTION

Colorless or pale yellow suspension, liquid.

PHARMACODYNAMICS

Toltrazuril is a triazinon derivative. It acts against coccidia of the genus *Isospora* and *Eimeria*. It is acting against all intracellular development stages of coccidia of the merogony (asexual multiplication) and gamogony (sexual phase). All stages are destroyed, thus the mode of action is coccidiocidal.

PHARMACOKINETICS**Piglet**

After oral administration toltrazuril is slowly absorbed. The main metabolite is characterised as toltrazuril sulfone. The elimination of toltrazuril is slow with a half-life elimination time around 3 days. The major route of excretion is via the faeces.

Calves

After oral administration in cattle toltrazuril is slowly absorbed. The elimination of toltrazuril is slow with a terminal half-life time of approximately 2.5 days. The main metabolite is characterised as toltrazuril sulfone. The major route of excretion is via the faeces.

Lambs

After oral administration toltrazuril is slowly absorbed in mammals. The main metabolite is characterised as toltrazuril sulfone. The elimination of toltrazuril is slow with an elimination half-life time of approximately 9 days. The major route of excretion is via the faeces.

INDICATION**Piglets**

For the prevention of clinical signs of coccidiosis in neonatal piglets on farms with a confirmed history of coccidiosis caused by *Isospora suis*

Calves

For the prevention of clinical signs of coccidiosis and reduction of coccidia shedding in housed calves replacing cows producing milk for human consumption (dairy cows) on farms with a confirmed history of coccidiosis caused by *Eimeria bovis* or *Eimeria zuernii*

Lambs

For the prevention of clinical signs of coccidiosis and reduction of coccidia shedding in lambs on farms with a confirmed history of coccidiosis caused by *Eimeria crandallis* and *Eimeria ovinoidalis*

DOSAGE & ADMINISTRATION**All Species**

The ready-to-use oral suspension must be shaken before use.

Piglets : Individual animal treatment. Each pig to be treated on day 3–5 of life with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 mL oral suspension per kg body weight. Due to the small volumes required to treat individual piglets, use of dosing equipment with a dose accuracy of 0.1 mL is recommended.

Treatment during an outbreak will be of limited value for the individual piglet because of damage to the small intestine having already occurred.

Calves : Each animal should be treated with a single oral dose of 15 mg toltrazuril/kg body weight corresponding to 3.0 mL oral suspension per 10 kg body weight.

Lambs : Each animal should be treated with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 mL oral suspension per kg body weight.

CONTRAINDICATIONS

For environmental reasons :

Do not use in calves weighing more than 80 kg bodyweight.

Do not use in fattening units such as veal or beef calves.

WARNINGS / PRECAUTIONS**Special warnings**

- As with any antiparasiticide frequent and repeated use of antiprotozoals from the same class may lead to the development of resistance.
- It is recommended to treat all animals in a pen
- Hygienic measures may reduce the risk of coccidiosis. It is therefore, recommended to improve concomitantly the hygienic conditions in the concerned facility, particularly dryness and cleanliness.
- To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.
- To alter the course of an established clinical coccidial infection, in individual animals already showing signs of diarrhoea, additional supportive therapy may be required.

Special precautions for use

- Special precautions for use in animals
- Special precautions to be taken by the person administering the veterinary medicinal product to animals. Wash any splashes from skin or eyes immediately with water.
- Other Precautions

The major metabolite of toltrazuril, toltrazuril sulfone (ponazuril), has been shown to be both persistent (half-life >1 year) and mobile in soil and to be toxic to plants.

For environmental reasons :

- In order to prevent any adverse effects on plants and possible contamination of groundwater manure from treated calves must not be spread onto land without dilution with manure from untreated cows.
Manure from treated calves must be diluted with at least 3 times the weight of manure from mature cows before it can be spread onto land.
- Lambs kept throughout the whole life span indoors under an intensive rearing system must not be treated beyond the age of 6 weeks or body weight of more than 20 kg at treatment.
Manure from these animals should only be applied to the same piece of land every third year.

DRUGS INTERACTIONS : None known.

PREGNANCY AND LACTATION

Laboratory studies in rat and rabbit have not produced any evidence of a teratogenic, foetotoxic or maternotoxic effect.

ADVERSE EFFECTS : None known.

OVERDOSE AND TREATMENT

A threefold overdose is well tolerated by healthy piglets and calves without signs of intolerance.

No signs of overdose have been observed in lamb safety studies with threefold overdose at a single treatment and twofold overdose treatment on 2 consecutive days.

STORAGE CONDITIONS : Store below 30°C. Protected from moisture and light.

WITHDRAWAL PERIOD(S) :

Piglets : Meat and offal - 77 days

Calves : Meat and offal - 63 days

Not permitted for use in lactating animals producing milk for human consumption.

Lambs : Meat and offal - 42 days

Not permitted for use in lactating sheep producing milk for human consumption.

DOSAGE FORMS AND PACKAGING AVAILABLE

100mL suspension packed into HDPE bottle.

DATE OF REVISION OF PACKAGE INSERT : 27 JUL 2018



Manufactured by
KBNP, INC.
235-9 Chusaro, Sinam, Yesan,
Chungnam, Korea



Distributed by
Nutri Pharmax Sdn.Bhd.
6, Lorong Serunai 1, Valdor Sungai Bakap,
14200 Sungai Jawi, Pulau Pinang, Malaysia
Tel: +604-582 9377