

PF TYLOSIN PREMIX 25% W/W POWDER

Composition:

Tylosin (as Tylosin Phosphate).....250mg

Exp. Qs.....1g

Description

White colour powder

Pharmacodynamic

Tylosin is a macrolide antibiotic produced by a strain of *Streptomyces fradiae*. It exerts its antimicrobial effect by inhibiting protein synthesis of susceptible micro-organisms.

The tylosin spectrum of activity includes Gram-positive bacteria, some Gram-negative strains such as *Pasteurella*, and *Mycoplasma* spp at concentrations of 16µg/ml or less.

Pharmacokinetic

Absorption: Tylosin reaches maximal blood levels between 1 and 3 hours after an oral dose. Minimal or no blood levels remain 24 hours after an oral dose.

Distribution: After oral doses were given to pigs, tylosin was found in all tissues, between 30 minutes and two hours after administration, except for the brain and spinal cord.

Biotransformation and Elimination: It has been shown that most of the material which is excreted is to be found in the faeces and consists of tylosin (factor A), relomycin (factor D) and dihydrodesmycosin.

Indication

Pigs:

For the prevention and control of enzootic pneumonia.

For the treatment and control of *Lawsonia intracellularis*, the organism associated with Porcine Intestinal Adenomatosis (Ileitis) and Porcine Haemorrhagic Enteropathy.

Route of Administration

Oral

Recommended dose

For incorporation into dry feed at the registered mill.

A manufacturer who is approved to incorporate directly at any concentration, veterinary medicinal products or premixtures containing such products must be responsible for mixing when incorporation is less than 2 kg per tonne for final feed.

For oral administration.

Pigs:

Prevention and control of enzootic pneumonia:

Treatment and control of *Lawsonia intracellularis*:

3-6 mg tylosin activity/kg bodyweight, which may normally be achieved by adding the product at the rate of 400 g per tonne, giving 100 g tylosin base per tonne. Feed as the only ration for 21 days.

The required levels of tylosin are obtained by mixing the appropriate quantity of PF Tylosin premix with 20-50 kg of a suitable feed component, prior to incorporation into the bulk of the feed to be prepared.

For the preparation of the medicated feed the body weight of the animals to be treated and their actual daily feed consumption should be taken into due account. Consumption may vary depending on factors like age, breed, husbandry system. To provide the required amount of active substance in mg per kg mixed feed the following calculation should be made:

...mg PF Tylosin premix/kg bodyweight/day	x	Average body weight (kg) of the animals to be treated	=	...mg PF Tylosin permix per kg/ mixed feed
Average daily amount of mixed feed intake /kg per animal				

The intake of medicated feed depends on the clinical condition of the animals. In order to obtain the correct dosage, the concentration of PF Tylosin Premix has to be adjusted accordingly.

Contraindication

Do not use in known cases of hypersensitivity.

Do not use in animals vaccinated with tylosin-sensitive vaccines either at the same time or within one week previously.

Interaction with other medicinal products

Lincosamides and aminoglycoside antibiotics antagonise the activity of tylosin.

Pregnancy and lactation

No studies have been conducted in the target species population. Use only according to the benefit risk assessment by the responsible veterinarian.

Side effects

None known.

Symptoms and treatment of overdose

PF Tylosin Premix has been shown to produce no adverse effects when fed to pigs at 600 ppm in the feed (six times the recommended maximum level) for 28 days.

Storage

Store in a cool dry place (<30°C). Strictly avoid light and heat exposure.

Warning and Precaution

Keep out of reach of children / Jauhi ubat dari kanak-kanak

Special warnings for each target species

None.

Special precautions for use in animals

For incorporation into medicated feeding stuffs only.

To ensure thorough dispersion of the product it should first be mixed with a small quantity of feed ingredients before incorporation into the final mix.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Tylosin may induce irritation. Macrolides, such as tylosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tylosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided.

To avoid exposure during preparation of the medicated feed, wear overalls, safety glasses, impervious gloves and wear either a disposable half mask respirator conforming to European Standard EN149 or a non-disposable respirator to European Standard EN140 with a filter to EN143. Wash hands after use.

In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water.

Do not handle the product if you are allergic to ingredients in the product.

If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

Only to be incorporated by a manufacturer who is approved to mix at a rate of below 2 kg per tonne final feed.

Shelf life

36 months

Shelf life after opening

24 hours

Shelf life after dilution or constitution: 24 hours

Packaging: 25kg

Withdrawal period

Meat: Pigs: Zero days.

Revised date

27/11/2017

Distributed by:

PANFAST MARKETING (M) SDN.BHD.

PENANG SCIENCE PARK 1102, MUKIM 13, LORONG PERINDUSTRIAN BUKIT

MINYAK 18, KAWASAN PERINDUSTRIAN BUKIT MINYAK,

14100 SIMPANG AMPAT, PENANG, MALAYSIA

TEL: +604-5016868 FAX: +604-5016866

Manufactured by:

SHENNONG ANIMAL HEALTH (M) SDN.BHD.

No.6, 8, 10, Jalan Industri USJ 1/19, Taman Perindustrian USJ 1

47610 SUBANG JAYA, Selangor Darul Ehsan, Malaysia.

Tel: 03-8011 4646 Fax : 03-8024 1166