

PF ROBENIDINE PREMIX 6.6% W/W POWDER

Composition:

Robenidine HCL.....66mg

Exp. Qs.....1g

Description

Light Yellowish Colour Powder

Pharmacodynamic

Robenidine has broad-spectrum activity against Eimeria species of birds.

Robenidine is most effective against late developing stages of first and second generation schizonts. It exhibits some activity against gamonts (sexual stages). Its action is first coccidiostatic and then coccidiocidal.

Pharmacokinetic

Robenidine hydrochloride is absorbed at limited extent and excreted rapidly. Unchanged robenidine is the major compound excreted by the chicken.

Indication

As an aid in the prevention of coccidiosis caused by Eimeria mitis, E. (Eimeria) brunetti, E. (Eimeria) tenella, E. (Eimeria) acervulina, E. (Eimeria) maxima, E. (Eimeria) necatrix and E. (Eimeria) praecox in broiler chickens.

Route of Administration

Oral

Recommended dose

33 mg/kg (milligrams per kilogram) (0.0033%) of robenidine hydrochloride in the complete feed. Feed this medicated feed as the sole ration.

Contraindication

Layer birds; hypersensitivity to robenidine.

Interaction with other medicinal products

No interactions or incompatibilities with feedingstuffs, carriers or other approved additives have been recorded and reported.

Pregnancy and lactation

Do not feed to laying hens, replacement chickens or breeding birds.

Side effects

It imparts unpleasant taste to the flesh of broiler birds if not withdrawn before slaughter.

Symptoms and treatment of overdose

Robenidine is safe for the chicken for fattening at doses corresponding to safety factors of about 6 with respect to the maximum dose proposed for use.

Storage

Store in a cool dry place (<30°C). Strictly avoid light and heat exposure.

Warning and Precaution

Keep out of reach of children / Jauhi ubat dari kanak-kanak

1. Do not use this medicated feed for treatment of outbreaks of Coccidiosis.

2. Consult a veterinarian or poultry pathologist if losses exceed 0.5% in a two day period.

3. Follow directions carefully.

Shelf life

36 months

Shelf life after opening

24 hours

Shelf life after dilution or constitution

24 hours

Withdrawal period

Treated chickens must not be slaughtered for use in food for at least 6 days after latest treatment with this medicated feed.

Packaging

25kg

Revised date

23/7/2018

Product Registration Holder:

PANFAST MARKETING (M) SDN.BHD.

PENANG SCIENCE PARK 1102, MUKIM 13, LORONG PERINDUSTRIAN BUKIT MINYAK 18, KAWASAN PERINDUSTRIAN BUKIT MINYAK, 14100 SIMPANG AMPAT, PENANG, MALAYSIA
TEL: +604-5016868 FAX: +604-5016866

Manufactured by:

SHENNONG ANIMAL HEALTH (M) SDN.BHD.

No.6, 8, 10, Jalan Industri USJ 1/19, Taman Perindustrian USJ 1 47610 SUBANG JAYA, Selangor Darul Ehsan, Malaysia.
Tel: 03-8011 4646 Fax : 03-8024 1166