

Veterinary Package Insert

TP-LINCO 400MG/G POWDER (40% Lincomycin Hydrochloride)

PRODUCT DESCRIPTION

White to almost white colour powder, which contains 400mg of lincomycin hydrochloride per gram.

PHARMACODYNAMICS & PHARMACOKINETICS

Lincomycin is a lincosamide antibiotic derived from *Streptomyces lincolensis* which inhibits protein synthesis. Lincomycin binds to the 50S sub-unit of the bacterial ribosome close to the peptidyl transfer centre and interferes with the peptide chain elongation process by causing premature peptidyl-tRNA dissociation from the ribosome.

Lincomycin is active against some Gram-positive bacteria (*Clostridium perfringens*) and mycoplasmas (*Mycoplasma hyopneumoniae*).

While the lincosamides are generally considered to be bacteriostatic agents, the activity depends on the sensitivity of the organism and concentration of the antibiotic. Lincomycin may be either bactericidal or bacteriostatic.

In pigs, lincomycin is rapidly absorbed following oral administration. The oral absorption of lincomycin was found to be 53%±19%.

Crossing the intestinal barrier, lincomycin is widely distributed to all tissues, especially the lungs and joint cavities; the volume of distribution is about 1 litre. The elimination half-life of lincomycin is greater than 3 hours. Approximately 50% of lincomycin is metabolised in the liver. Lincomycin undergoes enterohepatic circulation. Lincomycin is eliminated unchanged or in the form of various metabolites in bile and urine. High concentrations of the active form are observed in the intestine.

For chickens, metabolites comprised more than 75% of total residues in the liver. Unmetabolised lincomycin declined at a slightly faster half-life ($t_{1/2}$ = 5.8 hours) than total residue. Lincomycin and one unknown metabolite comprised >50% of the muscle residue at zero hours. The excreta contained mostly unmetabolised lincomycin (60-85%) during treatment.

INDICATION

Pigs:

Treatment and metaphylaxis of enzootic pneumonia caused by *Mycoplasma hyopneumoniae*.

The presence of the disease in the group must be established before the product is used.

Chickens:

Treatment and metaphylaxis of necrotic enteritis caused by *Clostridium perfringens*.

The presence of the disease in the group must be established before the product is used.

TARGET SPECIES

Pigs and chickens.

RECOMMENDED DOSAGE AND ADMINISTRATION

For use in drinking water.

Dosing guidance and recommended doses:

To ensure a correct dosage, body weight should be determined as accurately as possible to avoid underdosing.

The intake of medicated water depends on the physiological and clinical condition of the animals.

In order to obtain the correct dosage, the concentration of the lincomycin has to be adjusted accordingly. The uptake of water should be monitored frequently.

The medicated water should be the only source of drinking water for the animals for the entire duration of the treatment period. After the end of the medication period, the water supply system should be cleaned appropriately to avoid intake of sub-therapeutic amounts of the active substance.

Dosage:

Pigs:

Enzootic pneumonia: 10 mg lincomycin per kg of body weight (corresponding to 27.25 mg product per kg bodyweight) for 21 consecutive days.

Chickens:

Necrotic enteritis: 5 mg lincomycin per kg of body weight (corresponding to 13.63 mg product per kg bodyweight) for 7 consecutive days.

The concentration to be used depends on the actual body weight and the water consumption of the animals and can be calculated according to the following formula:

Dosage (mg product) per kg body weight and day	x	Mean body weight (kg) of animals to be treated	= ...mg product per litre drinking water
--	---	---	--

Average daily water intake (litre/animal)

CONTRAINDICATIONS

Do not use in cases of known hypersensitivity to the active substance or to any of the excipients.

Do not use in cases of known resistance to lincosamides.

Do not administer, and do not allow access to water containing lincomycin to horses, ruminants, guinea pigs, hamsters, chinchillas and rabbits as this could result in severe gastro-intestinal disturbance.

Do not use in cases of hepatic dysfunction.

WARNINGS & PRECAUTIONS.

Medicated drinking water uptake can be affected by the severity of the disease. In case of insufficient uptake of water, pigs should be treated parenterally.

The susceptibility of *Mycoplasma hyopneumoniae* to antimicrobial agents is difficult to test in vitro owing to technical constraints. In addition, there is a lack of clinical breakpoints for both *M. hyopneumoniae* and *C. perfringens*. Where possible, therapy should be based on local (regional, farm level) epidemiological information concerning the response of enzootic pneumonia/necrotic enteritis to treatment with lincomycin.

Special precautions for use in animals

Use of the veterinary medicinal product preferably should be based on identification of the target pathogen and susceptibility testing of the bacteria isolated from the animal.

Use of the veterinary medicinal product deviating from the instructions given in the summary of product characteristics may increase the prevalence of bacteria resistant to the lincomycin and may decrease the effectiveness of treatment with other lincosamides, macrolides and streptogramin B due to the potential for cross-resistance.

Repeated or prolonged use should be avoided by improving the farm management and hygiene practices.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

This product contains lincomycin and lactose, either of which can cause allergic reactions in some people. People with known hypersensitivity to lincomycin or any other lincosamide, or to lactose, should avoid contact with the veterinary medicinal product.

Care should be taken not to raise and inhale any dust.

Contact with skin and eyes should be avoided.

Personal protective equipment consisting of approved dust masks (either a disposable half-mask respirator or a non-disposable respirator with a filter), gloves and safety glasses should be worn when handling and mixing the product. If respiratory symptoms develop following exposure, seek medical advice and show this warning to the physician.

In case of accidental exposure to the skin, eyes or mucous membranes, wash the affected area thoroughly with plenty of water. If symptoms such as skin rash or persistent eye irritation appear after exposure, seek medical advice immediately and show the package leaflet or label to the physician.

Do not smoke, eat or drink when handling the product.

Wash hands and any exposed skin with soap and water immediately after use.

INTERACTIONS WITH OTHER MEDICATIONS

Antagonism may exist between lincomycin and macrolides such as erythromycin and other bactericidal antibiotics; concurrent use is therefore not recommended due to competitive binding at the 50S ribosomal subunit of the bacterial cell.

The bioavailability of lincomycin may decrease in the presence of gastric antacids or activated charcoal, pectin or kaolin.

Lincomycin can potentiate neuromuscular effects of anaesthetics and muscle relaxants.

PREGNANCY AND LACTATION

Laboratory studies in rats have not produced any evidence of teratogenic effects, although foetotoxicity has been reported. The safety of the veterinary medicinal product has not been established during pregnancy, lactation or lay in the target species. Use only according to the benefit/risk assessment by the responsible veterinarian.

ADVERSE EFFECTS

On rare occasions, pigs given lincomycin-medicated water may develop diarrhea or soft stools and/or mild swelling of the anus within the first two days

after onset of treatment. On rare occasions some pigs may show reddening of the skin and mild irritable behaviour. These conditions are usually self-correcting within 5-8 days without discontinuing the lincomycin treatment. Allergic/hypersensitive reactions occur on rare occasions.

OVERDOSE & TREATMENT

A dosage greater than 10 mg lincomycin per kg of body weight may cause diarrhoea and loose stools in pigs.

In case of accidental overdose, the treatment must be stopped and restarted at the recommended dose level.

There is no specific antidote, treatment is symptomatic.

WITHDRAWAL PERIOD

Meat and offal of pigs: 1 day
Meat and offal of chickens: 5 days
Not authorised for use in laying birds producing eggs for human consumption.

MAXIMUM RESIDUAL LIMIT (MRL)

Substance/ Drug Definition of residues in which MRL was set	Food	Maximum Residue Limits (MRLs) in µg/kg
Lincomycin	Chicken, muscle	100
	Chicken, skin & fat	50
	Chicken, liver	500
	Chicken, kidney	1500
	Chicken, eggs	50
	Pig, muscle	100
	Pig, skin & fat	50
	Pig, liver	500
	Pig, kidney	1500

STORAGE CONDITION

Store below 30°C. Protect from direct sunlight.

SHELF LIFE

3 years.

After 1st opening, reconstitution or dilution, use within 24 hours.

PACKING

100gm

500gm

1kg

15kg

THYE PHARMA SDN. BHD.

(manufacturer)

No.2 Lorong Industri Ringan Permatang
Tinggi 8,
Kawasan Industri Ringan Permatang Tinggi,
14000 Bukit Mertajam,
Pulau Pinang.

Revised September 2023