

DEXCHEM INDUSTRIES SDN.BHD (555717-U)

No.5, 11 & 11A, Lorong Cherok Tokun Jaya 2, Kawasan Perindustrian Cherok Tokun,
14000 BUKIT MERTAJAM PULAU PINANG, MALAYSIA

PRODUCT LITERATURE**1. Product Name:****ENFLODEX 200MG/ML SOLUTION****2. Each 1ml solution contains:****Active Ingredient:**

Enrofloxacin 200.00mg

Preservatives:

Benzyl Alcohol 62.5ml

3. Product Description:

Clear solution, pale yellow to off white in colour

4. Target Species:

Poultry (Chickens)

5. Pharmacodynamics:

Mode of action

Two enzymes essential in DNA replication and transcription, DNA gyrase and topoisomerase IV, have been identified as the molecular targets of fluoroquinolones. They modulate the topological state of DNA through cleaving and resealing reactions. Initially, both strands of the DNA double helix are cleaved. Then, a distant segment of DNA is passed through this break before the strands are resealed. Target inhibition is caused by non-covalent binding of fluoroquinolone molecules to an intermediate state in this sequence of reactions, in which DNA is cleaved, but both strands are retained covalently attached to the enzymes. Replication forks and

translational complexes cannot proceed beyond such enzyme-DNA-fluoroquinolone complexes, and inhibition of DNA and mRNA synthesis triggers events resulting in a rapid, drug concentration-dependant killing of pathogenic bacteria.

Antibacterial spectrum

Enrofloxacin is active against many Gram-negative bacteria, against Gram-positive bacteria and *Mycoplasma* spp.

6. Pharmacokinetics:

Enrofloxacin administered via drinking water to poultry is rapidly and very well absorbed.

Concentrations in target tissues like lungs, liver, kidney, intestine and muscle tissue, exceed plasma concentrations by far. In poultry enrofloxacin is poorly metabolised to its active metabolite ciprofloxacin (approximately 5%). Enrofloxacin is eliminated from the body at a half-life of 6 hours. Protein binding in poultry is approximately 25%.

7. Environmental Properties

NA

8. Indication:

Treatment of the infections caused by the following bacteria susceptible to enrofloxacin:

Chickens

Mycoplasma gallisepticum

Mycoplasma synoviae

Avibacterium paragallinarum

Pasteurella multocida

Escherichia coli

Enrofloxacin should be used where clinical experience, supported where possible by sensitivity testing of the causal organism, indicates enrofloxacin as the active substance of choice

9. Recommended Dose:

Chickens:

10 mg enrofloxacin/kg bodyweight per day for 3-5 consecutive days.

(Note: 0.05 ml of ENFLODEX 200MG/ML SOLUTION contains 10 mg enrofloxacin.)

Treatment for 3-5 consecutive days; for 5 consecutive days in mixed infections and chronic progressive forms. If no clinical improvement is achieved within 2-3 days, alternative antimicrobial therapy should be considered based on susceptibility testing.

Via the drinking water. Always make sure that the entire dose offered has been consumed. The medicated water should be made up fresh each day just before it is offered to the animals. The drinking water must be medicated throughout the treatment period, and no other water source should be available. Determine the bodyweight of the birds as accurately as possible in order to avoid underdosing.

Use only fresh pre-solutions, prepared every day before start of treatment. Pumping systems should be checked constantly to assure proper medication. Empty the water system and fill it with medicated water before starting the treatment.

Based on the recommended doses and the number and weight of the animals to be treated, the exact daily amount should be calculated according to the following formula:

Daily amount of ENFLODEX 200MG/ML SOLUTION (ml) per liter of drinking water :

Total number of chicken x Average body weight in kg x 0.05 = Total volume (ml) per day

ENFLODEX 200MG/ML SOLUTION may be put directly into the header tank or introduced via a water proportioner pump

10. Mode of Administration:

Oral

11. Contraindications:

Do not use for prophylaxis

Do not use when resistance/cross-resistance to (fluoro) quinolones is known to occur in the flock intended for treatment

Do not use in the case of known hypersensitivity to the active substance, other (fluoro) quinolones or to any of the excipients.

12. Warning & Precautions:

Special warnings for each target species

Treatment of *Mycoplasma spp* infections may not eradicate the organism.

Special precautions for use

Special precautions for use in animals

Official and local antimicrobial policies should be taken into account when the product is used.

Fluoroquinolones should be reserved for the treatment of clinical conditions which have responded poorly, or are expected to respond poorly, to other classes of antimicrobials. Since enrofloxacin was first authorised for use in poultry, there has been widespread reduction in susceptibility of *E.colito* fluoroquinolones and emergence of resistant organisms.

Whenever possible, fluoroquinolones should only be used based on susceptibility testing.

Use of the product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to the fluoroquinolones and may decrease the effectiveness of treatment with other quinolones due to the potential for cross resistance.

Special precautions to be taken by the person administering the medicinal product to animals

Those with known hypersensitivity to (fluoro)quinolones should avoid contact with this product. Avoid contact with skin and eyes.

Rinse any splashes from skin or eyes immediately with water.

Wash hands and exposed skin after use.

Do not eat, drink or smoke while using the product.

TO BE PRESCRIBED BY REGISTERED VETERINARY SURGEONS
ONLY.

13. Interactions with Other Medicaments

In vitro, an antagonism was shown, when combining fluoroquinolones with bacteriostatic antimicrobial agents such as macrolides or tetracyclines and phenicols. The simultaneous application of substances containing aluminium or magnesium can impair the absorption of enrofloxacin.

14. Pregnancy & Lactation:

Do not use in laying hens producing eggs for human consumption. Do not administer to layer replacement birds within 14 days of coming into lay.

15. Side effects:

None known.

16. Symptoms & Treatment of Overdose:

No adverse clinical symptoms were observed in chickens and turkeys treated respectively with doses up to 10 and 6 times higher than the therapy dose.

The use of fluoroquinolones during the growth phase combined with a marked and prolonged increase in the intake of drinking water, and hence active ingredient, possibly due to high temperatures, may potentially be associated with damage of the articular cartilage.

17. Withdrawal Period:

Chickens: Meat and offal: 7 days

Not authorized for use in birds producing eggs for human consumption.
Do not administer to layer replacement birds within 14 days of coming into lay.

18. Storage Condition:

Store below 30° C in dry place, protected from light.

19. Dosage Form:

Oral Solution

20. Packaging :

Packed in 1L plastic HDPE bottle with screw cap

21. Expiry date:

2 years from date of manufacturing

22. Shelf Life after first opening of container

To be used within 24 hours after first opening of container.

23. Shelf life after reconstitution or dilution

To be used within 24 hours after dilution.

Name & Address of the product registration holder,

DEXCHEM INDUSTRIES SDN.BHD^(555717-U)

No.5, 11 & 11A, Lorong Cherok Tokun Jaya 2, Kawasan Perindustrian Cherok Tokun,
14000 BUKIT MERTAJAM PULAU PINANG, MALAYSIA

Manufactured & Distributed by:

DEXCHEM INDUSTRIES SDN.BHD^(555717-U)

No 6 & 8, Jalan Industri Cherok To`Kun 2, Taman Industri Cherok To`kun,
14000 BUKIT MERTAJAM PULAU PINANG, MALAYSIA

Last update on: 01/09/2018