

OMS TILMICOSIN 200MG/G PREMIX POWDER

Composition

Each g contains:

Tilmicosin Phosphate223mg
(equivalent to Tilmicosin 200mg)

Product Description

OMS Tilmicosin 200mg/g Premix Powder is a creamy to beige powder.

Pharmacodynamics

Tilmicosin is a semi-synthetic antibiotic of the macrolide group, and is believed to affect protein synthesis. It has bacteriostatic action but at high concentrations it may be bactericidal. This antibacterial activity is predominantly against Gram-positive microorganism with activity against certain gram-negative ones and Mycoplasma of a bovine, porcine, ovine and avian origin. In particular its activity has been demonstrated against the following micro-organism:

Pigs: Mycoplasma hyopneumoniae, Pasteurella multocida, Actinobacillus pleuropneumoniae.

Scientific evidence suggests that macrolides act synergistically with the host immune system. Macrolides appear to enhance phagocyte killing of bacteria. Tilmicosin has been shown to inhibit *in vitro* the replication of the Porcine Reproductive and Respiratory Syndrome virus in alveolar macrophages in a dose dependent fashion.

Cross resistance between tilmicosin and other macrolides and lincomycin has been observed.

Pharmacokinetics

Absorption: Tilmicosin moves rapidly out of the serum into areas of low pH.

Distribution: Following oral administration, tilmicosin is distributed throughout the body with especially high levels found in the lung and in lung tissue macrophages. It is also distributed in the liver and kidney tissues.

Biotransformation: Several metabolites are formed, the predominant one being identified as T1. However the bulk of tilmicosin is excreted unchanged.

Elimination: Following oral administration, tilmicosin is excreted mainly via the bile into the faeces, but a small proportion is excreted via the urine.

Indication

Pigs: Treatment of respiratory disease caused by *Actinobacillus pleuropneumoniae*, *Mycoplasma hyopneumoniae*, *Pasteurella multocida* and other organisms sensitive to tilmicosin.

Mode of Administration

Orally. Mixed into feed.

Recommended Dosage

Orally. Mixed into feed.

Indication	Dose of tilmicosin	Duration of treatment	Inclusion rate in feed
Pigs: Treatment of respiratory disease	8-16 mg/kg bodyweight /day (equivalent to 200 to 400 ppm in the feed)	15-21 days	1-2 kg OMS Tilmicosin Premix/tonne

Contraindications

Horses or other Equidae, must not be allowed access to feeds containing tilmicosin. Horses fed with tilmicosin medicated feeds may present signs of toxicity with lethargy, anorexia, reduction of feed consumption, loose stools, colic, distension of the abdomen and death.

Do not use in case of hypersensitivity to tilmicosin or to any of the excipients.

Warnings and Precautions

Special warnings for each target species

Under practical conditions, the management of respiratory disease outbreaks recognises that acutely ill animals are inappetent and require parenteral therapy.

Special precautions for use

i) Special precautions for use in animals

Inappropriate use of the product may increase the prevalence of bacteria resistant to tilmicosin and may decrease the effectiveness of treatment with tilmicosin related substances.

Official, national and regional antimicrobial policies should be taken into account when the product is used.

Due to the likely variability (time, geographical) in the occurrence of the resistance of bacteria for tilmicosin, bacteriological sampling and susceptibility testing are recommended

ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals

Tilmicosin may induce irritation. Macrolides, such as tilmicosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tilmicosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided.

To avoid exposure during preparation of the medicated feed, wear overalls, safety glasses, impervious gloves and wear either a disposable half mask respirator. Do not eat, drink or smoke when handling this product. Wash hands after use.

In the case of accidental ingestion, wash out mouth immediately with water and seek medical advice. In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water.

Do not handle the product if you are allergic to ingredients in the product.

If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

Interactions with Other Medicaments

None known.

Pregnancy and Lactation

The safety of tilmicosin has not been established in boars used for breeding purposes.

Adverse Effects/Undesirable Effects

In very rare cases, feed intake may decrease (including feed refusal) in animals receiving medicated feed. This effect is transient.

Overdose and Treatment

No symptoms of overdose have been seen in pigs fed a ration containing levels of tilmicosin up to 80 mg/kg bodyweight (equivalent to 2000 ppm in the feed or ten times the recommended dose) for 15 days.

Withdrawal Period(s)

21 days

Storage condition

Store in a cool dry place, protected from sunlight. Do not store above 30°C.
Keep out of reach of children/Jauhkan dari kanak-kanak.

Shelf life

Shelf life as packaged for sale: 12 months

Shelf life after reconstitution into feed: 24 hours

Pack size(s)

Paper bag of 10kg

Product Registration Holder/ Manufacturer

Osmosis Nutrition Sdn. Bhd. (594876A)

Lot 16014(P.T. 24341), Jalan Nilam 3,

Bandar Nilai Utama,

71800 Nilai, Negeri Sembilan, Malaysia.

Tel: +606-799 9090 Fax: +606-799 9155

Date of revision of Package Insert

20/05/2024