

148 mm
75 mm



Product Description

Baycox 5% w/v Oral Suspension is a white to yellowish suspension. 1 ml of Baycox 5% w/v Oral Suspension contains 50 mg of toltrazuril as active ingredient, 2.1 mg of sodium propionate and 2.1 mg of sodium benzoate as preservatives.

Pharmacodynamics

Toltrazuril is a triazinone derivative. It acts against coccidia of the genus *Isospora* and *Eimeria*. It is acting against all intracellular development stages of coccidia of the merogony (asexual multiplication) and gamogony (sexual phase). All stages are destroyed, thus the mode of action is coccidiocidal.

Pharmacokinetics

Piglets: After oral administration toltrazuril is slowly absorbed with a bioavailability of $\geq 70\%$. The main metabolite is characterised as toltrazuril sulfone. The elimination of toltrazuril is slow with a half-life elimination time around 3 days. The major route of excretion is via the faeces.

Calves: After oral administration in cattle toltrazuril is slowly absorbed. The maximal plasma concentration ($C_{max} = 36.6$ mg/l) was observed between 24 and 48 hours (geometric mean 33.9 hours) following oral administration. The elimination of toltrazuril is slow with a terminal half-life time of approximately 2.5 days (64.2 hours). The main metabolite is characterised as toltrazuril sulfone. The major route of excretion is via the faeces.

Indication

Piglets: For the prevention of clinical signs of coccidiosis in neonatal piglets on farms with a confirmed history of coccidiosis caused by *Isospora suis*.

Calves: For the prevention of clinical signs of coccidiosis and reduction of coccidia shedding in housed calves replacing cows producing milk for human consumption (dairy cows) on farms with a confirmed history of coccidiosis caused by *Eimeria bovis* or *Eimeria zuernii*.

Recommended Dose

Piglets

Individual animal treatment. Each pig to be treated on day 3-5 of life with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight. Due to the small volumes required to treat individual piglets, use of dosing equipment with a dose accuracy of 0.1 ml is recommended. Treatment during an outbreak will be of limited value for the individual piglet because of damage to the small intestine having already occurred.

Calves

Each animal should be treated with a single oral dose of 15 mg toltrazuril/kg body weight corresponding to 3.0 ml oral suspension per 10 kg body weight. For the treatment of a group of animals of the same breed and same or similar age, the dosing should be done according to the heaviest animal of this group. To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.

Method of administration

The ready-to-use oral suspension must be shaken before use.

90208685

Contraindications

For environmental reasons:
Do not use in calves weighing more than 80 kg bodyweight. Do not use in fattening units such as veal or beef calves. For more details see sections Warning and Precautions.

Warning and Precautions

Special Warnings

As with any antiparasiticide frequent and repeated use of antiprotazoals from the same class may lead to the development of resistance. It is recommended to treat all animals in a pen. Hygienic measures may reduce the risk of coccidiosis. It is therefore, recommended to improve concomitantly the hygienic conditions in the concerned facility, particularly dryness and cleanliness.

To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.

To alter the course of an established clinical coccidial infection, in individual animals already showing signs of diarrhoea, additional supportive therapy may be required.

Special precautions to be taken by the person administering the veterinary medicinal product to animals
Wash any splashes from skin or eyes immediately with water.

Other Precautions

The major metabolite of toltrazuril, toltrazuril sulfone (ponazuril), has been shown to be both persistent (half-life >1 year) and mobile in soil and to be toxic to plants.

For environmental reasons:
In order to prevent any adverse effects on plants and possible contamination of groundwater manure from treated calves must not be spread onto land without dilution with manure from untreated cows. Manure from treated calves must be diluted with at least 3 times the weight of manure from mature cows before it can be spread onto land.

TO BE PRESCRIBED AND TREATED WITH BY REGISTERED VETERINARY SURGEONS ONLY.

Environmental properties

The metabolite of toltrazuril, toltrazuril sulfone (ponazuril) is a persistent (half-life >1 year) and mobile compound and has adverse effects on both the growth and emergence of plants. Given the persistent properties of ponazuril repeated spreading of manure from treated animals may lead to an accumulation in the soil and consequently a risk to plants. The accumulation of ponazuril in soil together with its mobility also leads to a risk of leaching to groundwater.

Interactions with Other Medicaments

None known

Pregnancy and Lactation

Laboratory studies in rat and rabbit have not produced any evidence of a teratogenic, foetotoxic or maternotoxic effect.

Adverse Reactions

None known.

Symptoms and Treatment of Overdose

A threefold overdose is well tolerated by healthy piglets and calves without signs of intolerance.

Withdrawal Period

Piglets: Meat and offal: 77 days
Calves: Meat and offal: 63 days

Not permitted for use in lactating animals producing milk for human consumption.

Storage Condition

Store in a dry place below 30°C.

The shelf life of this product after first opening of the container is 8 weeks.

Dosage Forms and packaging available

Baycox 5% w/v Oral Suspension is available in 100 ml bottle.

Marketing Authorisation Holder:
Elanco Malaysia Sdn Bhd, Unit 5.04, Level 5 & 6, Tower Block, The Bousteador, No. 10, Jalan PJU 7/6, Mutiara Damansara, 47800 Petaling Jaya, Selangor, Malaysia

Manufactured by:
KVP Pharma + Veterinär Produkte GmbH
Projensdorfer Straße 324, 24106 Kiel, Germany.

Date of revision: April 2022

Baycox is sold by Elanco or its affiliates and is not a product of Bayer.
The Product Name Baycox is owned by Bayer and used under license.
© 2022 Elanco or its affiliates

210 mm
Pharmaco-
Direction

• Item code placement is flexible
and can be moved to suit artwork

PRODUCT INFO

BLUE #:	502166APK	Product Name:	Baycox 5% w/v Oral Suspension
Item Code:	90208685	Component:	Package Inserts and Leaflets
Product Code:	LBAH	Pack Size:	100 ml
Previous Item Code	90204325		

ARTWORK INFO

Template:	PI_148x210mm_Kiel_v2a	Packaging Spec(s):	TPM_016192
Barcodes/Type:	Laetus: 29	Add. Info:	N/A
		Minimum Core Data Point Size:	6pt