



MINYAK ANGIN ZHUI FONG YEW 千里 追风油

Registration No. : MAL

外观 : 带有特征气味的褐色药油。
主治 : 缓解由风湿引起的肌肉和关节疼痛。
用法 : 将此药油搽于患处，每天3至4次。
给药途径 : 外用

Product Description : Brown-coloured oil with characteristic odour .
Indication : To relieve muscle and joint pain related to rheumatic pain .
Direction for use : Rub on site of pain 3 to 4 times daily .
Route of Administration : Topical

Pharmacodynamics : Menthol dilates the blood vessels causing a sensation of coldness , followed by analgesic effect . Menthol also acts as a penetration enhancer , increasing the penetration of drugs when applied on the skin , to give a faster onset of action . Methyl Salicylate is a salicylic acid derivative . Salicylates inhibit cyclooxygenase , thereby reducing the formation of prostaglandins , and cause platelet dysfunction . Methyl salicylate is used topically as a counter - irritant . Upon application , it is absorbed through the skin and is applied for the relief of pain in rheumatic conditions and painful muscle or joints .

Pharmacokinetics : After absorption , menthol is excreted in the urine and bile as a glucuronide . Methyl Salicylate may be absorbed through intact skin . The absorption of topical salicylates is proportional to the surface area involved , duration of exposure , concentration and skin integrity . Absorption characteristics of salicylates vary with the dose , formulation and route of administration . Percutaneous absorption is enhanced by exercise , heat , occlusion or disruption of the integrity of the skin or application to large areas of skin . Both the rate and extent of absorption increases after repeated application ; increasing the bioavailability . Methyl salicylate is extensively metabolized to salicylic acid in the dermal and subcutaneous tissues after topical application . At therapeutic levels , the half - life of salicylates is 2 - 4 hours . As salicylate level increase into the toxic range , the half - life can be greater than 18 hours .

Contraindications : Patient with known hypersensitivity to any of the ingredients in the product .

Warnings and Precautions : This product contains methyl salicylate and when applied or rub on the skin , can be absorbed through the skin into blood . For patients taking warfarin , excessive application on to the skin for muscle or joint pain may increase the chances of bleeding . Avoid contact with the eyes and mucous membrane . Do not apply to open wounds or damaged skin . Avoid excessive use during pregnancy and lactation . Do not apply over large areas of the body or bandage or use with heating pads . Initially apply a small amount to a limited skin area and observe whether there is discomfort or itch before applying to a larger skin area . Discontinue use if irritation develops . For external use only . Not to be taken .

Effects on Ability to Drive and Use Machines : Not known

Interactions with Other Medicaments : This product contains methyl salicylate and when applied or rubbed on to the skin , can be absorbed through the skin into the blood . For patients taking warfarin , excessive application on to the skin for muscle or joints pains may increase the chances of bleeding .

Statement on Usage During Pregnancy and Lactation : Safety for use in pregnancy has not been established; therefore the potential benefit of the product should be weighed against the possible risks to the mother and child .

SIDE EFFECT / ADVERSE REACTION : Skin irritation , redness or swelling , burning or stinging sensation . Adverse reaction is rare . Symptoms of dizziness and nausea may occur .

Symptoms and Treatment of Overdose : Mild chronic salicylate intoxication or salicylism usually occurs only after repeated use of large doses . Salicylism can also occur following excessive topical application of salicylates . Symptoms include dizziness , tinnitus , deafness , sweating , nausea and vomiting , headache and confusion and may be controlled by reducing the dosage .

药效学：请参阅英文版本。 药动学：请参阅英文版本。 禁忌：对此产品中任何成份过敏者。

警告和注意事项：

此产品含有水杨酸甲酯，当涂抹或擦在皮肤时，可以通过皮肤渗透进入血液。对于服用华法林的患者，当因肌肉或关节疼痛时而过度涂抹于皮肤时可能会增加出血的可能性。避免与眼睛和粘膜接触。不要涂抹于伤口或受损皮肤。避免在怀孕和哺乳期间过度使用。不要大面积涂抹在身体或用绷带包扎或使用加热垫。在涂抹于较大区域的皮肤前，涂抹少量在有限的皮肤区域以观察是否有不适或瘙痒。若刺激反应持续，请停止使用。外用忌食。

对驱动和使用机器的能力的影响：未知 与其他药物的相互作用：请参阅英文版本。 妊娠和哺乳期间使用声明：请参阅英文版本。

侧面效应/不良反应：皮肤刺激，发红或肿胀，灼热或刺痛感。不良反应很少见。可能会出现头晕和恶心的症状。过量症状和治疗：请参阅英文版本。

Storage Conditions : Store below 30°C .
Keep out of reach of children . Jauhkan daripada kanak - kanak .

Packing Size : 60ml

Shelf Life : 3 years from date of manufacture .

贮藏：

储存于30°C以下。放在儿童不能接触的地方。

保质期：自生产日期起3年。

Revision Date : 17 / 4 / 2017

Manufactured by :

WELLMETECH PHARMACEUTICAL INDUSTRIES SDN. BHD.

Lot 8058 , Kota Samarahan Industrial Park , 94300 Samarahan , Sarawak , Malaysia .

Distributed by :

EYH FONG TRADING

No . 10 , Hala Rapat Baru 21 , Kawasan Perindustrian Ringan Kinta Jaya , 31350 Ipoh , Perak .

Product Registration Holder :

CAP SEEN SHEI HERBS ENTERPRISE

No . 10 , Hala Rapat Baru 21 , Kawasan Perindustrian Ringan Kinta Jaya , 31350 Ipoh , Perak .