

NAME OF THE VETERINARY MEDICINAL PRODUCT

Previcox 57 mg chewable tablets for dogs
firocoxib

Manufacturer :

Boehringer Ingelheim Animal Health France, 4 Chemin du Calquet, 31000 Toulouse, France

Registration Holder and Repacker :

Rhone Ma Malaysia Sdn Bhd (200001023279), Lot 18A & 18B, Jalan 241, Seksyen 51A, 46100 Petaling Jaya, Selangor

STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Each chewable tablet contains:

Active substance:
firocoxib: 57 mg

Excipients:

Iron oxides (E172)
Caramel (E150d)

PRODUCT DESCRIPTION

Tan-brown, round, convex, tablets with a cross-shaped break line on one side. The tablets can be divided into 2 or 4 equal parts.

PHARMACODYNAMICS

Firocoxib is a non-steroidal anti-inflammatory drug (NSAID) belonging to the Coxib group, which acts by selective inhibition of cyclooxygenase-2 (COX-2)-mediated prostaglandin synthesis. Cyclooxygenase is responsible for generation of prostaglandins. COX-2 is the isoform of the enzyme that has been shown to be induced by pro-inflammatory stimuli and has been postulated to be primarily responsible for the synthesis of prostanoid mediators of pain, inflammation, and fever. Coxibs therefore display analgesic, anti-inflammatory and antipyretic properties. COX-2 is also thought to be involved in ovulation, implantation and closure of the *ductus arteriosus*, and central nervous system functions (fever induction, pain perception and cognitive function). In *in-vitro* canine whole blood assays, firocoxib exhibit approximately 380-fold selectivity for COX-2 over COX-1. The concentration of firocoxib required to inhibit 50% of the COX-2 enzyme (i.e. the IC₅₀) is 0.16 (±0.05) µM, whereas the IC₅₀ for COX-1 is 56(±7) µM.

PHARMACOKINETICS

Following oral administration in dogs at the recommended dose of 5mg per kg of bodyweight, firocoxib is rapidly absorbed and the time to maximal concentration (T_{max}) is 1.25 (±0.85) hours. The peak concentration (C_{max}) is 0.52 (±0.22) µg/ml (equivalent to approximately 1.5 µM), area under the curve (AUC 0-24) is 4.63 (±1.91) µg xhr/mL and oral bioavailability is 36.9 (±20.4) percent. The elimination half-life (t_{1/2}) is 7.59 (±1.53) hours. Firocoxib is approximately 96% bound to plasma proteins. Following multiple oral administrations, the steady state is reached by the third daily dose. Firocoxib is metabolised predominantly by dealkylation and glucuronidation in the liver. Elimination is principally in the bile and gastrointestinal tract.

INDICATIONS

For the relief of pain and inflammation associated with osteoarthritis in dogs.

For the relief of post-operative pain and inflammation associated with soft-tissue, orthopaedic and dental surgery in dogs.

DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Oral use. To be supplied only on veterinary prescription.

Osteoarthritis :


Administer 5 mg per kg bodyweight once daily as presented in the table below. Tablets can be administered with or without food. Duration of treatment will be dependent on the response observed. As field studies were limited to 90 days, longer term treatment should be considered carefully and regular monitoring undertaken by the veterinarian.

Relief of post-operative pain :

Administer 5 mg per kg bodyweight once daily as presented in the table below for up to 3 days as needed, starting approximately 2 hours prior to surgery. Following orthopaedic surgery and depending on the response observed, treatment using the same daily dosing schedule may be continued after the first 3 days, upon judgement of the attending veterinarian.

FOR ORAL USE AS PER TABLE BELOW.		
Body weight (kg)	Number of chewable tablets by size	
	57 mg	
3.0 – 5.5	0.5	5.2 – 9.5
5.6 – 7.5	0.75	5.7 – 7.6
7.6 – 10	1	5.7 – 7.5
10.1 – 13	1.25	5.5 – 7.1
13.1 – 16	1.5	5.3 – 6.5
16.1 – 18.5	1.75	5.4 – 6.2

Tablets can be divided into 2 or 4 equal parts to enable accurate dosing.



TARGET SPECIES

Dogs.

CONTRAINDICATIONS

Do not use in pregnant or lactating bitches.

Do not use in animals less than 10 weeks of age or less than 3kg bodyweight.

Do not use in animals suffering from gastrointestinal bleeding, blood dyscrasia or haemorrhagic disorders.

Do not use concomitantly with corticosteroids or other non-steroidal anti-inflammatory drugs (NSAID).

ADVERSE REACTIONS

Emesis and diarrhoea have occasionally been reported. These reactions are generally of a transitory nature and are reversible when the treatment is stopped. Renal and/or hepatic disorders have been reported in very rare cases in dogs administered the recommended treatment dose. Rarely, nervous system disorders have been reported in treated dogs. If adverse reactions like vomiting, repeated diarrhoea, faecal occult blood, sudden weight loss, anorexia, lethargy, degradation of renal or hepatic biochemistry parameters occur, use of the product should be stopped and the advice of a veterinarian should be sought. As with other NSAIDs, serious adverse effects can occur and in very rare cases, may be fatal.

WITHDRAWAL PERIOD

Not applicable.

SPECIAL PRECAUTIONS

Special precautions for use in animals:

The recommended dose, as indicated in the dosing table, should not be exceeded.

Use in very young animals, or animals with suspected or confirmed impairment of renal, cardiac or hepatic function may involve additional risk. If such use cannot be avoided, those dogs require careful veterinary monitoring.

Avoid use in any dehydrated, hypovolaemic or hypotensive animals, as there is a potential risk of increased renal toxicity. Concurrent administration of potentially nephrotoxic drugs should be avoided.

Use this product under strict veterinary monitoring where there is a risk of gastrointestinal bleeding, or if the animal previously displayed intolerance to NSAIDs. Renal and/or hepatic disorders have been reported in very rare cases in dogs administered the recommended treatment dose. It is possible that a proportion of such cases had sub-clinical renal or hepatic disease prior to the commencement of therapy. Therefore, appropriate laboratory testing to establish baseline renal or hepatic biochemistry parameters is recommended prior to and periodically during administration.

The treatment should be discontinued if any of these signs are observed: repeated diarrhoea, vomiting, faecal occult blood, sudden weight loss, anorexia, lethargy, degradation of renal or hepatic biochemistry parameters.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Wash hands after use of the product.

In case of accidental ingestion, seek medical advice immediately and show the package insert or the label to the physician.

Divided tablets should be returned to the original package.

Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

INTERACTIONS WITH OTHER MEDICAMENTS

Pre-treatment with other anti-inflammatory substances may result in additional or increased adverse effects and accordingly a treatment-free period with such drugs should be observed for at least 24 hours before the commencement of treatment with Previcox. The treatment-free period, however, should take into account the pharmacokinetic properties of the products used previously.

Previcox must not be administered in conjunction with other NSAIDs or glucocorticosteroids. Gastrointestinal tract ulceration may be exacerbated by corticosteroids in animals given non-steroidal anti-inflammatory drugs.

Concomitant treatment with molecules displaying action on renal flow, e.g. diuretics or Angiotensin Converting Enzyme (ACE) inhibitors, should be subject to clinical monitoring. Concurrent administration of potentially nephrotoxic drugs should be avoided as there might be an increased risk for renal toxicity. As anaesthetic drugs may affect renal perfusion, the use of parenteral fluid therapy during surgery should be considered to decrease potential renal complications when using NSAIDs peri-operatively.

Concurrent use of other active substances that have a high degree of protein binding may compete with firocoxib for binding and thus lead to toxic effects.

OVERDOSE AND TREATMENT

In dogs ten weeks of age, at the start of treatment, at dose rates equal or greater to 25 mg/kg/day (5 times the recommended dose) for three months, the following signs of toxicity were observed: bodyweight loss, poor appetite, changes in the liver (accumulation of lipid), brain (vacuolisation), duodenum (ulcers) and death. At dose rates equal or greater to 15 mg/kg/day (3 times the recommended dose) for six months, similar clinical signs were observed, albeit that the severity and frequency were less and duodenal ulcers were absent.

In those target animal safety studies, clinical signs of toxicity were reversible in some dogs following cessation of therapy.

In dogs seven months of age, at the start of treatment, at dose rates greater than or equal to 25 mg/kg/day (5 times the recommended dose) for six months,

gastrointestinal adverse effects, i.e. vomiting were observed.

Overdose studies were not conducted in animals over 14 months of age. If clinical signs of overdosing are observed, discontinue treatment.

PREGNANCY AND LACTATION

Do not use in pregnant or lactating bitches. Laboratory studies in rabbits have shown evidence of maternotoxic and faetotoxic effects at dose rates approximating the recommended treatment dose for the dog.

SPECIAL STORAGE PRECAUTIONS

Do not store above 30°C.

In order to protect from moisture, store in the original package.

Keep out of the reach of children.

Divided tablet may be stored for up to 1 month in the original package.

PACKING

57mg :

1 carton box containing three blister of 10 tablets.

1 bottle with 60 tablets.

FOR VETERINARY USE ONLY

Date of Revision : July 2021

5L PRE573-030 03/XXX



Boehringer
Ingelheim

LABEL CODE :

5LPRE573-030

PRODUCT : PREVICOX 57MG CHEWABLE TABLET FOR DOGS

PACK SIZE : 30 TABLETS, 60 TABLETS

COUNTRY : MY

DIMENSIONS :
cm : 17.0 x 55.0

VERSION

Version : 3

MMM/YY :

COLORS

Noir/Black

Please use the official PANTONE® (P) matching system for an accurate color representation.

Marketing approval:

COMPLETE NAME:

DATE + SIGNATURE:

Regulatory approval:

COMPLETE NAME:

DATE + SIGNATURE:

NOT PRINTABLE
NON IMPRIMABLE

TECHNICAL NOTE(S) - INFO(S) TECHNIQUE

- Artwork identification(s)
- Text areas / zones de textes
- Dieline / Découpe