

PF SULFAMONOMETHOXINE PREMIX 50% W/W POWDER

Composition:

Sulfamonomethoxine Sodium.....500mg

Exp. Qs.....1g

Description

White Powder

Pharmacodynamic

Bacteriostatic. Sulfonamides interfere with the biosynthesis of folic acid in bacterial cells; they compete with para-aminobenzoic acid (PABA) for incorporation in the folic acid molecule. By replacing the PABA molecule and preventing the folic acid formation required for DNA synthesis, the sulfonamides prevent multiplication of the bacterial cell. Only organisms that synthesize their own folic acid are susceptible; mammalian cells use preformed folic acid and, therefore, are not susceptible. Cells that produce excess PABA or environments with PABA, such as necrotic tissues, allow for resistance by competition with the sulfonamide.

Pharmacokinetic

Most sulfonamides are well absorbed. Delays in absorption may occur in adult ruminants or when sulfonamides are administered with food to monogastric animals. Sulfonamides are widely distributed throughout the body. They cross the placenta, and a few penetrate into the cerebrospinal fluid. Sulfonamides are primarily metabolized in the liver but metabolism also occurs in other tissues. Renal excretion is the primary route of elimination for most nonenteric sulfonamides.

Indication

Chickens: infectious coryza, coccidiosis, , leucocytozoonosis

Route of Administration

Oral

Recommended dose

Administer orally by mixing uniformly the following amounts as sulfamonomethoxine sodium per ton of feed.

Chickens (except laying hens):

Infectious Coryza 1,000 g (2,000g PF Sulfamonomethoxine Premix 50%)

Coccidiosis Disease 500 - 1,000 g (1,000 - 2,000g PF Sulfamonomethoxine Premix 50%)

Leucocytozoonosis disease 10 - 50 g (20 - 100g PF Sulfamonomethoxine Premix 50%)

Contraindication

Hypersensitivity to sulfonamides.

Interaction with other medicinal products

Drug interactions relating specifically to the use of sulfonamides in animals are rarely reported.

Pregnancy and lactation

Do not use for laying hen.

Side effects

Crystallization of sulfonamides can occur in the kidneys or urine with high doses of sulfonamide or when an animal is dehydrated.

Symptoms and treatment of overdose

Toxicities secondary to acute overdose of sulfonamides are not typically reported.

Storage

Store in a cool dry place (<30°C). Strictly avoid light and heat exposure.

Warning and Precaution

Keep out of reach of children / Jauhi ubat dari kanak-kanak (General)

- Because this drug is indication drug, please use it by prescription / instruction of veterinarian.
- This product should be used only for treatment of indications as stated for efficacy / effect.
- Please adhere strictly to prescribed dosage when using this drug.
- When using this drug, make sure to stop administration after the minimum period necessary for treatment of indication, do not do administer continuously for over weeks.
- This product should be used according to the use standard.
- If you use this drug for Leucocytozoonosis disease or atrophic rhinitis (AR), please obtain guidance from a veterinarian over the details of the use.

(Attention to users)

- In case of accidentally swallowing this drug, please get medical attention immediately.
- When mixing with feed, etc., do wear mask and take care not to inhale any dust.

Shelf life

36 months

Shelf life after opening

24 hours

Shelf life after dilution or constitution

24 hours

Withdrawal period

Chickens (except egg laying chicken): 7 days before slaughter to serve for food

Packaging

25kg

Revised date

14/8/2018

Product Registration Holder:

PANFAST MARKETING (M) SDN.BHD.

PENANG SCIENCE PARK 1102, MUKIM 13, LORONG PERINDUSTRIAN BUKIT

MINYAK 18, KAWASAN PERINDUSTRIAN BUKIT MINYAK,

14100 SIMPANG AMPAT, PENANG, MALAYSIA

TEL: +604-5016868 FAX: +604-5016866

Manufactured by:

SHENNONG ANIMAL HEALTH (M) SDN.BHD.

No.6, 8, 10, Jalan Industri USJ 1/19, Taman Perindustrian USJ 1

47610 SUBANG JAYA, Selangor Darul Ehsan, Malaysia.

Tel: 03-8011 4646 Fax : 03-8024 1166