

SN ANTI-COCCI 45 MG/ML

ORAL SOLUTION

(FOR VETERINARY USE ONLY)

COMPOSITION

Each 1000 ml contents:

Sulfaquinoxaline	45 g
Benzyl alcohol	18.5 ml

PRODUCT DESCRIPTION

This product is a yellowish color liquid.

PHARMACODYNAMICS

Sulfonamides are usually bacteriostatic agents when used alone. Sulfonamides interfere with the biosynthesis of folic acid in bacterial cells; they compete with para-aminobenzoic acid (PABA) for incorporation in the folic acid molecule. By replacing the PABA molecule and preventing the folic acid formation required for DNA synthesis, the sulfonamides prevent multiplication of the bacterial cell. Only organisms that synthesize their own folic acid are susceptible; mammalian cells use preformed folic acid and, therefore, are not susceptible. Cells that produce excess PABA or environments with PABA, such as necrotic tissues, allow for resistance by competition with the sulfonamide.

PHARMACOKINETICS

Most sulfonamides are well absorbed orally with the exception of the enteric sulfonamides, such as sulfaquinoxaline, which are minimally absorbed. Delays in absorption may occur in adult ruminants or when sulfonamides are administered with food to monogastric animals.

Sulfonamides are widely distributed throughout the body. They cross the placenta, and a few penetrate into the cerebrospinal fluid. Sulfonamides may be distributed into milk; however, they vary greatly in their ability to do so. The process depends on several factors, including protein binding and pKa values.

Sulfonamides are primarily metabolized in the liver but metabolism also occurs in other tissues. Biotransformation occurs mainly by acetylation, glucuronide conjugation, and aromatic hydroxylation in many species. The types of metabolites formed and the amount of each varies depending on the specific sulfonamide administered; the species, age, diet, and environment of the animal; the presence of disease; and, with the exception of pigs and ruminants, even the sex of the animal.

TARGET SPECIES

Chickens, Turkeys, Cattle and Calves

INDICATIONS

Cattle and calves:

For the control and treatment of outbreaks of coccidiosis caused by *Eimeria bovis* or *E. zurnii*.

Turkeys:

As an aid in the control of acute fowl cholera caused by *Pasteurella multocida* susceptible to sulfaquinoxaline and fowl typhoid caused by *Salmonella gallinarum* susceptible to sulfaquinoxaline.

As an aid in the control of outbreaks of coccidiosis caused by *Eimeria meleagridis* and *E. adenoides*.

Chickens:

As an aid in the control of acute fowl cholera caused by *Pasteurella multocida* susceptible to sulfaquinoxaline and fowl typhoid caused by *Salmonella gallinarum* susceptible to sulfaquinoxaline. As an aid in the control of outbreaks of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, and *E. brunetti*.

ADMINISTRATION & DOSAGE

Oral administration through drinking water. Prepare medicated drinking water fresh daily.

For control of Acute Fowl Cholera / Fowl Typhoid:

Chickens & Turkeys: Administer at the 0.04% level in drinking water (approximately 4.4 L product/ 500 L drinking water) for 2-3 days. Move birds to clean ground. If disease recurs, repeat treatment. Poultry that have survived fowl typhoid outbreaks should not be kept for laying house replacements or breeders unless tests show they are not carriers.

Medicated chickens must actually consume enough medicated water which provides a recommended dosage of approximately 22 to 100 mg per kg per day.

Medicated turkeys must actually consume enough medicated water which provides a recommended dosage of approximately 7.8 to 122 mg per kg per day.

For the control of outbreaks of Coccidiosis:

Chickens: Administer at the 0.04% level in drinking water (approximately 4.4 L product/ 500 L drinking water) for 2-3 days, then the repeat treatment as a 0.025% solution (approximately 2.8 L product/ 500 L drinking water) for 2 to 4 more days. The schedule may be repeated, if necessary. Medicated chickens must actually consume enough medicated water which provides a recommended dosage of approximately 22 to 100 mg per kg per day.

Turkeys: Administer at the 0.025% level in drinking water (approximately 2.8 L product/ 500 L drinking water) for 2 days - skip 3 days - give for 2 days - skip 3 days and give 2 more days. Repeat if necessary. Medicated turkeys must actually consume enough medicated water which provides a recommended dosage of approximately 7.8 to 122 mg per kg per day.

For the control and treatment of outbreaks of coccidiosis:

Cattle and calves: Administer at the 0.015% level in drinking water (approximately 1.7 L product/ 500 L drinking water) for 3-5 days. Medicated cattle and calves must actually consume enough medicated water which provides a recommended dosage of approximately 13 mg per kg per day in cattle and calves.

INTERACTION WITH OTHER MEDICAMENTS

Drug interaction relating specifically to the use of sulfonamides in animals are rarely reported in veterinary literature.

WITHDRAWAL PERIODS

- Meat: 10 days before slaughter.
- Do not medicate chickens or turkeys producing eggs for human consumption.

- Do not use in calves to be processed for veal or lactating dairy cattle.

WARNING AND PRECAUTIONS

- Do not give flushing mashes.
- May cause toxic reactions unless the drug is evenly mixed in water at dosages indicated and used according to directions.
- People who handle this medication should avoid contact with eyes, skin, or clothing to prevent eye and skin burns. In case of contact, the areas affected should be flushed for at least fifteen minutes; medical attention should be sought for eye exposure.
- Keep medicine out of reach of children.
- Jauhkan ubat dari kanak-kanak.

CONTRAINDICATIONS

Hypersensitive to sulfas, severe renal or hepatic impairment.

PREGNANCY AND LACTATION

This product must not be used on laying birds and lactating dairy cattle. Sulfonamides cross the placenta in pregnant animals. Some teratogenic effects have been seen when very high doses were given to pregnant mice and rats.

SIDE EFFECT

Prolonged administration of sulfaquinoxaline may result in depressed feed intake, deposition of crystals in the kidney, or interference with normal blood clotting.

Hemorrhagic syndrome (anorexia, epistaxis, hemoptysis, lethargy, pale mucous membranes, possibly death) has been reported in chickens but may occur in other species.

Sulfaquinoxaline is a vitamin K antagonist that inhibits vitamin K epoxide and vitamin K quinone reductase and causes an effect similar to that of coumarin anticoagulants.

OVERDOSE AND TREATMENT

Toxicities secondary to acute overdose of sulfonamides are not typically reported. Side effects may be more likely to occur with high doses and long-term administration, but are seen at recommended doses as well.

SHELF LIFE: 36 months

SHELF LIFE after first opening: 24 hours

SHELF LIFE after dilution: 24 hours

STORAGE

Store at temperature not exceeding 30°C. Strictly avoid light & heat exposure.

PACKAGING

The product filled 1000 ml in HDPE plastic bottle.

DATE OF REVISION: 14 May 2019



Product Registration Holder and Manufactured By:

SHENNONG ANIMAL HEALTH (M) SDN BHD

神農動物保健(馬)有限公司

No. 2, 4, 6, 8 & 10, Jalan Industri USJ 1/19,

Taman Perindustrian USJ 1,

47610 Subang Jaya, Selangor.

Tel: 03-8011 4646

Fax: 03-8024 1166